
Section 2.4

Solution Sets

Plan For Today

Today we will learn to describe and draw the solution set of an arbitrary system of linear equations Ax = b, using spans.



Recall: the solution set is the collection of all vectors x such that Ax = b is true.

Last time we discussed the set of vectors b for which Ax = b has a solution.

We also described this set using spans, but it was a different problem.

Everything is easier when b = 0, so we start with this case.

Definition


A system of linear equations of the form Ax = 0 is called **homogeneous.**

These are linear equations where everything to the right of the = is zero. The opposite is:

Definition

A system of linear equations of the form Ax = b with $b \neq 0$ is called **inhomogeneous.**

A homogeneous system always has the solution x=0. This is called the **trivial solution**. The nonzero solutions are called **nontrivial**.



Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 3 & 4 \\ 2 & -1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$$
?

We know how to do this: first form an augmented matrix and row reduce.

$$\begin{pmatrix} 1 & 3 & 4 & | & 0 \\ 2 & -1 & 2 & | & 0 \\ 1 & 0 & 1 & | & 0 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 1 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{pmatrix}.$$

The only solution is the trivial solution x = 0.

Observation

Since the last column (everything to the right of the =) was zero to begin, it will always stay zero! So it's not really necessary to write augmented matrices in the homogeneous case.

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -3 \\ 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equation}} \qquad x_1 - 3x_2 = 0$$

$$\xrightarrow{\text{parametric form}} \qquad \begin{cases} x_1 = 3x_2 \\ x_2 = x_2 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} \qquad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}.$$

This last equation is called the parametric vector form of the solution.

It is obtained by listing equations for all the variables, in order, including the free ones, and making a vector equation.


Example, continued

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix}$$
?

Answer: $x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ for any x_2 in **R**. The solution set is Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$.



[interactive]

Note: one free variable means the solution set is a line in \mathbf{R}^2 (2 = # variables = # columns).

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -1 & 2 \\ -2 & 2 & -4 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & -1 & 2 \\ 2 & -2 & 4 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equation}} \qquad x_1 - x_2 + 2x_3 = 0$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = x_2 - 2x_3 \\ x_2 = x_2 \\ x_3 = x_3 \end{cases}$$


$$\xrightarrow{\text{parametric vector form}} \qquad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}.$$

Example, continued

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & -1 & 2 \\ -2 & 2 & -4 \end{pmatrix}$$
?



[interactive]

Note: two free variables means the solution set is a plane in \mathbb{R}^3 (3 = # variables = # columns).

Question

What is the solution set of Ax = 0, where A =

$$\begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\stackrel{\text{equations}}{\longleftrightarrow} \begin{cases} x_1 & -8x_3 - 7x_4 = 0 \\ x_2 + 4x_3 + 3x_4 = 0 \end{cases}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x_1 = 8x_3 + 7x_4 \\ x_2 = -4x_3 - 3x_4 \\ x_3 = x_3 \\ x_4 = x_4 \end{cases}$$

parametric vector form
$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = x_3 \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix}.$$

Example, continued

Question

What is the solution set of Ax = 0, where

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix}?$$

Answer: Span
$$\left\{ \begin{pmatrix} 8 \\ -4 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 7 \\ -3 \\ 0 \\ 1 \end{pmatrix} \right\}$$
.

[not pictured here]

Note: two free variables means the solution set is a plane in \mathbf{R}^4 (4 = # variables = # columns).

Parametric Vector Form

Homogeneous systems

Let A be an $m \times n$ matrix. Suppose that the free variables in the homogeneous equation Ax = 0 are, for example, x_3, x_6 , and x_8 .

- 1. Find the reduced row echelon form of A.
- 2. Write the parametric form of the solution set, including the redundant equations $x_3 = x_3$, $x_6 = x_6$, and $x_8 = x_8$. Put equations for all of the x_i in order.
- 3. Make a single vector equation from these equations by putting x_3 , x_6 , and x_8 as coefficients of vectors v_3 , v_6 , and v_8 , respectively.

The solutions to Ax = 0 will then be expressed in the form

$$x = x_3v_3 + x_6v_6 + x_8v_8$$

for some vectors v_3 , v_6 , v_8 in \mathbb{R}^n , and any scalars x_3 , x_6 , x_8 .

In this case, the solution set to Ax = 0 is

Span
$$\{v_3, v_6, v_8\}$$
.

The equation above is called the parametric vector form of the solution.

We emphasize the fact that the set of solutions to Ax = 0 is a span.

Poll

How many solutions can there be to a homogeneous system with more equations than variables?

- A. 0
- B. 1
- **C**. ∞

The trivial solution is always a solution to a homogeneous system, so answer A is impossible.

This matrix has only one solution to Ax = 0: [interactive]

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

This matrix has infinitely many solutions to Ax = 0: [interactive]

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$$

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & -3 & | & -3 \\ 2 & -6 & | & -6 \end{pmatrix} \quad \overset{\text{row reduce}}{\longrightarrow} \quad \begin{pmatrix} 1 & -3 & | & -3 \\ 0 & 0 & | & 0 \end{pmatrix}$$

$$\xrightarrow{\text{equation}} \qquad x_1 - 3x_2 = -3$$

$$\xrightarrow{\text{parametric form}} \qquad \begin{cases} x_1 & = 3x_2 - 3 \\ x_2 & = x_2 + 0 \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} \qquad x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

The only difference from the homogeneous case is the constant vector $p = \binom{-3}{0}$.

Note that p is itself a solution: take $x_2 = 0$.

Inhomogeneous Systems

Example, continued


Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -3 \\ 2 & -6 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -3 \\ -6 \end{pmatrix}?$$

Answer:
$$x = x_2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}$$
 for any x_2 in **R**.

This is a *translate* of Span $\left\{ \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$: it is the parallel line through $p = \begin{pmatrix} -3 \\ 0 \end{pmatrix}$.



It can be written

$$\mathsf{Span}\left\{ \begin{pmatrix} \mathbf{3} \\ \mathbf{1} \end{pmatrix} \right\} + \begin{pmatrix} -3 \\ 0 \end{pmatrix}.$$

[interactive]

Inhomogeneous Systems Example

Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -1 & 2 \\ -2 & 2 & -4 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 1 \\ -2 \end{pmatrix}?$$

$$\begin{pmatrix} 1 & -1 & 2 & 1 \\ -2 & 2 & -4 & -2 \end{pmatrix} \quad \overset{\text{row reduce}}{\longrightarrow} \quad \begin{pmatrix} 1 & -1 & 2 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\stackrel{\text{equation}}{\longrightarrow} \quad x_1 - x_2 + 2x_3 = 1$$

$$\underset{\text{parametric form}}{\longrightarrow} \quad \begin{cases} x_1 = x_2 - 2x_3 + 1 \\ x_2 = x_2 \\ x_3 = x_3 \end{cases}$$

$$\underset{\text{parametric vector form}}{\longrightarrow} \quad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = x_2 \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

Inhomogeneous Systems

Example, continued


Question

What is the solution set of Ax = b, where

$$A = \begin{pmatrix} 1 & -1 & 2 \\ -2 & 2 & -4 \end{pmatrix} \quad \text{ and } \quad b = \begin{pmatrix} 1 \\ -2 \end{pmatrix}?$$

and
$$b = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$
?

Answer: Span
$$\left\{ \begin{pmatrix} 1\\1\\0 \end{pmatrix}, \begin{pmatrix} -2\\0\\1 \end{pmatrix} \right\} + \begin{pmatrix} 1\\0\\0 \end{pmatrix}$$
.



The solution set is a translate of

Span
$$\left\{ \begin{pmatrix} 1\\1\\0 \end{pmatrix}, \begin{pmatrix} -2\\0\\1 \end{pmatrix} \right\}$$
:

it is the parallel plane through

$$p = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

[interactive]

Homogeneous vs. Inhomogeneous Systems

Key Observation


The set of solutions to Ax = b, if it is nonempty, is obtained by taking one **specific** or **particular solution** p to Ax = b, and adding all solutions to Ax = 0.

Why? If Ap = b and Ax = 0, then

$$A(p+x) = Ap + Ax = b + 0 = b,$$

so p + x is also a solution to Ax = b.

We know the solution set of Ax = 0 is a span. So the solution set of Ax = b is a *translate* of a span: it is *parallel* to a span. (Or it is empty.)



This works for *any* specific solution p: it doesn't have to be the one produced by finding the parametric vector form and setting the free variables all to zero, as we did before.

[interactive 1] [interactive 2]

Solution Sets and Spans of Columns

Very Important

Let A be an $m \times n$ matrix. There are now two *completely different* things you know how to describe using spans:

- ▶ The **solution set**: for fixed *b*, this is all *x* such that Ax = b.
 - This is a span if b = 0, or a translate of a span in general (if it's consistent).
 - ▶ Lives in \mathbf{R}^n .
 - Computed by finding the parametric vector form.
- ► The span of the columns: this is all b such that Ax = b is consistent.
 - This is the span of the columns of A.
 - Lives in R^m.

Don't confuse these two geometric objects!

Much of the first midterm tests whether you understand both.

[interactive]

Summary

- ▶ The solution set to a **homogeneous** system Ax = 0 is a span. It always contains the **trivial solution** x = 0.
- ▶ The solution set to a **nonhomogeneous** system Ax = b is either empty, or it is a translate of a span: namely, it is a translate of the solution set of Ax = 0.
- ▶ The solution set to Ax = b can be expressed as a translate of a span by computing the **parametric vector form** of the solution.
- ▶ The solution set to Ax = b and the span of the columns of A (from the previous lecture) are two completely different things, and you have to understand them separately.