

Lecture Notes 1

1 Curves

1.1 Definition and Examples

A (parametrized) *curve* (in Euclidean space) is a mapping $\alpha: I \rightarrow \mathbf{R}^n$, where I is an interval in the real line. We also use the notation

$$I \ni t \xrightarrow{\alpha} \alpha(t) \in \mathbf{R}^n,$$

which emphasizes that α sends each element of the interval I to a certain point in \mathbf{R}^n . We say that α is (of the class of) C^k provided that it is k times continuously differentiable. We shall always assume that α is continuous (C^0), and whenever we need to differentiate it we will assume that α is differentiable up to however many orders that we may need.

Some standard examples of curves are a *line* which passes through a point $p \in \mathbf{R}^n$, is parallel to the vector $v \in \mathbf{R}^n$, and has constant speed $\|v\|$

$$[0, 2\pi] \ni t \xrightarrow{\alpha} p + tv \in \mathbf{R}^n;$$

a *circle* of radius \mathbf{R} in the plane, which is oriented counterclockwise,

$$[0, 2\pi] \ni t \xrightarrow{\alpha} (r \cos(t), r \sin(t)) \in \mathbf{R}^2;$$

and the right handed *helix* (or corkscrew) given by

$$\mathbf{R} \ni t \xrightarrow{\alpha} (r \cos(t), r \sin(t), t) \in \mathbf{R}^3.$$

Other famous examples include the *figure-eight* curve

$$[0, 2\pi] \ni t \xrightarrow{\alpha} (\sin(t), \sin(2t)) \in \mathbf{R}^2,$$

¹Last revised: September 11, 2004

the *parabola*

$$\mathbf{R} \ni t \xrightarrow{\alpha} (t, t^2) \in \mathbf{R}^2,$$

and the *cubic curve*

$$\mathbf{R} \ni t \xrightarrow{\alpha} (t, t^2, t^3) \in \mathbf{R}^3.$$

Exercise 1. Sketch the cubic curve (*Hint:* First draw each of the projections into the xy , yz , and zx planes).

Exercise 2. Find a formula for the curve which is traced by the motion of a fixed point on a wheel of radius r rolling with constant speed on a flat surface (*Hint:* Add the formula for a circle to the formula for a line generated by the motion of the center of the wheel. You only need to make sure that the speed of the line correctly matches the speed of the circle).

Exercise 3. Let $\alpha: I \rightarrow \mathbf{R}^n$, and $\beta: J \rightarrow \mathbf{R}^n$ be a pair of differentiable curves. Show that

$$\left(\langle \alpha(t), \beta(t) \rangle \right)' = \langle \alpha'(t), \beta(t) \rangle + \langle \alpha(t), \beta'(t) \rangle$$

and

$$\left(\|\alpha(t)\| \right)' = \frac{\langle \alpha(t), \alpha'(t) \rangle}{\|\alpha(t)\|}.$$

(*Hint:* The first identity follows immediately from the definition of the inner-product, together with the ordinary product rule for derivatives. The second identity follows from the first once we recall that $\|\cdot\| := \langle \cdot, \cdot \rangle^{1/2}$).

Exercise 4. Show that if α has unit speed, i.e., $\|\alpha'(t)\| = 1$, then its velocity and acceleration are orthogonal, i.e., $\langle \alpha(t), \alpha''(t) \rangle = 0$.

Exercise 5. Show that if the position vector and velocity of a planar curve $\alpha: I \rightarrow \mathbf{R}^2$ are always perpendicular, i.e., $\langle \alpha(t), \alpha'(t) \rangle = 0$, for all $t \in I$, then $\alpha(I)$ lies on a circle centered at the origin of \mathbf{R}^2 .

Exercise 6. Use the fundamental theorem of Calculus for real valued functions to show:

$$\alpha(b) - \alpha(a) = \int_a^b \alpha'(t) dt.$$

Exercise 7. Prove that

$$\|\alpha(b) - \alpha(a)\| \leq \int_a^b \|\alpha'(t)\| dt.$$

(*Hint:* Use the fundamental theorem of calculus and the Cauchy-Schwartz inequality to show that for any unit vector $u \in \mathbf{R}^n$,

$$\langle \alpha(b) - \alpha(a), u \rangle = \int_a^b \langle \alpha'(t), u \rangle dt \leq \int_a^b \|\alpha'(t)\| dt.$$

Then set $u := (\alpha(b) - \alpha(a)) / \|\alpha(b) - \alpha(a)\|$.

1.2 Reparametrization

We say that $\beta: J \rightarrow \mathbf{R}^n$ is a *reparametrization* of $\alpha: I \rightarrow \mathbf{R}^n$ provided that there exists a smooth bijection $\theta: I \rightarrow J$ such that $\alpha(t) = \beta(\theta(t))$. In other words, the following diagram commutes:

$$\begin{array}{ccc} & \mathbf{R}^n & \\ \alpha \nearrow & & \nwarrow \beta \\ I & \xrightarrow{\theta} & J \end{array}$$

For instance $\beta(t) = (\cos(2t), \sin(2t))$, $0 \leq t \leq \pi$, is a reparametrization $\alpha(t) = (\cos(t), \sin(t))$, $0 \leq t \leq 2\pi$, with $\theta: [0, 2\pi] \rightarrow [0, \pi]$ given by $\theta(t) = t/2$.

The *geometric quantities* associated to a curve do not change under reparametrization. These include length and curvature as we define below.

1.3 Length and Arclength

By a *partition* P of an interval $[a, b]$ we mean a collection of points $\{t_0, \dots, t_n\}$ of $[a, b]$ such that

$$a = t_0 < t_1 < \dots < t_n = b.$$

The *approximation of the length of α with respect to P* is defined as

$$\text{length}[\alpha, P] := \sum_{i=1}^n \|\alpha(t_i) - \alpha(t_{i-1})\|,$$

and if $\text{Partition}[a, b]$ denotes the set of all partitions of $[a, b]$, then the *length* of α is given by

$$\text{length}[\alpha] := \sup \{ \text{length}[\alpha, P] \mid P \in \text{Partition}[a, b] \},$$

where ‘sup’ denotes the supremum or the least upper bound.

Exercise 8. Show that the shortest curve between any pairs of points in \mathbf{R}^n is the straight line segment joining them. (*Hint:* Use the triangle inequality).

We say that a curve is *rectifiable* if it has finite length.

Exercise* 9 (Nonrectifiable curves). Show that there exists a curve $\alpha: [0, 1] \rightarrow \mathbf{R}^2$ which is not rectifiable (*Hint:* One such curve, known as the *Koch curve* (Figure 1), may be obtained as the limit of a sequence of curves $\alpha_i: [0, 1] \rightarrow \mathbf{R}^2$ defined as follows. Let α_0 trace the line segment $[0, 1]$. Consider an equilateral triangle of sides $1/3$ whose base rests on the middle third of $[0, 1]$. Deleting this middle third from the interval and the triangle yields the curve traced by α_1 . Repeating this procedure on each of the 4



Figure 1:

subsegments of α_1 yields α_2 . Similarly α_{i+1} is obtained from α_i . You need to show that α_i converge to a (continuous) curve, which may be done using the Arzela-Ascoli theorem. It is easy to see that this limit has infinite length, because the length of α_i is $(4/3)^i$. Another example of a nonrectifiable curve $\alpha: [0, 1] \rightarrow \mathbf{R}^2$ is given by $\alpha(t) := (t, t \sin(\pi/t))$, when $t \neq 0$, and $\alpha(t) := (0, 0)$ otherwise. The difficulty here is to show that the length is infinite.)

If a curve is C^1 , then its length may be computed as the following exercise shows. Note also that the following exercise shows that a C^1 curve over a compact domain is rectifiable.

Exercise* 10 (Length of C^1 curves). Show that if $\alpha: I \rightarrow \mathbf{R}^n$ is a C^1 curve, then

$$\text{length}[\alpha] = \int_I \|\alpha'(t)\| dt.$$

(*Hints:* It suffices to show that (i) $\text{length}[\alpha, P]$ is less than the above integral, for all $P \in \text{Partition}[a, b]$, and (ii) there exists a sequence P_n of partitions such that $\lim_{n \rightarrow \infty} \text{length}[\alpha, P_n]$ is equal to the integral. The first part follows quickly from Exercise 7. To prove the second part, let P_n be a partition given by $t_i := a + i(b-a)/n$. Any other partition where the maximum length of the subsegments converges to zero would do as well. Then apply the mean value theorem, and recall the definition of the integral as the limit of Riemann sums. See do Carmo Exc. 8, Sec. 1-3, and the solution on p. 475.)

Exercise 11. Compute the length of a circle of radius r , and the length of one cycle of the curve traced by a point on a circle of radius r rolling on a straight line.

Exercise 12 (Invariance of length under reparametrization). Show that if β is a reparametrization of α , then $\text{length}[\beta] = \text{length}[\alpha]$, i.e., length is invariant under reparametrization (*Hint:* you only need to recall the chain rule together with the integration by substitution.)

Let $L := \text{length}[\alpha]$. The *arclength* function of α is a mapping $s: [a, b] \rightarrow [0, L]$ given by

$$s(t) := \int_a^t \|\alpha'(u)\| du.$$

Thus $s(t)$ is the length of the subsegment of α which stretches from the initial time a to time t .

Exercise 13 (Regular curves). Show that if α is a *regular* curve, i.e., $\|\alpha'(t)\| \neq 0$ for all $t \in I$, then $s(t)$ is an invertible function, i.e., it is one-to-one (*Hint:* compute $s'(t)$).

Exercise 14 (Reparametrization by arclength). Show that every regular curve $\alpha: [a, b] \rightarrow \mathbf{R}^n$, may be reparametrized by arclength (*Hint:* Define $\beta: [0, L] \rightarrow \mathbf{R}^n$ by $\beta(t) := \alpha(s^{-1}(t))$, and use the chain rule to show that $\|\beta'\| = 1$; you also need to recall that since $f(f^{-1}(t)) = t$, then, again by chain rule, we have $(f^{-1})'(t) = 1/f'(f^{-1}(t))$ for any smooth function f with nonvanishing derivative.)

1.4 Cauchy's integral formula and curves of constant width

Let $\alpha: I \rightarrow \mathbf{R}^2$ be a curve and $u(\theta) := (\cos(\theta), \sin(\theta))$ be a unit vector. The projection of α into the line passing through the origin and parallel to u is given by $\alpha_u(t) := \langle \alpha(t), u \rangle u$.

Exercise 15 (Cauchy's integral formula). Show that if $\alpha: I \rightarrow \mathbf{R}^2$ has length L , then the average length of the projections α_u , over all directions, is $2L/\pi$, i.e.,

$$\frac{1}{2\pi} \int_0^{2\pi} \text{length}[\alpha_{u(\theta)}] d\theta = \frac{2L}{\pi}.$$

(*Hint:* First prove this fact for the case when α traces a line segment. Then a limiting argument settles the general case, once you recall the definition of length. Also, there is a purely analytic and more elegant proof for the case when α is C^1 . In this case the integrand is simply $\int_I \|\alpha'(t)\| |\cos(\theta)| dt$.)

As an application of the above formula we may obtain a sharp inequality involving *width* of *closed* curves. The width of a set $X \subset \mathbf{R}^2$ is the distance between the closest pairs of parallel lines which contain X in between them. For instance the width of a circle of radius r is $2r$. A curve $\alpha: [a, b] \rightarrow \mathbf{R}^2$ is said to be closed provided that $\alpha(a) = \alpha(b)$. We should also mention that α is a C^k closed curve provided that the (one-sided) derivatives of α match up at a and b .

Exercise 16 (Width and length). Show that if $\alpha: [a, b] \rightarrow \mathbf{R}^2$ is a closed curve with width w and length L , then

$$w \leq \frac{L}{\pi}.$$

Note that the above inequality is sharp, since for circles $w = L/\pi$. Are there other curves satisfying this property? The answer may surprise you. For any unit vector $u(\theta)$, the width of a set $X \subset \mathbf{R}^2$ in the direction u , w_u , is defined as the distance between the closest pairs of lines which contain X in between them. We say that a closed curve in the plane has *constant width* provided that w_u is constant in all directions.

Exercise 17. Show that if the equality in Exercise 16 holds then α is a curve of constant width.

The last exercise would have been insignificant if circles were the only curves of constant width, but that is not the case:

Exercise 18 (Reuleaux triangle). Consider three disks of radius r whose centers are on an equilateral triangle of sides r , see Figure 2. Show that the curve which bounds the intersection of these disks has constant width. Also show that similar constructions for any regular polygon yield curves of constant width.




Figure 2:

It can be shown that of all curves of constant width w , Reuleaux triangle has the least area. This is known as the Blaschke-Lebesgue theorem. A recent proof of this result has been obtained by Evans Harrell.

Note that the Reuleaux triangle is not a C^1 regular curve for it has sharp corners. To obtain a C^1 example of a curve of constant width, we may take a curve which is a constant distance away from the Reuleaux triangle. Further, a C^∞ example may be constructed by taking an *evolute* of a *deltoid*, see Gray p. 177.