

## A SHORT REMARK ABOUT $L^2$ -SPACES

In this section we establish that the space of all square integrable functions form a Hilbert space. To start, consider all continuous functions on some interval  $I$  which may be the half line or the whole real line and define

$$L^2(I) = \left\{ f : \int_I |f(x)|^2 dx < \infty \right\} .$$

It is quite easy to verify that  $L^2(I)$  is a linear space with inner product

$$(f, g) = \int_I \overline{f(x)} g(x) dx .$$

Unfortunately, this spaces is not complete. Consider  $I = [-1, 1]$  and the sequence of functions  $f^{(k)}(x) = -1$  for  $-1 \leq x \leq -\frac{1}{k}$ ,  $f^{(k)}(x) = kx$  for  $-\frac{1}{k} \leq x \leq \frac{1}{k}$  and  $f^{(k)}(x) = 1$  for  $\frac{1}{k} \leq x \leq 1$ . Clearly these functions are continuous for each  $k = 1, 2, \dots$ . If  $\ell \geq k$  we have that

$$\int_I |f^{(\ell)}(x) - f^{(k)}(x)|^2 dx = \int_{-\frac{1}{k}}^{\frac{1}{k}} |f^{(\ell)}(x) - f^{(k)}(x)|^2 dx \leq 4 \times \frac{2}{k}$$

from which we see that  $f^{(k)}$  is a Cauchy sequence. The limit of this sequence, however, is not a continuous function and the limit is not in our linear space. There is a process of completing this space at the price that the integral has to be interpreted according to Lebesgue.

The idea is the following. Consider a positive function  $f$  and we want to give a definition of

$$\int_I f(x) dx .$$

We interpret this integral as the area underneath the graph of  $f$  over  $I$ . One way of approximating this area is according to Riemann which you have learned in your analysis course. Another one is to look at the **length** of the **level sets** of the function  $f$  which is given by

$$\{x \in I : f(x) > t\} .$$

If we denote by  $|\{x \in I : f(x) > t\}|$  the length of these level sets we can think of the area as

$$\int_0^M |\{x \in I : f(x) > t\}| dt , \tag{1}$$

where  $M$  is the maximal value of  $f$ . Note that  $|\{x \in I : f(x) > t\}|$  is a decreasing function of  $t$  and hence it is Riemann integrable.

Now observe, that the level sets can be quite crazy sets that do not necessarily have a length. So the first step is to state properties that such sets must have in order to have a chance of making sense out of this integral. We call such sets **measurable** and require the following:

- a) If  $A \subset I$  is measurable, so is its complement  $A^c$ .
  - b)  $I$  is measurable.
  - c) If  $A_1, A_2, \dots$  is a countable family of measurable sets, then their union is also measurable.
- Any collection of sets that have the above properties we call a **sigma algebra**.

In a further step we now define what we mean by the volume of such sets, i.e., the **measure** of such sets. A **measure**  $\mu$  is a function from a sigma algebra  $\Sigma$  into the positive real numbers that has the following properties

- a)  $\mu(A) \leq \mu(B)$  if  $A \subset B$  and  $A, B \in \Sigma$ .
- b) Let  $A_1, A_2, \dots$  be a **countable** collection of **disjoint** sets in  $\Sigma$ . Then

$$\mu(\cup_{j=1}^{\infty} A_j) = \sum_{j=1}^{\infty} \mu(A_j) .$$

This last property is called **countable additivity** of the the measure  $\mu$ . This property is the key in establishing completeness of spaces of integrable functions.

A consequence of the countable additivity are the following two statements:

- a) If  $A_1 \subset A_2 \subset \dots$  is an increasingly nested sequence of sets in  $\Sigma$ , then

$$\lim_{N \rightarrow \infty} \mu(\cup_{j=1}^N A_j) = \mu(\cup_{j=1}^{\infty} A_j)$$

and

- b) If  $A_1 \supset A_2 \supset \dots$  is a decreasingly nested sequence of set in  $\Sigma$ , then

$$\lim_{N \rightarrow \infty} \mu(\cap_{j=1}^N A_j) = \mu(\cap_{j=1}^{\infty} A_j) .$$

Now we close in on our definition of the integral. A function  $f : I \rightarrow \mathbb{R}_+$  is **measurable** if the sets  $\{x \in I : f(x) > t\}$  are measurable for all  $t \in \mathbb{R}$ .

Given a non-negative measurable function  $f$  and a measure  $\mu$  we say that the function is **summable** or **integrable** if

$$\int_I f(x) \mu(dx) := \int \mu(\{x \in I : f(x) > t\}) dt < \infty$$

where, as before, the last integral is a Riemann integral, since the function  $t \rightarrow \mu(\{x \in I : f(x) > t\})$  is decreasing.

**Remark 0.1.** *There could be sets that have zero measure. Thus modifying the function on a set of zero measure would not affect the integral. We say that a certain property holds **almost everywhere with respect to**  $\mu$  if the set where the property does not hold has zero  $\mu$  measure.*

There are two important theorems that follow from these definitions.

**Theorem 0.2. Monotone convergence** *Let  $f^{(k)}$  be a sequence of summable functions and assume  $f^{(k)}(x) \leq f^{(k+1)}(x)$  for almost all  $x \in I$  and assume that  $\int_I [f^{(k)}(x) - f^{(1)}(x)] \mu(dx)$  is uniformly bounded. Then the limit*

$$\lim_{k \rightarrow \infty} f^{(k)}(x) := f(x)$$

*exists for almost every  $x$  and is measurable. Moreover,*

$$\lim_{k \rightarrow \infty} \int_I f^{(k)}(x) \mu(dx) = \int_I f(x) \mu(dx) ,$$

*in the sense that if one of the quantities is infinite so is the other.*

The other important theorem is

**Theorem 0.3. Dominated convergence** *Let  $f^{(k)}(x)$  be a sequence of summable functions that converges almost everywhere with respect to  $\mu$  to a function  $f$ . If there exists a summable function  $G(x)$  such that*

$$|f^{(k)}(x)| \leq G(x)$$

*for all  $k = 1, 2, 3, \dots$ , then  $f(x)$  is measurable, summable and*

$$\lim_{k \rightarrow \infty} \int_I f^{(k)}(x) \mu(dx) = \int_I f(x) \mu(dx) .$$

The big question is whether such  $\sigma$  algebras and measures exist. This is the hard part of the theory and you are referred to the books on measure theory. On the real line there exists a unique translation invariant measure  $\mathcal{L}$ , the Lebesgue measure. Translation invariant means that  $\mathcal{L}(B) = \mathcal{L}(A)$  whenever  $B$  is a translate of  $A$ .

The beauty of all this is that it works in great generality. We can replace the interval  $I$  by any set  $\Omega$  and  $\mu$  be any measure on a sigma algebra of subsets of  $\Omega$ . The theorems stated above continue to hold in this case too.

**Definition 0.4.  $L^2(\Omega, \mu)$ -space** *This space consists of all square summable functions  $f : \Omega \rightarrow \mathbb{C}$ .*

We are ready to state the important

**Theorem 0.5. Riesz-Fischer** *The space  $L^2(\Omega, \mu)$  endowed with the inner product*

$$(f, g) = \int_{\Omega} \overline{f(x)} g(x) \mu(dx)$$

*is a Hilbert space.*

*Proof.* Let  $f^{(k)}$  be a Cauchy sequence in  $L^2(\Omega, \mu)$ . This means that for any  $\varepsilon > 0$  there exists  $N$  so that for all  $k, \ell > N$

$$\|f^{(k)} - f^{(\ell)}\| < \varepsilon .$$

Hence, there exists  $k_1$  so that for all  $\ell > k_1$

$$\|f^{(k_1)} - f^{(\ell)}\| < \frac{1}{2} .$$

Likewise, there exists  $k_2 > k_1$  so that for all  $\ell > k_2$

$$\|f^{(k_2)} - f^{(\ell)}\| < \frac{1}{2^2} .$$

Continuing this way we find a sequence  $k_1, k_2, k_3, \dots$  such that for all  $j = 1, 2, \dots$

$$\|f^{(k_j)} - f^{(k_{j+1})}\| < \frac{1}{2^j} .$$

Now consider the sequence  $f^{(k_j)}$  and write

$$f^{(k_j)} = f^{(k_1)} + [f^{(k_2)} - f^{(k_1)}] + [f^{(k_3)} - f^{(k_2)}] + \dots + [f^{(k_j)} - f^{(k_{j-1})}] .$$

If we set

$$F^{(j)} = |f^{(k_1)}| + |f^{(k_2)} - f^{(k_1)}| + |f^{(k_3)} - f^{(k_2)}| + \dots + |f^{(k_j)} - f^{(k_{j-1})}| ,$$

we obviously have that

$$|f^{(k_j)}| \leq F^{(j)} .$$

The sequence  $F^{(j)}(x)$  is a monotone increasing sequence and hence converges to a function  $F(x)$ . This implies that the sequence  $f^{(k_j)}(x)$  converges to some function  $f(x)$  since the partial sums converge absolutely. Further,

$$\|F^{(j)}\| \leq \|f^{(k_1)}\| + \|f^{(k_2)} - f^{(k_1)}\| + \|f^{(k_3)} - f^{(k_2)}\| + \dots + \|f^{(k_j)} - f^{(k_{j-1})}\|$$

which is bounded above by

$$\|f^{(k_1)}\| + \frac{1}{2} + \frac{1}{2^2} + \dots + \frac{1}{2^j} < \|f^{(k_1)}\| + 1 .$$

Hence by the monotone convergence theorem we find that  $F$  is square summable and

$$|f^{(k_j)}(x) - f(x)| \leq |f^{(k_j)}(x)| + |f(x)| \leq 2F(x) .$$

Since  $f^{(k_j)}(x)$  converges to  $f(x)$  for every  $x$  we have by the dominated convergence theorem that

$$\lim_{j \rightarrow \infty} \int_{\Omega} |f^{(k_j)}(x) - f(x)|^2 \mu(dx) = 0 .$$

In other words we have for the *subsequence*  $k_j$  that

$$\lim_{j \rightarrow \infty} \|f^{(k_j)} - f\| = 0 .$$

We have to show that the whole sequence converges. For this, fix and  $\varepsilon > 0$  and pick  $N$  such that for  $k_j > N$ ,  $\|f^{(k_j)} - f\| < \varepsilon/2$  and for  $\ell > N$ ,  $\|f^{(k_j)} - f^{(\ell)}\| < \varepsilon/2$ . Then for all  $\ell > N$

$$\|f^{(\ell)} - f\| \leq \|f^{(k_j)} - f^{(\ell)}\| + \|f^{(k_j)} - f\| < \varepsilon .$$

□