

**Math 4305, Fall 2014,  
Midterm 1, Practice solutions**

**Problem 1** Suppose that the matrix below is the augmented matrix of a system of linear equations

$$\begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 3 & 2 \\ 0 & -2 & 0 & 1 & 3 \\ 0 & 0 & 1 & k & h \end{pmatrix}$$

a)(6 points) For what values of  $h$  and  $k$ , this system has no solution.

**Solution:** By interchanging  $r_2$  and  $r_3$ , and  $r_4 - r_3$ , one arrives the REF of the matrix:

$$\begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & -2 & 0 & 1 & 3 \\ 0 & 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & k-3 & h-2 \end{pmatrix}$$

Now, it's easy to see that, the system has no solution if and only if the rightmost column is pivot. This happens if and only if  $k = 3$  and  $h \neq 2$ .

b) (7 points) For what values of  $h$  and  $k$ , this system has a unique solution. Find the solution.

**Solution:** Based on the REF derived from part a), the system has a unique solution if and only if  $k \neq 3$ . In this case, the system is consistent without free variable. In order to solve the system, we row reduce the REF into RREF. This is achieved by  $\frac{1}{k-3}r_4$ ,  $-\frac{1}{2}r_2$ ,  $r_3 - 3r_4$ ,  $r_2 + \frac{1}{2}r_4$  and  $r_1 - 2r_2$ . The RREF is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 4 - Y \\ 0 & 1 & 0 & 0 & -\frac{3}{2} + \frac{1}{2}Y \\ 0 & 0 & 1 & 0 & 2 - 3Y \\ 0 & 0 & 0 & 1 & Y \end{pmatrix},$$

where  $Y = \frac{h-2}{k-3}$ . So the solution is

$$x_1 = 4 - Y, x_2 = -\frac{3}{2} + \frac{1}{2}Y, x_3 = 2 - 3Y \text{ and } x_4 = Y.$$

c)(7 points) For what values of  $h$  and  $k$ , this system has infinitely many solutions. Describe the set of all solutions using parametric vector form.

**Solution:** From part a), the system has infinitely many solutions if and only if  $k = 3$  and  $h = 2$ . In this case, the system is consistent with a free variable  $x_4$ . The REF is now

$$\begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & -2 & 0 & 1 & 3 \\ 0 & 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

To solve the system, we row reduce the above matrix to RREF by  $r_1 + r_2$  and  $-\frac{1}{2}r_2$ :

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 4 \\ 0 & 1 & 0 & -\frac{1}{2} & -\frac{3}{2} \\ 0 & 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Thus,  $x_1 = 4 - x_4$ ,  $x_2 = -\frac{3}{2} + \frac{1}{2}x_4$ ,  $x_3 = 2 - 3x_4$  and  $x_4$  is free. So the solution is described by

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 4 \\ -\frac{3}{2} \\ 2 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -1 \\ \frac{1}{2} \\ -3 \\ 1 \end{pmatrix}.$$

**Problem 2** Let  $\mathbf{v} = (1, 0, 1)^t$ . Define the linear transformation  $T : \mathbf{R}^3 \rightarrow \mathbf{R}^3$  by  $T(\mathbf{x}) = \mathbf{v} \times \mathbf{x}$ . Where  $\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix}$ .

a) Find the standard matrix  $A$  of  $T$ .

**Solution**  $A = [\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3]$ , where  $\mathbf{a}_i = T(\mathbf{e}_i)$ .

$$T(\mathbf{e}_1) = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, T(\mathbf{e}_2) = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, T(\mathbf{e}_3) = \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix}.$$

We thus have

$$A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

b) Find a basis of  $im(A)$ .

**Solution:** We do the interchange of  $r_1$  and  $r_2$ , then  $r_3 + r_2$ , we thus reach the REF of  $A$ :

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore, we know that a basis of  $im(A)$  is  $\{\mathbf{a}_1, \mathbf{a}_2\}$ .

c) What's the dimension of  $\ker(A)$ ?

**Solution** By the Rank Theorem, we know that

$$\dim \ker(A) = 3 - \dim \operatorname{im}(A) = 1.$$

**Problem 3** Consider an  $m \times n$  matrix  $A$  and an  $n \times m$  matrix  $B$  (with  $n \neq m$ ) such that  $AB = I_m$ . Are the columns of  $B$  linearly independent? What about columns of  $A$ ?

**Solution:** If columns of  $B$  are linearly dependent, so are columns of  $AB$ , which contradicts to  $AB = I_m$ . Or, we could show that columns of  $B$  are linearly independent directly. To show this, we assume there is a vector  $\mathbf{x} \in \mathbf{R}^m$ , such that  $B\mathbf{x} = 0$ . Then we have  $\mathbf{x} = I_m\mathbf{x} = AB\mathbf{x} = A0 = 0$ . Thus, if  $AB = I_m$ , then columns of  $B$  are linearly independent. Furthermore, we know that  $n > m$ . Since  $A$  is  $m \times n$ , columns of  $A$  are linearly dependent.

**Problem 4** Let  $S = \{(x, y) : xy \geq 0\}$  be a subset of the plane  $\mathbf{R}^2$ . Is  $S$  a subspace of  $\mathbf{R}^2$ ?

**Solution:**  $S$  is not a subspace of  $\mathbf{R}^2$ . One can easily verify that it is not closed for addition. Choose  $\mathbf{v} = (-1, 0)$  and  $\mathbf{u} = (0, 1)$ , both are in  $S$ , however,  $\mathbf{v} + \mathbf{u} = (-1, 1)$  is not in  $S$ .

**Problem 5** For which values of the constant  $k$  is the following matrix invertible? Find the inverse.

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & k \\ 1 & 4 & k^2 \end{pmatrix}$$

**Solution:** Row reduce the matrix into REF by  $r_2 - r_1$ ,  $r_3 - r_1$ ,  $r_3 - 3r_2$ ,

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & k-1 \\ 0 & 0 & k^2-3k+2 \end{pmatrix}.$$

The matrix is invertible if  $k^2 - 3k + 2 \neq 0$ . Thus, if  $k \neq 1$  and  $k \neq 2$ , the matrix is invertible. For  $k \neq 1$  and  $k \neq 2$ , we denote the nonzero quantity  $k^2 - 3k + 2$  by  $N$ , let  $M = k - 1$ , thus Gauss-Jordan algorithm will give the inverse

$$\frac{1}{N} \begin{pmatrix} 2M + 2N - 2 & 3 - 3M - N & M - 1 \\ -N - 2M & N + 3M & -M \\ 2 & -3 & 1 \end{pmatrix}.$$