

RATIONALITY OF THE $SL(2, \mathbb{C})$ -REIDEMEISTER TORSION IN DIMENSION 3

JEROME DUBOIS AND STAVROS GAROUFALIDIS

ABSTRACT. If M is a finite volume complete hyperbolic 3-manifold with one cusp, the geometric component X_M of its $SL(2, \mathbb{C})$ -character variety is an affine complex curve, which is smooth at the discrete faithful representation ρ_0 . Porti defined a non-abelian Reidemeister torsion in a neighborhood of ρ_0 in X_M and observed that it is an analytic map. In the present paper we prove that this non-abelian Reidemeister torsion is the germ of a unique rational function on X_M . This implies the existence of a polynomial relation of the torsion of a representation and the trace of the meridian or the longitude, and explains why the torsion of the discrete faithful representation lies in the trace field of M . We postulate that the coefficients of the $1/N^k$ -asymptotics of the Parametrized Volume Conjecture for M are elements of the field of rational functions on X_M .

CONTENTS

1. Introduction	1
1.1. The volume of an $SL(2, \mathbb{C})$ -representation and the A -polynomial	1
1.2. The $SL(2, \mathbb{C})$ -character variety of M and its field of rational functions	2
1.3. The Reidemeister torsion of an $SL(2, \mathbb{C})$ -representation	2
1.4. Examples	3
1.5. Problems	4
2. The character variety of hyperbolic 3-dimensional manifolds	5
2.1. Four favors of the character variety, après Dunfield	5
2.2. The coefficient field of augmented representations	6
2.3. Augmented representations and the shape field	6
2.4. Ideal triangulations and the gluing equations variety	7
3. The non-abelian Reidemeister torsion	8
3.1. An explanation of the rationality of the Reidemeister torsion in dimension 3	8
3.2. Proof of Theorem 2	8
3.3. The dependence of the Reidemeister torsion on the admissible curve and the A -polynomial	11
Acknowledgment	11
References	11

1. INTRODUCTION

1.1. The volume of an $SL(2, \mathbb{C})$ -representation and the A -polynomial. A well-known numerical invariant of a 3-dimensional finite volume hyperbolic manifold M with a cusp is its *volume*, a positive real number. A complete invariant of the hyperbolic structure of M is a discrete faithful representation of $\pi_1(M)$ into $PSL(2, \mathbb{C})$ (well-defined up to conjugation) which is also a topological invariant, as follows from Mostow rigidity Theorem. Every $PSL(2, \mathbb{C})$ -representation ρ of $\pi_1(M)$ has a real-valued volume $\text{Vol}(\rho)$; see [Dn,

Date: August 10, 2009 .

S.G. was supported in part by NSF.

1991 *Mathematics Classification.* Primary 57N10. Secondary 57M25.

Key words and phrases: knots, A -polynomial, Reidemeister torsion, volume, character variety, 3-manifolds, hyperbolic geometry, invariant trace field.

Ch.2] and also [F, FK]. When a representation varies in a 1-parameter family ρ_t , the variation of the volume $\frac{d}{dt} \text{Vol}(\rho_t)$ depends only on the restriction of ρ_t to the boundary torus ∂M . This is a general principle of Atiyah-Patodi-Singer, and in our special case it also follows from Schalfi's formula. This raises the question: which $\text{PSL}(2, \mathbb{C})$ -representations of ∂M extend to a representation of M ? The answer is given by an algebraic condition between the eigenvalues of a meridian and longitude of ∂M . This condition is the vanishing of the so-called *A-polynomial* of M ; see [CCGLS]. The *A-polynomial* of M encodes important informations about

- (a) the hyperbolic geometry of M , and determines the variation of the volume of the hyperbolic structure of M .
- (b) the topology of M and more precisely about the slopes of incompressible surfaces in the knot complement, as follows from Culler-Shalen theory; see [CCGLS].

More recently, the *A-polynomial* (or rather, its extension that includes the images of all components of the character variety) is conjecturally linked in two different ways to a *quantum knot invariant*, namely the *colored Jones polynomials* of a knot in 3-space (for a definition of the latter, which we will not use in the present paper, see [Tu] and [GL1]):

- (a) There is an A_q -polynomial in two q -commuting variables which encodes a minimal order linear q -difference equation for the sequence of colored Jones polynomials; see [GL1]. The AJ Conjecture of [Ga1] states that when $q = 1$, the A_q -polynomial coincides with the *A-polynomial*.
- (b) There is a parametrized version of the Volume Conjecture which links the variation of the limit in the Volume Conjecture to the *A-polynomial*; see [GM, GL2].

Aside from conjectures, the following result of [DG] and [BZ] (based on foundational work of Kronheimer-Mrowka) shows that the *A-polynomial* detects the unknot.

Theorem 1. [BZ, DG] *The A-polynomial of a nontrivial knot in 3-space is nontrivial.*

1.2. The $\text{SL}(2, \mathbb{C})$ -character variety of M and its field of rational functions. For historical reasons that simplify the linear algebra, it is useful to consider $\text{SL}(2, \mathbb{C})$ (rather than $\text{PSL}(2, \mathbb{C})$)-representations of $\pi_1(M)$. The discrete faithful representation of M lifts to a $\text{SL}(2, \mathbb{C})$ -representation $\rho_0: \pi_1(M) \rightarrow \text{SL}(2, \mathbb{C})$; see [Cu]. To understand how the $\text{SL}(2, \mathbb{C})$ -representation ρ_0 of $\pi_1(M)$ varies, we consider the unique component X_M of the *$\text{SL}(2, \mathbb{C})$ -character variety* of M that contains ρ_0 . It is well-known that X_M is an affine curve defined over \mathbb{Q} and that ρ_0 is a smooth point of X_M ; see [CCGLS]. Moreover, the coordinate ring $\mathbb{Q}[X_M]$ is generated by tr_γ for all $\gamma \in \pi_1 M$, where tr_γ is the so called *trace-function* defined by:

$$(1) \quad \text{tr}_\gamma: X_M \longrightarrow \mathbb{C}, \quad \text{tr}_\gamma(\rho) = \text{tr}(\rho(\gamma)).$$

Here $\text{tr}(A) = \sum_i a_{ii}$ denotes the trace of a square matrix $A = (a_{ij})$. Let $\mathbb{Q}(X_M)$ denote the field of rational functions of X_M . For a detailed discussion on character varieties, one can refer to Shalen's survey [Sh], to [BDR-V, Sec.10] and also to [CCGLS, Go].

1.3. The Reidemeister torsion of an $\text{SL}(2, \mathbb{C})$ -representation. Another important numerical invariant of a representation of a manifold is its *Reidemeister torsion*, which comes in several combinatorial or analytic flavors, see Milnor's survey [Mi] or Turaev's monograph [T] for details. Combinatorially, torsion is defined in terms of ratios of determinants of matrices assigned to based, acyclic complexes, which themselves are associated with a cell decomposition of a manifold and an acyclic representation. One can define torsion for all (not necessarily acyclic) representations of a manifold as an element of a top exterior power of a twisted (co)homology group, and one can obtain a complex number after choosing a basis for the twisted (co)homology. Porti [Po] defined a Reidemeister torsion for the adjoint representation associated to an $\text{SL}(2, \mathbb{C})$ -representation ρ of $\pi_1(M)$ when ρ is in a *neighborhood* U of $\rho_0 \in X_M$. Such representations are not acyclic and a basis for the twisted homology (and thus the torsion) depends on an *admissible curve* γ , i.e., a simple closed curve γ in ∂M which is not nullhomologous in ∂M (see Porti's monograph [Po, Chap. 3] for details). Thus, the *non-abelian Reidemeister torsion* is a map:

$$(2) \quad \tau_\gamma: U \longrightarrow \mathbb{C}.$$

Moreover, Porti [Po] observed that τ_γ is an analytic map. Our main result shows that *the non-abelian torsion τ_γ is the germ of a unique element of $\mathbb{Q}(X_M)$.*

Theorem 2. *For every admissible curve γ , the non-abelian Reidemeister torsion $\tau_\gamma : U \rightarrow \mathbb{C}$ is the germ of a unique element of $\mathbb{Q}(X_M)$, which is regular at ρ_0 .*

The above theorem allows us to define torsion to representations that are not near the discrete faithful one.

Corollary 1.1. *For every admissible curve γ , $\tau_\gamma(\rho_0)$ lies in the trace field of M .*

Recall that if M is the complement of a knot in a $\mathbb{Z}/2$ -homology sphere, then the trace field and the invariant trace field of M coincide; see [NR, Theorem 2.2]. Our next theorem shows that τ_γ is an *algebraic function* of tr_γ . This follows easily from the fact that τ_γ and tr_γ are rational functions on X_M and that $\mathbb{Q}(X_M)$ has transcendence degree 1, since X_M is an affine curve defined over \mathbb{Q} .

Theorem 3. *For every admissible curve γ , there exists a polynomial $T_\gamma(\tau, y) \in \mathbb{Z}[\tau, y]$, called the T_γ -polynomial, so that*

$$(3) \quad T_\gamma(\tau_\gamma, \text{tr}_\gamma) = 0.$$

Let us make some remarks regarding Theorems 2 and 3.

Remark 1.2. The dependence of the torsion function τ_γ on γ is determined by the A -polynomial; see Equation (17). Thus, T_γ is determined by T_μ and the A -polynomial of M .

Remark 1.3. Theorem 2 is an instance of a well-recorded phenomenon: many classical and quantum invariants of knotted 3-dimensional objects are algebraic. For a detailed discussion regarding conjectures and facts, see [Ga2]. For a quick explanation of the algebraicity in dimension 3, see Section 3.1 below.

1.4. Examples. In this section, we illustrate Theorem 3 for the complement of the figure eight knot 4_1 , and the complement of the 5_2 knot.

Example 1.4. Consider the complement M of the figure eight knot 4_1 with a meridian-longitude system (μ, λ) . The non-abelian Reidemeister torsion (with respect to the longitude λ) on the character variety X_M is given by (see [Po] or [Db]):

$$\tau_\lambda = \sqrt{17 + 4 \text{tr}_\lambda}.$$

with the convention that we choose the positive square root near the discrete faithful representation ρ_0 with $\text{tr}_\lambda(\rho_0) = -2$. Thus $T_\lambda(\tau_\lambda, \text{tr}_\lambda) = 0$ where

$$T_\lambda(\tau, y) = 17 + 4y - \tau^2.$$

The A -polynomial of the figure eight knot is given by (see [CCGLS]):

$$A(\epsilon_\lambda, e_\mu) = -2 + (e_\mu^4 + e_\mu^{-4}) - (e_\mu^2 + e_\mu^{-2}) + (e_\lambda^2 + e_\lambda^{-2}).$$

Given that $\text{tr}_\lambda = e_\lambda + e_\lambda^{-1}$, $\text{tr}_\mu = e_\mu + e_\mu^{-1}$, the vanishing of the A -polynomial implies that

$$\text{tr}_\lambda = \text{tr}_\mu^4 - 5 \text{tr}_\mu^2 + 2.$$

For details, see [DHY]. On the other hand, the torsion with respect to the meridian is given by (see Equation (16)):

$$\tau_\mu = \tau_\lambda \cdot \left(\frac{\text{tr}_\lambda^2 - 4}{\text{tr}_\mu^2 - 4} \right)^{1/2} \cdot \frac{\partial \text{tr}_\mu}{\partial \text{tr}_\lambda} = \frac{1}{2} \sqrt{(\text{tr}_\mu^2 - 5)(\text{tr}_\mu^2 - 1)}.$$

Thus $T_\mu(\tau_\mu, \text{tr}_\mu) = 0$ where

$$T_\mu(\tau, z) = -5 + 6z^2 - z^4 + 4\tau^2.$$

At the discrete faithful representation ρ_0 , we have $\text{tr}_\lambda(\rho_0) = -2$ [Ca, Cor.2.4] and $\text{tr}_\mu(\rho_0) = \pm 2$ giving that

$$\tau_\lambda(\rho_0) = 3, \quad \tau_\mu(\rho_0) = i\sqrt{3}.$$

Example 1.5. We will repeat the previous example for the twist knot 5_2 . The non-abelian Reidemeister torsion (with respect to the longitude λ) for 5_2 is given by (see [DHY]):

$$\tau_\lambda = (5 \operatorname{tr}_\mu^4 - 37 \operatorname{tr}_\mu^2 + 36) u + (7 - 5 \operatorname{tr}_\mu^2) u^2,$$

where u satisfies the polynomial equation

$$(2 \operatorname{tr}_\mu^2 - 9) - (\operatorname{tr}_\mu^4 - 7 \operatorname{tr}_\mu^2 + 14) u + (2 \operatorname{tr}_\mu^2 - 9) u^2 - u^3 = 0.$$

The selected root that we choose near the discrete faithful representation is $u = \frac{4}{\mathfrak{c}-2}$ where \mathfrak{c} is the cusp-shape; see [DHY, Eqn.(3.9)]. Eliminating u from the above equations, it follows that $T_\lambda(\tau_\lambda, \operatorname{tr}_\mu) = 0$ where

$$\begin{aligned} T_\lambda(\tau, y) &= \tau^3 + \tau^2(-47 + 14y^2 - 5y^4) \\ &\quad + \tau(-5138 + 10057y^2 - 7830y^4 + 3213y^6 - 640y^8 + 50y^{10}) \\ &\quad - 120447 + 339345y^2 - 371691y^4 + 203917y^6 \\ &\quad - 60090y^8 + 8850y^{10} - 500y^{12}. \end{aligned}$$

At the discrete faithful representation ρ_0 , we have $\operatorname{tr}_\mu(\rho_0) = \pm 2$ giving that $\tau = \tau_\lambda(\rho_0)$ is the root of the equation

$$\tau^3 - 71\tau^2 + 2802\tau - 28075 = 0, \quad \tau = 28.4932\dots + i 34.5189\dots$$

1.5. Problems. In this section we list a few problems and future directions.

Problem 1.6. Is the T_λ -polynomial of a hyperbolic knot nontrivial?

Remark 1.7. The volume and the Reidemeister torsion appear as the *classical* and *semiclassical* limit in a parametrized version of the Volume Conjecture; see for example [GM]. Physics arguments suggest that the non-commutative A -polynomial and the Reidemeister torsion is determined by the A -polynomial and the volume of the manifold alone. However, computations with twist knots suggest that the A and T_λ -polynomials seem to be independent from each other. Perhaps this discrepancy can be explained by the difference between on-shell and off-shell physics computations.

Let us now formulate a speculation regarding the *Parametrized Volume Conjecture* of Gukov-Murakami and Le-Garoufalidis; see [GM, GL2]. If K is a knot in S^3 , let $J_{K,N}(q) \in \mathbb{Q}[q^{\pm 1}]$ denote the quantum group invariant of K colored by the N -dimensional irreducible representation of \mathfrak{sl}_2 , and normalized to be 1 at the unknot. For fixed $\alpha \in \mathbb{C}$, the Parametrized Volume Conjecture studies the asymptotics of the sequence $(J_{K,N}(e^{\alpha/N}))$ for $N = 1, 2, \dots$. For suitable α near $2\pi i$, and for hyperbolic knots K , one expects an asymptotic expansion of the form

$$J_{K,N}(e^{\alpha/N}) \sim e^{\frac{NCS(\rho_\alpha)}{2\pi i}} N^{3/2} c_0(\alpha) \left(1 + \sum_{k=1}^{\infty} \frac{c_k(\alpha)}{N^k} \right)$$

where $\rho_\alpha \in X_M$ denotes a representation near ρ_0 with $\operatorname{tr}_\mu(\rho_\alpha) = e^\alpha + e^{-\alpha}$; see [DGLZ, GL2].

Problem 1.8. For every k , and with suitable normalization, show that $c_k(\alpha)$ are germs of unique elements of the field $\mathbb{Q}(X_M)$.

Our next problem concerns the extension of Theorem 2 to simple complex Lie groups $G_{\mathbb{C}}$, rather than $\operatorname{SL}(2, \mathbb{C})$. Physics arguments regarding the 1-loop computation of *perturbative Chern-Simons theory* suggest that an extension of Theorem 2 to arbitrary complex simple groups $G_{\mathbb{C}}$ is possible. It is reasonable to expect that an extension of the non-abelian Reidemeister torsion is possible (see for example [BH1, BH2]), and that Theorem 2 extends.

Problem 1.9. Extend Theorem 2 to arbitrary simple complex Lie groups $G_{\mathbb{C}}$.

2. THE CHARACTER VARIETY OF HYPERBOLIC 3-DIMENSIONAL MANIFOLDS

2.1. Four favors of the character variety, après Dunfield. The careful reader may observe that the volume function is defined for $\mathrm{PSL}(2, \mathbb{C})$ representations of a 1-cusped hyperbolic manifold M , whereas the Reidemeister torsion is defined for $\mathrm{SL}(2, \mathbb{C})$ -representations of M . The proof of Theorem 2 requires a new variant of a representation, the so-called *augmented representation* that comes in two flavors: the $\mathrm{PSL}(2, \mathbb{C})$ and the $\mathrm{SL}(2, \mathbb{C})$ one. For an excellent discussion, we refer the reader to [Dn, Sec.2-3] and [BDR-V, Sec.10]. Much of the results of this section the second author learnt from N. Dunfield, whom we thank for his guidance. Naturally, we are responsible for any comprehension errors.

Let us define the four versions of the character variety of M . Let $R(M, \mathrm{SL}(2, \mathbb{C}))$ denote the set of all homomorphisms of $\pi_1(M)$ into $\mathrm{SL}(2, \mathbb{C})$ and let $X_{M, \mathrm{SL}(2, \mathbb{C})} = R(M, \mathrm{SL}(2, \mathbb{C})) / \mathrm{SL}(2, \mathbb{C})$ denote the algebro-geometric quotient, where $\mathrm{SL}(2, \mathbb{C})$ acts by conjugation. It is easy to see that $R(M, \mathrm{SL}(2, \mathbb{C}))$ and $X_{M, \mathrm{SL}(2, \mathbb{C})}$ are affine varieties defined over \mathbb{Q} .

Let $\overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ denote the subvariety of $R(M, \mathrm{SL}(2, \mathbb{C})) \times P^1(\mathbb{C})$ consisting of pairs (ρ, z) where z is a fixed point of $\rho(\pi_1(\partial M))$. Let $\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})}$ denote the algebro-geometric quotient of $\overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ under the diagonal action of $\mathrm{SL}(2, \mathbb{C})$ by conjugation and Möbius transformations respectively. We will call elements $(\rho, z) \in \overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ *augmented representations*. Their images in the augmented character variety $\overline{X}(M, \mathrm{SL}(2, \mathbb{C}))$ will be called *augmented characters* and will be denoted by square brackets $[(\rho, z)]$. Likewise, replacing $\mathrm{SL}(2, \mathbb{C})$ by $\mathrm{PSL}(2, \mathbb{C})$, we can define the character variety $X_{M, \mathrm{PSL}(2, \mathbb{C})}$ and its augmented version $\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$.

The advantage of the augmented character variety $\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})}$ is that given $\gamma \in \pi_1(\partial M)$ there is a regular function e_γ which sends $[(\rho, z)]$ to the eigenvalue of $\rho(\gamma)$ corresponding to z . In contrast, in $X_{M, \mathrm{SL}(2, \mathbb{C})}$ only the trace $e_\gamma + e_\gamma^{-1}$ of $\rho(\gamma)$ is well-defined. Likewise, in $\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$ (resp. $X_{M, \mathrm{SL}(2, \mathbb{C})}$) only e_γ^2 (resp. $e_\gamma^2 + e_\gamma^{-2}$) is defined.

From now on, we will restrict to a geometric component of the $\mathrm{PSL}(2, \mathbb{C})$ character variety of M and its lifts. The four character varieties associated to M fit in a commutative diagram

$$(4) \quad \begin{array}{ccc} \overline{X}_{M, \mathrm{SL}(2, \mathbb{C})} & \longrightarrow & \overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})} \\ \downarrow & & \downarrow \\ X_{M, \mathrm{SL}(2, \mathbb{C})} & \longrightarrow & X_{M, \mathrm{PSL}(2, \mathbb{C})} \end{array}$$

where the vertical maps are forgetful maps $[(\rho, z)] \longrightarrow [\rho]$ and the horizontal maps are induced by the projection $\mathrm{SL}(2, \mathbb{C}) \longrightarrow \mathrm{PSL}(2, \mathbb{C})$. The vertical maps are generically 2:1 at the geometric components. The horizontal maps are discussed in [Dn, Cor.3.2].

The notation X_M of Section 1 matches the notation $X_M = X_{M, \mathrm{SL}(2, \mathbb{C})}$ of this section.

The next lemma describes the coordinates rings of the four versions of the character variety.

- Lemma 2.1.** (1) *The coordinate ring of $X_{M, \mathrm{SL}(2, \mathbb{C})}$ is generated by tr_γ for all $\gamma \in \pi_1(M)$.*
(2) *The coordinate ring of $X_{M, \mathrm{PSL}(2, \mathbb{C})}$ is generated by tr_γ^2 for all $\gamma \in \pi_1(M)$.*
(3) *The coordinate ring of $\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})}$ is generated by tr_γ for all $\gamma \in \pi_1(M)$ and by e_γ for $\gamma \in \pi_1(\partial M)$.*
(4) *The coordinate ring of $\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$ is generated by tr_γ^2 for all $\gamma \in \pi_1(M)$ and by e_γ^2 for $\gamma \in \pi_1(\partial M)$.*

The commutative diagram (4) gives an inclusion of fields of rational functions:

$$(5) \quad \begin{array}{ccc} \mathbb{Q}(\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})}) & \longleftarrow & \mathbb{Q}(\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}) \\ \uparrow & & \uparrow \\ \mathbb{Q}(X_{M, \mathrm{SL}(2, \mathbb{C})}) & \longleftarrow & \mathbb{Q}(X_{M, \mathrm{PSL}(2, \mathbb{C})}) \end{array}$$

where the horizontal extensions are of degree 2.

2.2. The coefficient field of augmented representations. A crucial part in the proof of Theorem 2 is the choice of a coefficient field of an $\mathrm{SL}(2, \mathbb{C})$ -representation of $\pi_1(M)$. In this section, we show that the notion of an augmented representation solves fits well with the choice of a coefficient field.

First, let us describe the problem. Given a subgroup Γ of $\mathrm{SL}(2, \mathbb{C})$, we can define its *trace field* $\mathbb{Q}(\Gamma)$ (resp. its *coefficient field* $E(\Gamma)$) by $\mathbb{Q}(\mathrm{tr}(A) \mid A \in \Gamma)$ (resp. the field generated over \mathbb{Q} by the entries of all elements A of Γ). The trace field but *not* the coefficient field of Γ is obviously invariant under conjugation of Γ in $\mathrm{SL}(2, \mathbb{C})$. In general, it is not possible to choose a conjugate of Γ to be a subgroup of $\mathrm{SL}(2, \mathbb{Q}(\Gamma))$. Nonetheless, the following lemma shows that not all is lost.

Lemma 2.2. ([Ma, Prop. 3.3][MR, Cor. 3.2.4]) *If Γ is non-elementary, then Γ is conjugate to $\mathrm{SL}(2, K)$ where $K = \mathbb{Q}(\Gamma)(e)$ is an extension of degree $[K : \mathbb{Q}(\Gamma)] \leq 2$, and e can be chosen to be an eigenvalue of a loxodromic element of Γ .*

For the definition of a *non-elementary* subgroup of $\mathrm{SL}(2, \mathbb{C})$ and of a *loxodromic* element, see [Ma, MR]. The proof of the lemma uses the theory of 4-dimensional *quaternion algebras*.

We want to apply Lemma 2.2 to a representation $\rho \in R(M, \mathrm{SL}(2, \mathbb{C}))$. Recall that the discrete faithful representation ρ_0 of $\pi_1(M)$ is non-elementary, and that the subset of characters of elementary representations in the geometric component $X_{M, \mathrm{SL}(2, \mathbb{C})}$ is Zariski closed, and therefore, finite; see [MR].

Given a representation $\rho \in R(M, \mathrm{SL}(2, \mathbb{C}))$, let $\mathbb{Q}(\rho)$ and $E(\rho)$ denote the trace field and the coefficient field of the subgroup $\rho(\pi_1(M)) \subset \mathrm{SL}(2, \mathbb{C})$ respectively. Likewise, if $(\rho, z) \in \overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ is an augmented representation, let $\mathbb{Q}(\rho, z)$ denote the field generated over \mathbb{Q} by $\mathrm{tr}_g(\rho)$ for $g \in \pi_1(M)$ and e_γ for $\gamma \in \pi_1(\partial M)$. Similarly, we define the coefficient field $E(\rho, z)$ associated to the augmented representation (ρ, z) .

The next lemma follows from lemma 2.2 and the above discussion.

Lemma 2.3. (1) *If $\rho \in R(M, \mathrm{SL}(2, \mathbb{C}))$ is generic (i.e., non-elementary) then a conjugate of ρ is defined over a quadratic extension of $\mathbb{Q}(\rho)$.*
 (2) *If $(\rho, z) \in \overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ is generic (i.e., non-elementary) then there exists $N \in \mathrm{SL}(2, \mathbb{C})$ so that $N^{-1}\rho N$ is defined over $E(\rho, z)$.*
 (3) *There exists $N \in \mathrm{SL}(2, \mathbb{C})$ such that if (ρ, z) is near the discrete faithful representation (ρ_0, z_0) , then $N^{-1}(\rho, z)N$ is defined over $E(\rho, z)$.*

An alternative version of the above Lemma is possible; see Lemma 2.6 below.

2.3. Augmented representations and the shape field. There is an alternative description of the field $\mathbb{Q}(\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})})$ in terms of shape parameters of ideal triangulations of M , which is useful in applications. For completeness, we discuss it in this section and the next. Let us first describe $\overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$ in terms of *pseudo-developing maps*, discussed in detail in [Dn, Sec.2.5]. Given $\rho \in R_{M, \mathrm{PSL}(2, \mathbb{C})}$, consider a ρ -equivariant map $\widetilde{M} \rightarrow \mathbb{H}^3$, where \mathbb{H}^3 denotes the 3-dimensional hyperbolic space. Since ∂M is a 2-torus, it lifts to a disjoint collection of planes \mathbb{R}^2 in the universal cover \widetilde{M} . Let \overline{M} denote the space obtained by cutting \widetilde{M} along these planes, and crushing them into points. Set-theoretically, the set $\overline{M} \setminus \widetilde{M}$ of ideal points is in 1-1 correspondence with the *cusps* of M in \mathbb{H}^3 , i.e., with the coset $\pi_1(M)/\pi_1(\partial M)$. An augmented representation $(\rho, z) \in \overline{R}_{M, \mathrm{PSL}(2, \mathbb{C})}$ gives a $\pi_1(M)$ -equivariant map

$$D_{(\rho, z)} : \overline{M} \rightarrow \overline{\mathbb{H}^3}$$

where $\overline{\mathbb{H}^3} = \mathbb{H}^3 \cup \mathbb{CP}^1$ is the compactification of hyperbolic space by adding a sphere \mathbb{CP}^1 at infinity. Such a map is a pseudo-developing map in [Dn, Sec.2.5]. An augmented character $[(\rho, z)] \in \overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$ does not have a unique pseudo-developing map, however every two are homotopic, for example using a straight line homotopy $tf(x) + (1-t)g(x)$ in \mathbb{H}^3 . Thus, there is a well-defined map:

$$(6) \quad \overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})} \rightarrow \{\text{Pseudo-developing maps of } M, \text{ modulo homotopy rel boundary}\}$$

Consider a 4-tuple of distinct points $(A, B, C, D) \in (\overline{M} \setminus \widetilde{M})^4$, and an augmented character $[(\rho, z)] \in \overline{X}_{M, \mathrm{PSL}(2, \mathbb{C})}$. Then, $D_{[(\rho, z)]}$ sends A, B, C, D to four points A', B', C', D' in $\mathbb{C} \cup \{\infty\} = \mathbb{CP}^1 = \partial\mathbb{H}^3$, and

consider their *cross-ratio*

$$cr_{A,B,C,D}[(\rho, z)] = \frac{(A' - D')(B' - C')}{(A' - C')(B' - D')}.$$

If A', B', C', D' are distinct, then $cr_{A,B,C,D}[(\rho, z)] \in \mathbb{C}$, else $cr_{A,B,C,D}[(\rho, z)]$ is undefined. This gives a rational map

$$cr_{A,B,C,D} : \overline{X}_{M, \text{PSL}(2, \mathbb{C})} \longrightarrow \mathbb{C}.$$

Let \mathbb{Q}^{dev} denote the field over \mathbb{Q} generated by $cr_{A,B,C,D}$ for all 4-tuples of distinct points of $\overline{M} \setminus \widetilde{M}$.

Lemma 2.4. *We have*

$$\mathbb{Q}^{\text{dev}} = \mathbb{Q}(\overline{X}_{M, \text{PSL}(2, \mathbb{C})}).$$

The proof will be given in the next section.

2.4. Ideal triangulations and the gluing equations variety. A convenient way to construct the unique hyperbolic structure on M , and its small incomplete hyperbolic deformations is using an *ideal triangulation* $\mathcal{T} = (\mathcal{T}_1, \dots, \mathcal{T}_s)$ of M which recovers the complete hyperbolic structure. For a detailed description of ideal triangulations, see [BP] and also [BDR-V, App.10]. An ideal triangulation \mathcal{T} which is compatible with the discrete faithful representation has nondegenerate shape parameters $z_j \in \mathbb{C} \setminus \{0, 1\}$ for $j = 1, \dots, s$. Such a triangulation always exists; for example subdivide the canonical Epstein-Penner decomposition of M by adding ideal triangles; see [EP, BP, PP]. Once we choose shape parameters for each ideal tetrahedron, one can use them to give a hyperbolic metric (in general incomplete) in the universal cover \widetilde{M} , once a compatibility condition along the edges of \mathcal{T} is satisfied. This compatibility condition defines the so-called *Gluing Equations* variety $\mathcal{G}(\mathcal{T})$. In the appendix of [BDR-V], Dunfield describes a map

$$(7) \quad \mathcal{G}(\mathcal{T}) \longrightarrow \overline{R}_{M, \text{PSL}(2, \mathbb{C})}$$

which projects to an injection

$$(8) \quad \mathcal{G}(\mathcal{T}) \longrightarrow \overline{X}_{M, \text{PSL}(2, \mathbb{C})}$$

Consider the field $\mathbb{Q}(z_1, \dots, z_s)$ over \mathbb{Q} generated by the shape parameters z_1, \dots, z_s . A priori, $\mathbb{Q}(z_1, \dots, z_s)$ depends on M . The next lemma describes the fields of rational functions of augmented representations in terms of the shape field.

Lemma 2.5. *We have*

$$(9) \quad \mathbb{Q}(\overline{X}_{M, \text{PSL}(2, \mathbb{C})}) = \mathbb{Q}(z_1, \dots, z_s)$$

and

$$(10) \quad \mathbb{Q}(\overline{X}_{M, \text{SL}(2, \mathbb{C})}) = \mathbb{Q}(z_1, \dots, z_s, e_\lambda, e_\mu)$$

Proof. The shape parameters z_j for $j = 1, \dots, s$ are coordinate functions on the curve $\mathcal{G}(\mathcal{T})$. In addition, the squares e_λ^2 and e_μ^2 of the eigenvalues of a meridian-longitude pair (λ, μ) of ∂M are rational functions of the shape parameters z_j . Since the map in Equation (8) is an inclusion of a curve into another, it follows that their fields of rational functions are equal. This proves Equation (9).

Equation (10) follows from Lemma 2.3 and the fact that $e_\lambda^2, e_\mu^2 \in \mathbb{Q}(z_1, \dots, z_s)$. \square

As a consequence, Lemma 2.4 follows verbatim by the proof of [MR, Lem.5.5.2]. \square

Let us end this section with an alternative version of Lemma 2.3 using shape fields. Recall from [Dn, Sec.2] that the map in Equation (7) can be defined as follows. Fix a solution (z_1, \dots, z_s) of the Gluing Equations of \mathcal{T} . Lift \mathcal{T} to an ideal triangulation of \widetilde{M} , and then map the lift of one ideal tetrahedron to a fixed ideal tetrahedron of \mathbb{H}^3 of the same shape, and then use $\pi_1(M)$ -equivariance to send every other ideal tetrahedron to an appropriate ideal tetrahedron of \mathbb{H}^3 , using face-pairings. There is a consistency condition, which is satisfied since we are using a solution to the Gluing Equations. This defines a developing map and

a corresponding $\mathrm{PSL}(2, \mathbb{C})$ -representation ρ . In [BDR-V, App. 10], Dunfield describes how to define not only a representation in $\mathrm{PSL}(2, \mathbb{C})$, but also an augmented one (ρ, z) .

The combinatorial structure of \mathcal{T} gives a presentation of $\Pi = \pi_1(M)$ in terms of *face-pairings*:

$$(11) \quad \Pi = \langle g_1, \dots, g_s \mid r_1, \dots, r_{s-1} \rangle.$$

Each generator of Π is represented by a path in the 1-skeleton of the dual triangulation of \mathcal{T} ; see [MR, Chap. 5] or [R, Ch.11]. The entries of $\rho(g_j)$, for $j = 1, \dots, s$, are given by face-pairings, and are explicit matrices with entries in $\mathbb{Q}(z_1, \dots, z_s)$; see [MR, Chap. 5]. The above discussion proves the following version of Lemma 2.3.

Lemma 2.6. (1) *The image of the map in Equation (7) is defined over $\mathbb{Q}(z_1, \dots, z_s)$.*
 (2) *Generically, a lift of the image of the map in Equation (7) to $\overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ is defined over $\mathbb{Q}(z_1, \dots, z_s, e_\lambda, e_\mu)$.*

3. THE NON-ABELIAN REIDEMEISTER TORSION

3.1. An explanation of the rationality of the Reidemeister torsion in dimension 3. Before we prove the rationality of the torsion stated in Theorem 2, let us give the main idea which is rather simple, and defer the technical details for the next section.

The starting point is a hyperbolic manifold M with one cusp. The character variety $\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})}$ depends only on $\pi_1(M)$ but we view it in a specific birational equivalent way by using a combinatorial decomposition of M into ideal tetrahedra. Every such manifold is obtained by a combinatorial face-pairing of a finite collection \mathcal{T} of nondegenerate (but perhaps flat, or negatively oriented) ideal tetrahedra $\mathcal{T}_1, \dots, \mathcal{T}_s$. The hyperbolic shape of a nondegenerate ideal tetrahedron is determined by a complex number $z \in \mathbb{C} \setminus \{0, 1\}$, up to the action of a finite group of order 6. The discrete faithful representation ρ_0 assigns hyperbolic shapes z_j to the tetrahedra \mathcal{T}_j for $j = 1, \dots, s$. These shapes satisfy the so-called *Gluing Equations*, which is a collection of polynomial equations in z_j and $1 - z_j$ to make the metric match along the edges of the ideal tetrahedra. The Gluing Equations define a variety $\mathcal{G}(\mathcal{T})$ which of course depends on \mathcal{T} . When the discrete faithful representation ρ_0 slightly deforms in ρ_t (i.e., bends, in the language of Thurston) this causes the shapes z_j of \mathcal{T}_j to deform to $z_j(t)$. For small enough t , the new shapes still satisfy the Gluing Equations. Consequently, for every t , the shapes $z_j(t)$ for $j = 1, \dots, s$ are algebraically dependent, and so is any algebraic function of the shapes.

In the case of the A -polynomial, the squares $e_\lambda(t)^2$ and $e_\mu(t)^2$ of the eigenvalues $e_\lambda(t)$ and $e_\mu(t)$ of a meridian-longitude pair of $T^2 = \partial M$ are rational functions in $z_j(t)$ (in fact, monomials in $z_j(t)$ and $1 - z_j(t)$ with integer exponents), thus $(e_\lambda(t), e_\mu(t))$ are algebraically dependent. This dependence defines the A -polynomial.

In the case of Reidemeister torsion and Theorem 2, the torsion $\tau_\mu(\rho_t)$ of the relevant chain complex is defined over the field $\mathbb{Q}(z_1(t), \dots, z_s(t), e_\lambda(t), e_\mu(t))$. In other words all matrices that compute the torsion (and thus the ratios of their determinants) have entries in the field $\mathbb{Q}(z_1(t), \dots, z_s(t), e_\lambda(t), e_\mu(t))$.

3.2. Proof of Theorem 2. In this section, we will prove Theorem 2. Let M be a one-cusp finite-volume complete hyperbolic 3-manifold. Choose an ideal triangulation $\mathcal{T} = (\mathcal{T}_1, \dots, \mathcal{T}_s)$ compatible with the discrete faithful representation of M as described above, and let (z_1, \dots, z_s) denote the shape parameters of \mathcal{T} . Let E denote the following field:

$$\mathbb{K} = \mathbb{Q}(z_1, \dots, z_s, e_\lambda, e_\mu) = \mathbb{Q}(\overline{X}_{M, \mathrm{SL}(2, \mathbb{C})})$$

where the last equality follows from Lemma 2.5.

Let J denote an open interval in \mathbb{R} that contains 0, and consider a 1-parameter family $t \in J \mapsto z(t) = (z_1(t), \dots, z_s(t)) \in \mathcal{G}(\mathcal{T})$ of solutions of the Gluing Equations, with image $(\rho'_t, z'_t) \in \overline{R}(M, \mathrm{PSL}(2, \mathbb{C}))$ under the map in Equation (7) and with lift $(\rho_t, z_t) \in \overline{R}(M, \mathrm{SL}(2, \mathbb{C}))$ where ρ_0 is a lift to $\mathrm{SL}(2, \mathbb{C})$ of the discrete faithful representation of M . Fix γ an essential curve in the boundary torus ∂M .

We will explain how to define the Reidemeister torsion $\tau_\gamma(\rho_t)$ (for complete definitions the reader can refer to Porti's monograph [Po] and to Turaev's book [T]), and why it coincides with the evaluation of an element of \mathbb{K} at ρ_t .

The 2-skeleton of the combinatorial dual W to \mathcal{T} is a 2-dimensional CW -complex which is a spine of M ; see [BP]. Mostow rigidity Theorem implies that every homotopy equivalence of M is homotopic to a homeomorphism (even to an isometry), and Chapman's theorem concludes that every homotopy equivalence of M is simple; [Co]. Thus, W is simple homotopy equivalent to M , and we can use W to compute $\tau_\gamma(\rho_t)$. The ideas of the definition of the non-abelian torsion $\tau_\gamma(\rho_t)$ are the following:

- (a) Consider the universal cover \widetilde{W} of W and the integral chain complex $C_*(\widetilde{W}; \mathbb{Z})$ of \widetilde{W} for $* = 0, 1, 2$. The fundamental group $\Pi = \pi_1(W) = \pi_1(M)$ acts on \widetilde{W} by covering transformations. This action turns the complex $C_*(\widetilde{W}; \mathbb{Z})$ into a $\mathbb{Z}[\Pi]$ -module. The Lie algebra $\mathfrak{sl}_2(\mathbb{C})$ also can be viewed as a $\mathbb{Z}[\Pi]$ -module by using the composition $Ad \circ \rho_t$, where Ad denotes the adjoint representation of $\mathfrak{sl}_2(\mathbb{C})$. We let $\mathfrak{sl}_2(\mathbb{C})_{\rho_t}$ denote this $\mathbb{Z}[\Pi]$ -module. The *twisted chain complex* of W is the \mathbb{C} -vector space:

$$(12) \quad C_*^{\rho_t} = C_*(\widetilde{W}; \mathbb{Z}) \otimes_{\mathbb{Z}[\Pi]} \mathfrak{sl}_2(\mathbb{C})_{\rho_t}.$$

- (b) The twisted chain complex $C_*^{\rho_t}$ computes the so-called *twisted homology* of W which is denoted by $H_*^{\rho_t}$. The betti numbers of $H_*^{\rho_t}$ are given by (because ρ_t lies in a neighborhood of the discrete and faithful representation and thus is regular, see [Po, Chap. 3]):

$$\dim_{\mathbb{C}}(H_0^{\rho_t}) = 0, \quad \dim_{\mathbb{C}}(H_1^{\rho_t}) = 1, \quad \dim_{\mathbb{C}}(H_2^{\rho_t}) = 1.$$

- (c) For $i = 1, 2$ construct elements \mathbf{h}_i^t in $C_i^{\rho_t}$, which project to bases of the twisted homology groups $H_i^{\rho_t}$.
- (d) Then, the torsion $\tau_\gamma(\rho_t)$ is an explicit ratio of determinants; see [Db] or [Po, Chap. 3] and Equation (15) below.

We now give the details of the definition of the non-abelian Reidemeister torsion and prove Theorem 2. To clarify the presentation, suppose that V_t is a 1-parameter family of \mathbb{C} -vector spaces for $t \in J$. We will say that V_t is defined over \mathbb{K} if there exists a vector space $V_{\mathbb{K}}$ over \mathbb{Q} such that $V_t = (V_{\mathbb{K}} \otimes_{\mathbb{Q}} E(\rho_t, z_t)) \otimes_{\mathbb{Q}} \mathbb{C}$ for all $t \in J$, where $E(\rho_t, z_t)$ is the coefficient field of (ρ_t, z_t) , defined in Section 2.2. Likewise, a 1-parameter family of \mathbb{C} -linear transformations $T_t \in \text{Hom}_{\mathbb{C}}(V_t, W_t)$ is defined over \mathbb{K} if $T \in \text{Hom}_{\mathbb{Q}}(V_{\mathbb{K}}, W_{\mathbb{K}}) \otimes_{\mathbb{Q}} \mathbb{C}$. In concrete terms, a 1-parameter family of matrices (resp. vectors) is defined over E if its entries (resp. coordinates) lie in \mathbb{K} .

Lemma 2.6 implies the following.

Claim 3.1. *The 1-parameter family (ρ_t, z_t) ($t \in J$) is defined over \mathbb{K} .*

Consider the presentation Π in Equation (11) of $\pi_1(M)$ given by face-pairings. A coordinate description of the chain complex $C_*^{\rho_t}$ is given by (see [Db])

$$0 \longrightarrow \mathfrak{sl}_2(\mathbb{C})^{s-1} \xrightarrow{d_2^{\rho_t}} \mathfrak{sl}_2(\mathbb{C})^s \xrightarrow{d_1^{\rho_t}} \mathfrak{sl}_2(\mathbb{C}) \longrightarrow 0$$

for $* = 0, 1, 2$ where the boundary operators are given by

$$d_1^{\rho_t}(x_1, \dots, x_s) = \sum_{j=1}^s (1 - g_j) \circ x_j, \quad \text{and} \quad d_2^{\rho_t}(x_1, \dots, x_{s-1}) = \left(\sum_{j=1}^{s-1} \frac{\partial r_j}{\partial g_k} \circ x_j \right)_{1 \leq k \leq s}.$$

Here $g \circ x = Ad_{\rho_t(g)}(x)$ and $\frac{\partial r_j}{\partial g_k}$ denotes the *Fox derivative* of r_j with respect to g_k . The above description of $C_*^{\rho_t}$ and Claim 3.1 imply the following.

Claim 3.2. *The 1-parameter family $C_*^{\rho_t}$ ($t \in J$) is defined over \mathbb{K} .*

Next, we construct a 1-parameter family of basing elements \mathbf{h}_i^t for $i = 1, 2$ and show that it is defined over \mathbb{K} . Let $\{e_1^{(i)}, \dots, e_{n_i}^{(i)}\}$ be the set of i -dimensional cells of W . We lift them to the universal cover and we choose an arbitrary order and an arbitrary orientation for the cells $\{\tilde{e}_1^{(i)}, \dots, \tilde{e}_{n_i}^{(i)}\}$. If $\mathcal{B} = \{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$ is an orthonormal basis of $\mathfrak{sl}_2(\mathbb{C})$, then we consider the corresponding (geometric) basis over \mathbb{C} :

$$\mathbf{c}_{\mathcal{B}}^i = \left\{ \tilde{e}_1^{(i)} \otimes \mathbf{a}, \tilde{e}_1^{(i)} \otimes \mathbf{b}, \tilde{e}_1^{(i)} \otimes \mathbf{c}, \dots, \tilde{e}_{n_i}^{(i)} \otimes \mathbf{a}, \tilde{e}_{n_i}^{(i)} \otimes \mathbf{b}, \tilde{e}_{n_i}^{(i)} \otimes \mathbf{c} \right\}$$

of $C_i^{\rho_t}$. We fix a generator P^{ρ_t} of $H_0^{\rho_t}(\partial M) \subset C_0^{\rho_t}$ i.e., $P^{\rho_t} \in \mathfrak{sl}_2(\mathbb{C})$ is such that $Ad_{\rho_t(g)}(P^{\rho_t}) = P^{\rho_t}$ for all $g \in \pi_1(\partial M)$.

Claim 3.3. *The 1-parameter family P^{ρ_t} ($t \in J$) is defined over \mathbb{K} .*

Proof. Observe that P^{ρ_t} is a generator of the intersection

$$\ker(Ad_{\rho_t(\mu)} - \mathbf{1}) \cap \ker(Ad_{\rho_t(\lambda)} - \mathbf{1}).$$

Since this family of vector spaces and linear maps is defined over \mathbb{K} (by Claim 3.2), the result follows. \square

The canonical inclusion $j: \partial M \rightarrow M$ induces (see [Po, Corollary 3.23]) an isomorphism

$$j_*: H_2^{\rho_t}(\partial M) \rightarrow H_2^{\rho_t}(M) \simeq H_2^{\rho_t}(W) = \ker d_2^{\rho_t} \subset C_2^{\rho_t}.$$

Moreover, one can prove that (see [Po, Proposition 3.18])

$$H_2^{\rho_t}(\partial M) \cong H_2(\partial M; \mathbb{Z}) \otimes \mathbb{C}.$$

More precisely, let $[[\partial M]] \in H_2(\partial M; \mathbb{Z})$ be the fundamental class induced by the orientation of ∂M , one has $H_2^{\rho_t}(\partial M) = \mathbb{C}[[\partial M]] \otimes P_t^{\rho_t}$. The *reference generator* of $H_2^{\rho_t}(M)$ is defined by

$$(13) \quad \mathbf{h}_2^t = j_*([[\partial M]]) \otimes P_t^{\rho_t} \in C_2^{\rho_t}.$$

Claim 3.3 implies that

Claim 3.4. *The 1-parameter family \mathbf{h}_2^t ($t \in J$) is defined over \mathbb{K} .*

Since ρ_t is near ρ_0 and γ is admissible, the inclusion $\iota: \gamma \rightarrow M$ induces (see [Po, Definition 3.21]) an *isomorphism*

$$\iota^*: H_1^{\rho_t}(\gamma) \rightarrow H_1^{\rho_t}(M) \simeq H_1^{\rho_t}(W) = \ker d_1^{\rho_t} / \text{im } d_2^{\rho_t}.$$

The *reference generator* of the first twisted homology group $H_1^{\rho_t}(M)$ is defined by

$$(14) \quad \mathbf{h}_1^t = \iota_*([\gamma] \otimes P_t^{\rho_t}) \in C_1^{\rho_t}.$$

Claim 3.3 implies that:

Claim 3.5. *The 1-parameter family \mathbf{h}_1^t ($t \in J$) is defined over \mathbb{K} .*

Using the bases described above, the non-abelian Reidemeister torsion of the 1-parameter family ρ_t is defined by:

$$(15) \quad \tau_\gamma(\rho_t) = \text{Tor}(C_*^{\rho_t}(W; \mathfrak{sl}_2(\mathbb{C})_{\rho_t}), \mathbf{c}_{\mathcal{B}}^*, \mathbf{h}_t^*) \in \mathbb{C}^*.$$

The torsion $\tau_\gamma(\rho_t)$ is an invariant of M which is *well defined up to a sign*. Moreover, if ρ_t and $\tilde{\rho}_t$ are two 1-parameter family of representations which pointwise have the same character then $\tau_\gamma(\rho_t) = \tau_\gamma(\tilde{\rho}_t)$. Finally, one can observe that $\tau_\gamma(\rho_t)$ does not depend on the choice of the invariant vector P^{ρ_t} (see [Db]).

The above discussion implies that

Claim 3.6. *For every essential curve $\gamma \in \partial M$, the 1-parameter family $\tau_\gamma(\rho_t)$ ($t \in J$) is defined over \mathbb{K} .*

In other words, there exist $\hat{\tau}_\gamma \in \mathbb{Q}(\overline{X}_{M, \text{SL}(2, \mathbb{C})})$ such that for (ρ_t, z) near (ρ_0, z_0) we have $\tau_\gamma(\rho) = \hat{\tau}_\gamma(\rho_t, z)$. Since the left hand side does not depend on z , it follows from Section 2.1 that $\hat{\tau}_\gamma \in \mathbb{Q}(X_{M, \text{SL}(2, \mathbb{C})})$. This concludes the proof of Theorem 2. \square

3.3. The dependence of the Reidemeister torsion on the admissible curve and the A -polynomial.

In this section, we discuss the dependence of the non-abelian Reidemeister torsion on the admissible curve. Although this discussion is independent of the proof of Theorem 2, it might be useful in other contexts. Recall that the non-abelian Reidemeister torsion is defined in terms of the twisted chain complex in Equation (12) which is not acyclic. Thus, it requires the choice of distinguished bases \mathbf{h}_i for $i = 1, 2$. Such bases can be chosen once an *admissible curve* $\gamma \in \partial M$ is chosen; see [Po, Chap. 3]. Porti proves that for every homotopically non-trivial curve γ in ∂M , the holonomy representation ρ_0 is γ -regular. The same holds for representations ρ near ρ_0 . A well-known application of Thurston's *Hyperbolic Dehn Surgery Theorem* implies that $\rho_0 \in X_M$ is a smooth point of X_M and that a neighborhood U of ρ_0 is parametrized by the polynomial function tr_γ ; see for example [NZ] and [Po, Cor. 3.28]. Choose a meridian-longitude pair (μ, λ) in ∂M , set $\text{tr}_\mu(\rho_i) = e_\mu + e_\mu^{-1}$, $\text{tr}_\lambda(\rho) = e_\lambda + e_\lambda^{-1}$, and consider the A -polynomial $A_M = A_M(e_\mu, e_\lambda) \in \mathbb{Z}[e_\mu^{\pm 1}, e_\lambda^{\pm 1}]$ of M . For a detailed discussion on the A -polynomial of M and its relation to the various views of the character variety, see the appendix of [BDR-V].

With the above notation, Porti proves that the dependence of the torsion on the admissible curve γ is controlled by the A -polynomial. More precisely, we have [Po, Cor. 4.9, Prop. 4.7]:

$$(16) \quad \tau_\mu = \tau_\lambda \cdot \left(\frac{\text{tr}_\lambda^2 - 4}{\text{tr}_\mu^2 - 4} \right)^{1/2} \cdot \frac{\partial \text{tr}_\mu}{\partial \text{tr}_\lambda}$$

$$(17) \quad = \tau_\lambda \cdot (\text{res}^* \circ (\Delta^*)^{-1}) \left(\frac{e_\lambda \partial A_M / \partial e_\lambda}{e_\mu \partial A_M / \partial e_\mu} \right),$$

where $\text{res}^*: X_{M, SL(2, \mathbb{C})} \rightarrow X_{\partial M, SL(2, \mathbb{C})}$ is the restriction-map induced by the usual inclusion $\partial M \hookrightarrow M$, and Δ^* works as follows on the trace field

$$\Delta^*(\text{tr}_\gamma) = e_\gamma + e_\gamma^{-1}.$$

Acknowledgment. A first draft of the paper was discussed during a workshop on the Volume Conjecture in Strasbourg 2007. The authors wish to thank their organizers, S. Baseilhac, F. Costantino and G. Massuyeau for their hospitality, and M. Heusener, R. Kashaev and W. Neumann for enlightening conversations. S.G. wishes to thank N. Dunfield for numerous useful conversations, suggestions and for a careful reading of a first draft.

REFERENCES

- [APS] M.F. Atiyah, V.K. Patodi and I.M. Singer, *Spectral asymmetry and Riemannian geometry I-III*, Math. Proc. Cambridge Philos. Soc. **77** (1975) 43–69, *ibid* **78** (1975) 405–432 *ibid* **79** (1976) 71–99.
- [BP] R. Benedetti and C. Petronio, *Lectures on hyperbolic geometry*, Universitext. Springer-Verlag (1992).
- [Bo] D.W. Boyd, *Mahler's measure and invariants of hyperbolic manifolds*, in Number Theory for the Millenium (M.A. Bennett et al. ed.) A.K. Peters Boston (2001).
- [BDR-V] ———, N.M. Dunfield and F. Rodriguez-Villegas, *Mahler's Measure and the Dilogarithm (II)*, preprint 2003 [arXiv:0308041](https://arxiv.org/abs/0308041).
- [BZ] S. Boyer and X. Zhang, *Every nontrivial knot in S^3 has nontrivial A -polynomial*, to appear in Proc. Amer. Math. Soc.
- [BH1] D. Burghelena and S. Haller, *Complex-valued Ray-Singer torsion*, J. Funct. Anal. **248** (2007) 27–78.
- [BH2] ——— and ———, *Torsion, as a function on the space of representations*, in \mathbb{C}^* -algebras and elliptic theory II, Trends Math., Birkhäuser (2008) 41–66.
- [Ca] D. Calegari, *Real places and torus bundles*, Geom. Dedicata **118** (2006) 209–227.
- [Ch] A. Champanerkar, *A -polynomials and Bloch invariants of hyperbolic 3-manifolds*, thesis, Columbia University 2003.
- [Co] M. Cohen, *A course in simple-homotopy theory*, Graduate Texts in Mathematics **10**, Springer-Verlag (1973).
- [CCGLS] D. Cooper, M. Culler, H. Gillet, D. Long and P. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118** (1994) 47–84.
- [Cu] M. Culler, *Lifting representations to covering groups*, Adv. in Math. **59** (1986) 64–70.
- [DGLZ] T. Dimofte, S. Gukov, J. Lenells and D. Zagier, *Exact Results for Perturbative Chern-Simons Theory with Complex Gauge Group*, preprint 2009 [arXiv:0903.2472](https://arxiv.org/abs/0903.2472).
- [Db] J. Dubois, *Non-abelian twisted Reidemeister torsion for fibered knots*, Canad. Math. Bull. **49** (2006) 55–71
- [DHY] J. Dubois, V. Huynh and Y. Yamaguchi, *Non-abelian Reidemeister torsion for twist knots*, J. of Knot Theory and Its Ramifications **18** (2009) 1–39

- [Dn] N. Dunfield, *Cyclic surgery, degrees of maps of character curves, and volume rigidity for hyperbolic manifolds*, Invent. Math. **136** (1999), 623–657.
- [DG] ——— and S. Garoufalidis, *Non-triviality of the A-polynomial for knots in S^3* , Algebr. Geom. Topol. **4** (2004) 1145–1153.
- [EP] D.B. Epstein and R.C. Penner, *Euclidean decompositions of noncompact hyperbolic manifolds*, J. Differential Geom. **27** (1988) 67–80.
- [F] S. Francaviglia, *Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds*, Int. Math. Res. Not. **9** (2004) 425–459.
- [FK] ——— and B. Klaff, *Maximal volume representations are Fuchsian*, Geom. Dedicata **117** (2006) 111–124.
- [GL1] ——— and T.T.Q. Le, *The colored Jones function is q-holonomic*, Geom. and Topology **9** (2005) 1253–1293.
- [GL2] ——— and ———, *Asymptotics of the colored Jones function of a knot*, preprint 2005 math.GT/0508100, to appear in Geom. and Topology.
- [Ga1] ———, *On the characteristic and deformation varieties of a knot*, Proceedings of the CassonFest, Geometry and Topology Monographs **7** (2004) 291–309.
- [Ga2] ———, *Chern-Simons theory, analytic continuation and arithmetic*, Acta Math. Vietnamica, **33** (2008) 335–362.
- [Go] W.M. Goldman, *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. Math. **85** (1986) 263–302.
- [GM] S. Gukov and H. Murakami, $SL(2, \mathbb{C})$ Chern-Simons theory and the asymptotic behavior of the colored Jones polynomial, preprint 2006 arXiv:math/0608324.
- [MR] C. Maclachlan and A.W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Graduate Texts in Mathematics **219** Springer-Verlag (2003).
- [Ma] A.M. Macbeath, *Commensurability of co-compact three-dimensional hyperbolic groups*, Duke Math. J. **50** (1983) 1245–1253.
- [Mi] J. Milnor, *Whitehead torsion*, Bull. AMS **72** (1966) 358–426.
- [NR] W.D. Neumann and A.W. Reid, *Arithmetic of hyperbolic manifolds*, Topology '90, Ohio State Univ. Math. Res. Inst. Publ., **1** (1992) 273–310.
- [NZ] ——— and D. Zagier, *Volumes of hyperbolic three-manifolds*, Topology **24** (1985) 307–332.
- [PP] C. Petronio and J. Porti, *Negatively oriented ideal triangulations and a proof of Thurston's hyperbolic Dehn filling theorem*, Expo. Math. **18** (2000) 1–35.
- [Po] J. Porti, *Torsion de Reidemeister pour les variétés hyperboliques*, Memoirs AMS **612** (1997).
- [R] J.G. Ratcliffe, *Foundations of hyperbolic manifolds*, Graduate Texts in Mathematics **149** Springer-Verlag (1994).
- [Sh] P.B. Shalen, *Representations of 3-manifold groups*, Handbook of Geometric Topology (2002) 955–1044.
- [Th] W. Thurston, *The geometry and topology of 3-manifolds*, 1979 notes, available from MSRI at www.msri.org/publications/books/gt3m/
- [Tu] V. Turaev, *The Yang-Baxter equation and invariants of links*, Inventiones Math. **92** (1988) 527–553.
- [T] V. Turaev, *Torsions of 3-dimensional manifolds*, Progress in Mathematics **208** Birkhuser Verlag (2002).

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PARIS DIDEROT–PARIS 7, UFR DE MATHÉMATIQUES, CASE 7012, BÂTIMENT CHEVALERET, 2, PLACE JUSSIEU, 75205 PARIS CEDEX 13 FRANCE,
<http://www.institut.math.jussieu.fr/~dubois/>
E-mail address: dubois@math.jussieu.fr

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA,
<http://www.math.gatech.edu/~stavros>
E-mail address: stavros@math.gatech.edu