

# Representations of Integers as the Sum of $k$ Terms

**Paul Erdős**

*Hungarian Academy of Sciences, Hungary*

**Prasad Tetali**

*Department of Computer Science, Courant Institute of Mathematical Sciences,  
New York, NY 10012*

## ABSTRACT

A set of natural numbers is called an *asymptotic basis* of order  $k$  if every number (sufficiently large) can be expressed as a sum of  $k$  distinct numbers from the set. In this paper we prove that, for every fixed  $k$ , there exists an asymptotic basis of order  $k$  such that the number of representations of  $n$  is  $\Theta(\log n)$ .

*Key Words: Sidon sequences, representations of integers, random constructions*

## 1. INTRODUCTION

Let  $\mathcal{S}$  be a sequence of natural numbers, and let  $r_k(n)$  denote the number of representations of  $n$  in the form

$$n = a_1 + a_2 + \cdots + a_k, \quad 0 < a_1 < \cdots < a_k (a_i \in \mathcal{S})$$

$\mathcal{S}$  is said to be an *asymptotic basis* of order  $k$  if there exists a natural number  $n_0$  such that  $r_k(n) > 0$  for  $n > n_0$ . In 1932 Sidon raised the following question: Does there exist an asymptotic basis of order 2 such that, for every  $\epsilon > 0$ ,  $\lim_{n \rightarrow \infty} r_2(n)/n^\epsilon = 0$ ? In 1956 Erdős [2] proved that there exist positive constants  $c_1$  and  $c_2$  such that

$$c_1 \log n \leq r_2(n) \leq c_2 \log n \text{ for large } n,$$

which more than answered the question. For a thorough exposition of the above result and several other results of similar nature, the reader is strongly recommended to refer to [5].

In this paper we prove a more general version of Sidon's original question using probabilistic techniques, old and new. Although the following general version was believed to be true, no complete proof appears in the literature.

**Theorem 1.** *For every fixed  $k$  there exists an asymptotic basis of order  $k$  such that  $r_k(n) = \Theta(\log n)$ .*

We shall actually conclude that in a suitable probability (measure) space almost all sequences are of *the right kind*.

## 2. PROBABILISTIC TOOLS

### Disjointness Lemma

Let  $A_1, A_2, \dots$  be events in a probability space. The following elementary lemma is often useful in proving the probability of occurrence of "large" subsets of mutually independent events is "small." We refer to this as the *disjointness lemma* in view of the nature of its application in this paper.

**Lemma 1 (Disjointness Lemma).** *If  $\sum_i \Pr[A_i] \leq \mu$ , then*

$$\sum_{\substack{\{A_1, \dots, A_l\} \\ \text{mutually} \\ \text{independent}}} \Pr[A_1 \wedge A_2 \wedge \dots \wedge A_l] \leq \mu^l / l!$$

*Proof.* Let  $\Sigma'$  denote the summation in the statement of the lemma. Then by mutual independence of the probabilities,

$$\begin{aligned} \sum' \Pr[A_1 \wedge \dots \wedge A_l] &= \sum' \Pr[A_1] \cdots \Pr[A_l] \\ &\leq \sum_{\text{all } \{A_1, \dots, A_l\}} \Pr[A_1] \cdots \Pr[A_l] \\ &= \frac{1}{l!} \sum_{\{A_1, \dots, A_l\}} \Pr[A_1] \cdots \Pr[A_l] \\ &= \frac{1}{l!} \left( \sum_i \Pr[A_i] \right)^l \\ &= \frac{1}{l!} \mu^l. \quad \blacksquare \end{aligned}$$

### Erdős-Rado $\Delta$ -system

Let  $n$  and  $l$  be positive integers,  $l \geq 3$ . We say that  $l$  sets form a  $\Delta$ -system (of size  $l$ ), if they have pairwise the same intersection. Let  $\mathcal{F}$  be a family of sets, each of size  $n$ . Then

REPRESE

Lemm

The p  
be fo

Correl

In a p  
events  
events  
inequ  
the pr  
notati

Let  
Pr{x ∈  
x ∈ Ω.  
univer  
then

Lemm

The p  
origina

Borel-

The k  
probat  
followi  
probat

Lemm:  
space.

then w

3. PRC

Let k t  
and th.

**Lemma 2** ( $\Delta$ -System Lemma).  *$\mathcal{F}$  contains a  $\Delta$ -system of size  $l$  whenever*

$$|\mathcal{F}| \geq (l-1)^n N! \rightarrow n$$

The proof and a more general version of the theorem (for infinite set systems) can be found in [3].

**Correlation Inequality**

In a probability space, we know that the probability of a (finite) conjunction of events is exactly the product of the individual probabilities, provided that the events are *mutually independent*. In a similar spirit, the following *correlation inequality* allows us to estimate the probability of a conjunction by the product of the probabilities, provided that the events have *low correlation*! We need some notation to state the lemma precisely.

Let  $\Omega$  be a finite set of elements, and  $S$  be a random subset of  $\Omega$  given by  $\Pr[x \in S] = p_x$ , where we assume these events to be mutually independent over  $x \in \Omega$ . Let  $A_i$  denote the event  $X_i \subset S$ , where  $X_1, \dots, X_n$  are subsets of the universal set  $\Omega$ . Write  $i \sim j$  if  $i \neq j$  and  $X_i \cap X_j \neq \emptyset$ . Further let all  $\Pr[A_i] \leq 1/2$ , then

**Lemma 3** (Correlation Inequality).

$$\Pr\left[\bigwedge \bar{A}_i\right] \leq \left[\prod \Pr[\bar{A}_i]\right] e^{2 \sum_{i \sim j} \Pr[A_i \cap A_j]}.$$

The proof requires only elementary probability theory and appears in [1]. The original proof of the result was given by Svante Janson and appears in [6].

**Borel–Cantelli Lemma**

The last tool we make use of in this paper is one of great importance in probability theory. We merely state the well known Borel–Cantelli lemma in the following; the proof can be found in [5] and also in any standard book on probability theory.

**Lemma 4** (Borel–Cantelli). *Let  $\{A_i\}$  be a sequence of events in a probability space. If*

$$\sum_{j=1}^{\infty} \Pr(A_j) < \infty,$$

*then with probability 1, at most a finite number of the events  $A_j$  can occur.*

**3. PROOF OF THE MAIN THEOREM**

Let  $k$  be fixed. Our aim is to show that there exists an asymptotic basis of order  $k$  and that the number of representations is roughly  $\log n$  for  $n$  large. Consider a

sequence  $\mathcal{S}$  of natural numbers where the probability  $p(z)$  of  $z$  occurring in  $\mathcal{S}$  is given by

$$p(z) = C \frac{(\log z)^{1/k}}{z^{(k-1)/k}} \text{ for } z > z_0$$

$$= 0 \text{ otherwise,}$$

where  $z_0$  is the smallest constant such that  $p(z) \leq 1/2$ . We shall choose  $C$  so that  $C^k D_k > 3$ , where

$$D_k = [2^{1/k} - 1][3^{1/k} - 2^{1/k}] \cdots [(k-1)^{1/k} - (k-2)^{1/k}] \left(\frac{k^{k-1}}{k-1}\right)^{(k-1)/k},$$

a *not-so-nice* positive absolute constant!

Let  $r_k(n)$  be the random variable denoting the number of representations of  $n$  as a sum of  $k$  distinct numbers from  $\mathcal{S}$ . Then

**Lemma 5.**

$$\mu = E[r_k(n)] = \Theta(\log n)$$

*Proof.*

$$\mu = \sum_{\substack{x_1 + \cdots + x_k = n \\ 1 \leq x_1 < \cdots < x_k < n}} \Pr[x_1] \cdots \Pr[x_k]$$

$$= \sum^* C \left(\frac{\log x_1}{x_1^{k-1}}\right)^{1/k} \cdots C \left(\frac{\log x_k}{x_k^{k-1}}\right)^{1/k}$$

where  $\Sigma^*$  is short hand for  $\Sigma_{\substack{x_1 + \cdots + x_k = n \\ 1 \leq x_1 < \cdots < x_k < n}}$ . We estimate the above sum by breaking it into two parts ( $\mu_1 + \mu_2$ ), depending on whether  $x_1 > n/(\log n)$  or not. (It may be noted that "log  $n$ " here may be replaced by any positive function that approaches infinity sufficiently slowly as  $n \rightarrow \infty$ .) We prove the lemma by showing that  $\mu_1 = \Theta(\log n)$  and that  $\mu_2 = o(\log n)$ . For this purpose, let  $\delta_n = 1/(\log n)$ . Then

$$\mu_1 = C^k \sum_{n\delta_n < x_1 < n}^* \frac{[(\log x_1) \cdots (\log x_k)]^{1/k}}{(x_1 \cdots x_k)^{(k-1)/k}}$$

$$= C^k (1 + o(1)) (\log n) \sum_{n\delta_n < x_1 < n}^* \frac{1}{(x_1 \cdots x_k)^{(k-1)/k}}$$

$$= C^k (1 + o(1)) (\log n) \cdot (SUM) \text{ (say),}$$

where the numerator comes out as a common term, in view of the range of summation. It now suffices to show that  $SUM = \Theta(1)$ . Note that  $n/k < x_k < n$ , since  $x_i$ 's form an increasing sequence. As  $n/k < x_k$ , we bound  $SUM$  from above by

$$SUM < \frac{1}{(n/k)^{(k-1)/k}} \sum_{n\delta_n < x_1 < n}^* \frac{1}{(x_1 \cdots x_{k-1})^{(k-1)/k}}$$

REPRESENT/

We can  
( $x_1, \dots,$

We now

In the ot  
by

For a lo  
width  $\alpha$   
( $x_1 \cdots x_k$   
 $x_{k-1} < (k$

SUM :

and once

We can bound this tricky summation from above by summation over all  $(x_1, \dots, x_{k-1})$ , with  $1 \leq x_i \leq n$ . That is,

$$\begin{aligned} SUM &< \frac{1}{(n/k)^{(k-1)/k}} \sum_{\substack{1 \leq x_i \leq n \\ i=1, \dots, k-1}} \frac{1}{(x_1 \cdots x_{k-1})^{(k-1)/k}} \\ &= \left(\frac{k}{n}\right)^{(k-1)/k} \left(\sum_{1 \leq x_1 \leq n} \frac{1}{(x_1)^{(k-1)/k}}\right)^{k-1}. \end{aligned}$$

We now estimate this discrete sum by integrating over the same range:

$$\begin{aligned} SUM &< \left(\frac{k}{n}\right)^{(k-1)/k} \left(\int_1^n \frac{1}{x_1^{(k-1)/k}} dx_1 + O(1)\right)^{k-1} \\ &= \left(\frac{k}{n}\right)^{(k-1)/k} (kx_1^{1/k}|_1^n + O(1))^{k-1} \\ &= \frac{k^{(k-1)(k+1)/k}}{n^{(k-1)/k}} [n^{(k-1)/k} + o(n^{(k-1)/k})] \\ &= [1 + o(1)]k^{(k-1)(k+1)/k}. \end{aligned}$$

In the other direction, set  $\alpha = 1/k(k-1)$ . As  $x_k < n$ , we bound  $SUM$  from below by

$$SUM > \frac{1}{n^{(k-1)/k}} \sum_{n\delta_n < x_1 < n}^* \frac{1}{(x_1 \cdots x_{k-1})^{(k-1)/k}}.$$

For a lower bound on this summation, our idea is to restrict the  $x_i$ 's to strips of width  $\alpha n$ , i.e., we bound the summation from below by summation over  $(x_1 \cdots x_{k-1})$ , with restrictions  $n\delta_n < x_1 < \alpha n$ ,  $\alpha n < x_2 < 2\alpha n, \dots, (k-2)\alpha n < x_{k-1} < (k-1)\alpha n$ . (Our choice of  $\alpha$  ensures that  $x_{k-1} < x_k$ .) Thus

$$SUM > \frac{1}{n^{(k-1)/k}} \sum_{n\delta_n < x_1 < \alpha n} \frac{1}{x_1^{(k-1)/k}} \cdots \sum_{(k-2)\alpha n < x_{k-1} < (k-1)\alpha n} \frac{1}{x_{k-1}^{(k-1)/k}}$$

and once again, estimating by integrals,

$$\begin{aligned} &= \frac{1}{n^{(k-1)/k}} \left(\int_{n\delta_n}^{\alpha n} \frac{1}{x_1^{(k-1)/k}} + O(1)\right) \cdots \left(\int_{(k-2)\alpha n}^{(k-1)\alpha n} \frac{1}{x_{k-1}^{(k-1)/k}} + O(1)\right) \\ &= \frac{1}{n^{(k-1)/k}} (kx_1^{1/k}|_{n\delta_n}^{\alpha n} + O(1)) \cdots (kx_{k-1}^{1/k}|_{(k-2)\alpha n}^{(k-1)\alpha n} + O(1)) \\ &= [1 + o(1)]k^{k-1} \alpha^{(k-1)/k} [1 - (\alpha\delta_n)^{1/k}] [2^{1/k} - 1] \cdots [(k-1)^{1/k} - (k-2)^{1/k}] \\ &= D_k + o(1) \end{aligned}$$

where

range of  $x_k < n$ ,  
om above

breaking  
(It may  
tion that  
showing  
/(log n).

so that

ons of n

D TOTAL

in  $\mathcal{P}$  is

$$D_k = [2^{1/k} - 1][3^{1/k} - 2^{1/k}] \cdots [(k-1)^{1/k} - (k-2)^{1/k}] \left(\frac{k^{k-1}}{k-1}\right)^{(k-1)/k}$$

a positive absolute constant!

Consider now the second part:

$$\begin{aligned} \mu_2 &= C^k \sum_{1 < x_1 < n\delta_n}^* \frac{[(\log x_1) \cdots (\log x_k)]^{1/k}}{(x_1 \cdots x_k)^{(k-1)/k}} \\ &< C^k (\log n) \sum_{1 < x_1 < n\delta_n}^* \frac{1}{(x_1 \cdots x_k)^{(k-1)/k}} \\ &= C^k (\log n) \cdot (\text{sum}) \quad (\text{say}). \end{aligned}$$

It now suffices to show that  $\text{sum} = o(1)$ , since we are trying to prove  $\mu_2 = o(\log n)$ . The calculation is similar to what we did for  $SUM$  and proceeds as follows:

$$\begin{aligned} \text{sum} &< \frac{1}{(n/k)^{(k-1)/k}} \sum_{1 \leq x_1 \leq n\delta_n}^* \frac{1}{(x_1 \cdots x_{k-1})^{(k-1)/k}} \\ &< \frac{1}{(n/k)^{(k-1)/k}} \sum_{\substack{1 \leq x_1 \leq n\delta_n \\ 1 \leq x_i \leq n \\ i=2, \dots, k-1}} \frac{1}{(x_1 \cdots x_{k-1})^{(k-1)/k}} \\ &= \left(\frac{k}{n}\right)^{(k-1)/k} \left(\sum_{1 \leq x_1 \leq n\delta_n} \frac{1}{x_1^{(k-1)/k}}\right) \left(\sum_{1 \leq x_2 \leq n} \frac{1}{x_2^{(k-1)/k}}\right)^{k-2} \\ &= \left(\frac{k}{n}\right)^{(k-1)/k} \left(\int_1^{n\delta_n} \frac{1}{x_1^{(k-1)/k}} dx_1 + O(1)\right) \left(\int_1^n \frac{1}{x_2^{(k-1)/k}} dx_2 + O(1)\right)^{k-2} \\ &= \left(\frac{k}{n}\right)^{(k-1)/k} (kx_1^{1/k}|_1^{n\delta_n} + O(1))(kx_2^{1/k}|_1^n + O(1))^{k-2} \\ &= \frac{k^{(k-1)(k-2)/k}}{n^{(k-1)/k}} [\delta_n^{1/k} n^{(k-1)/k} + o(\delta_n^{1/k} n^{(k-1)/k})] \\ \text{sum} &= O(\delta_n^{1/k}). \end{aligned}$$

This shows that

$$\begin{aligned} \mu_2 &= C^k (\log n) O(\delta_n^{1/k}) \\ &= o(\log n). \end{aligned}$$

Thus we have established

$$[b_1 + o(1)] \log n < \mu_1 < [b_2 + o(1)] \log n \text{ and } \mu_2 = o(\log n)$$

and hence

$$[b_1 + o(1)] \log n < \mu < [b_2 + o(1)] \log n,$$

where  $b_1 = D_k C^k$  and  $b_2 = C^k k^{(k-1)(k+1)/k}$ .

3.1. Fi  
Let  $S_1$   
 $n$  in b

We sa  
lemm:  
disjoi

Lemm

Proof  
assoc  
lemm

Th  
repr  
of pa

Lemm

Proof

To  
 $b_1 =$   
yiel

### 3.1. First Part of the Main Theorem

Let  $S_1 = \{a_1, \dots, a_k\}$  and  $S_2 = \{b_1, \dots, b_k\}$  be two different representations of  $n$  in basis  $\mathcal{S}$ ; that is,  $S_1 \neq S_2$ ,  $S_1, S_2 \subset \mathcal{S}$  and

$$a_1 + \dots + a_k = b_1 + \dots + b_k = n.$$

We say  $S_i$  and  $S_j$  are *disjoint* if they share no element in common. The following lemmas guarantee (with high probability) that a maximal collection of pairwise disjoint representations has size at most  $O(\log n)$ .

**Lemma 6.**

$$\sum_{\substack{\{S_1, \dots, S_{6\mu}\} \\ \text{pairwise} \\ \text{disjoint}}} \Pr[S_1 \wedge \dots \wedge S_{6\mu}] \leq \mu^{6\mu} / (6\mu)!.$$

*Proof.* Pairwise disjointness of the sets implies mutual independence of the associated events. The proof is hence a simple application of the *disjointness lemma* (Lemma 1) with  $l = 6\mu$ . ■

This in turn bounds the probability of having more than  $6\mu$  pairwise disjoint representations of  $n$  as follows. Let  $r_k^*(n)$  denote the size of a maximal collection of pairwise disjoint representations. Then

**Lemma 7.** For  $C > (1/3D_k)^{1/k}$ ,

$$\Pr[r_k^*(n) > 6\mu] \leq n^{-2+o(1)}.$$

*Proof.* Using Lemma 6 and the bound  $m! > m^m e^{-m}$

$$\begin{aligned} \Pr[r_k^*(n) > 6\mu] &< \frac{1}{(6\mu)!} (\mu)^{6\mu} \\ &< \frac{1}{(6\mu/e)^{6\mu}} (\mu)^{6\mu} \\ &= \left(\frac{e}{6}\right)^{6\mu} \\ &< \left(\frac{e}{6}\right)^{[6b_1+o(1)]\log n} \\ &= n^{-6b_1+o(1)}. \end{aligned}$$

To complete the proof of the lemma, we have to show that  $b_1 \geq 1/3$ . As  $b_1 = D_k C^k$ , we pick  $C$  large enough (so that  $D_k C^k > 1/3$ ) to make  $b_1 > 1/3$ , yielding

$$\Pr[r_k^*(n) > 6\mu] \leq n^{-2+o(1)}. \quad \blacksquare$$

**Lemma 7a.** A.a.  $\exists n_0$  s.t.  $r_k^*(n) \leq 6\mu$ , for  $n > n_0$ .

And onc  
with  $1 \leq$

*Proof.* Use Borel–Cantelli lemma! First, a few words about the *probability space* we are dealing with. Let  $\Omega$  denote universal set of all subsequences of the sequence of natural numbers. Our *random set*  $\mathcal{S}$  is the subsequence formed by choosing number  $z$  with probability  $p(z)$ . Thus the triple  $(\Omega, \mathcal{S}, p(\cdot))$  is our *probability space*. (For a rigorous proof that this is a *valid* probability space, Ref. [5], pp. 141–142.)

Now let  $A_n$  denote the event  $r_k^*(n) > 6\mu$ . By Lemma 7, we have

$$\sum_{n=1}^{\infty} \Pr(A_n) = \sum_{n=1}^{\infty} n^{-2+o(1)} < \infty.$$

Substitut

By the Borel–Cantelli lemma, this implies that with probability 1, at most a *finite* number of  $A_n$  occur. Hence the lemma. ■

To prove the first part of the main theorem, we must extend Lemma 7a and show that  $r_k(n)$  is  $O(\log n)$ . For this purpose we need a bound [of  $O(1)$ ] on the number of representations of  $n$  as a sum of  $k - 1$  (distinct) numbers. (The puzzled reader is advised to skip the next few lemmas and read the proof of Theorem 2 for further motivation.) But we need to generalize some of our notation before we can proceed. Let  $r_l(n)$  denote the number of representations of  $n$  as a sum of  $l$  distinct numbers from  $\mathcal{S}$ . And let  $r_l^*(n)$  denote the size of a maximal collection of *pairwise disjoint* such representations. Further, let  $\mu_l$  denote the expectation of  $r_l(n)$  (note that  $\mu_k$  is simply  $\mu$ )

Let  $S^l$

Lemma 5

**Lemma 8.** For  $2 \leq l \leq (k - 1)$ ,

$$\mu_l \leq n^{-1+l/k+o(1)}.$$

*Proof.* (independ  
from the  
(ii) Fron

*Proof.* The line of calculation is similar to that for  $\mu$ , only simpler. Let  $l$  be fixed. Let  $\Sigma^*$  denote  $\sum_{\substack{x_1+\dots+x_l=n \\ 1 \leq x_1 < \dots < x_l < n}}$ . Then by definition,

$$\begin{aligned} \mu_l &= \sum_{\substack{x_1+\dots+x_l=n \\ 1 \leq x_1 < \dots < x_l < n}} \Pr[x_1] \cdots \Pr[x_l] \\ &= \sum^* C \left( \frac{\log x_1}{x_1^{k-1}} \right)^{1/k} \cdots C \left( \frac{\log x_l}{x_l^{k-1}} \right)^{1/k} \\ &= n^{o(1)} \sum^* \frac{1}{(x_1 \cdots x_l)^{(k-1)/k}} \\ &= n^{o(1)} SUM_l. \end{aligned}$$

As  $l \leq k$

As  $n/l < x_l$ ,

$$SUM_l = n^{-(k-1)/k+o(1)} \sum^* \frac{1}{(x_1 \cdots x_{l-1})^{(k-1)/k}}.$$

Once

And once again we estimate crudely by the summation over all  $(x_1, \dots, x_{l-1})$ , with  $1 \leq x_i \leq n, i = 1, 2, \dots, l-1$ :

$$\begin{aligned} \text{SUM}_l &\leq n^{-(k-1)/k+o(1)} \sum_{\substack{1 \leq x_i \leq n \\ i=1, \dots, l-1}} \frac{1}{(x_1 \cdots x_{l-1})^{(k-1)/k}} \\ &= n^{-(k-1)/k+o(1)} \left( \sum_{1 \leq x_1 \leq n} \frac{1}{x_1^{(k-1)/k}} \right)^{l-1} \\ &= n^{-(k-1)/k+o(1)} (n^{1/k+o(1)})^{l-1} \\ &= n^{-1+l/k+o(1)}. \end{aligned}$$

Substituting this in the estimate for  $\mu_l$  above, we get

$$\begin{aligned} \mu_l &\leq n^{o(1)} (n^{-1+l/k+o(1)}) \\ &= n^{-1+l/k+o(1)}. \end{aligned}$$

Let  $S^l$  denote a representation of  $n$  as a sum of  $l$  distinct numbers. Then:

**Lemma 9.** For  $2 \leq l \leq (k-1)$ ,

- (i)  $\sum_{\substack{\{S_1^l, \dots, S_{2k}^l\} \\ \text{pairwise} \\ \text{disjoint}}} \Pr[S_1^l \wedge \dots \wedge S_{2k}^l] < \frac{\mu_l^{2k}}{(2k)!}$
- (ii)  $\Pr[r_l^*(n) > 2k] < n^{-2+o(1)}$ .

*Proof.* (i) Note that when  $S_i^l$  and  $S_j^l$  are disjoint,  $\Pr[S_i^l]$  and  $\Pr[S_j^l]$  are mutually independent probabilities. So the first part of the lemma follows immediately from the *disjointness lemma*.

(ii) From part (i) it follows that

$$\begin{aligned} \Pr[r_l^*(n) > 2k] &< \frac{\mu_l^{2k}}{(2k)!} \\ &< \frac{1}{(2k)!} (n^{-1+l/k+o(1)})^{2k} \\ &= n^{-2k+2l+o(1)} \end{aligned}$$

As  $l \leq k-1$ , we can conclude

$$\Pr[r_l^*(n) > 2k] < n^{-2+o(1)}.$$

Once again, by the Borel-Cantelli lemma, the above assertion implies that

$$\text{a.a. for } 2 \leq l \leq (k-1) \quad \exists n_l \text{ s.t. } n > n_l \Rightarrow r_l^*(n) < 2k.$$

But for any finite  $n_l$ , there are at most a finite [certainly  $< \binom{n_l}{l}$ ] number of representations as a sum of  $l$  numbers. Therefore,

$$\text{a.a. for } 2 \leq l \leq (k-1) \exists c_l \text{ s.t. } \forall n, r_l^*(n) < c_l.$$

At this point we remind the reader that we set out to prove that  $r_{k-1}(n)$  is at most a constant. And we make critical use of the  $\Delta$ -system lemma (Lemma 2) to prove this.

**Lemma 10.**

$$\text{a.a. } \exists c \text{ s.t. } r_{k-1}(n) < c, \forall n.$$

*Proof.* We know

$$\text{a.a. for } 2 \leq l \leq (k-1) \exists c_l \text{ s.t. } \forall n, r_l^*(n) < c_l. \quad (*)$$

Set  $c_{\max} = \max_l \{c_l\}$ . We claim (whenever  $c_l$ 's exist),

$$\forall n r_{k-1}(n) \leq (c_{\max})^{k-1} (k-1)!$$

Suppose not, i.e., the claim is false for some  $n = n'$ . Then by the  $\Delta$ -system lemma, there exists a  $\Delta$ -system  $\{S_1^{k-1}, \dots, S_{\Delta}^{k-1}\}$  of size  $\Delta = c_{\max} + 1$ . Let the common intersection of the system be  $R$ . So  $R = \{x_1, \dots, x_r\}$ , where  $0 \leq r \leq k-2$ . If  $\sum_i x_i = m$ , then removing the common intersection  $R$  from each set will yield  $r_{k-1-r}^*(n' - m) \geq c_{\max} + 1$ . This is impossible in view of (\*) and the definition of  $c_{\max}$ . This proves the lemma and, in fact, also shows that  $c \leq c_{\max}^{k-1} (k-1)!$  ■

We are now ready to prove one side of the main theorem.

**Theorem 2.** *A.a.*  $\exists c \exists n_0$  s.t.  $r_k(n) < [6b_2ck + o(1)] \log n, n > n_0$ .

*Proof.* Let  $\hat{S}$  be any maximal collection of pairwise disjoint representations of  $n$  as a sum of  $k$  distinct numbers. By our notation,  $|\hat{S}| = r_k^*(n)$ . By Lemma 7a, we have

$$\text{A.a. } \exists n_0 \text{ s.t. } r_k^*(n) < 6\mu, n > n_0.$$

This implies there are at most  $(k) \times (6\mu)$  numbers in our collection  $\hat{S}$ . [There are (almost always) at most  $6\mu$  sets in the collection, and each set has exactly  $k$  numbers.] As  $\hat{S}$  is maximal, any representation of  $n$  must use at least one number from the collection. However, the number of representations of  $n$  which use  $x$  is precisely  $r_{k-1}(n-x)$ . By Lemma 10 we know  $r_{k-1}(n-x) < c$ . Thus the total number of representations of  $n$  is at most  $(6k\mu) \times c$ . Hence the theorem follows by substituting the bound for  $\mu$ .

3.2. Sect

In the fir like to pr that  $r_k^*(n)$  lemmas.

Let  $S_i$  from  $\mathcal{S}$ . valid repi in the fol

Lemma 1

*Proof.* 1 numbers.

Consider

$$S_i = (z_i)$$

Let  $\sum_i z_i$

$$\sum_{|S_i \cap S_j| = l} 1$$

Fortunate for  $1 \leq l \leq$

Since  $m_0$  : above sur  $n/2$ , (iii)

3.2. Second part of the main theorem

In the first part, we have proved that  $r_k(n) < C_2 \log n$ , for large  $n$ . We now would like to prove that  $r_k(n) > C_1 \log n$ , for large  $n$ . We prove the stronger inequality, that  $r_k^*(n) > C_1 \log n$ , for large  $n$ . Before we can prove this, we need two more lemmas.

Let  $S_i$  and  $S_j$  be two valid representations of  $n$  as a sum of  $k$  distinct numbers from  $\mathcal{S}$ . With abuse of notation, we let  $\Pr[S_i]$  denote the probability that  $S_i$  is a valid representation. We would like to show that these events have *low correlation* in the following precise sense.

**Lemma 11.**

$$\sum_{i \sim j} \Pr[S_i \wedge S_j] = o(1).$$

*Proof.* Note that  $i \sim j$  implies  $S_i$  and  $S_j$  share at least 1 number and at most  $k - 2$  numbers.

$$\sum_{i \sim j} \Pr[S_i \wedge S_j] = \sum_{l=1}^{k-2} \sum_{|S_i \cap S_j|=l} \Pr[S_i \wedge S_j].$$

Consider  $S_i, S_j$  such that  $|S_i \cap S_j| = l$ . Say,

$$S_i = (z_1, \dots, z_l, x_1, x_2, \dots, x_{k-l}) \text{ and } S_j = (z_1, \dots, z_l, y_1, y_2, \dots, y_{k-l}).$$

Let  $\sum_i z_i$  be  $m$ . Then  $\sum_i x_i = \sum_i y_i = n - m$ . So

$$\begin{aligned} \sum_{|S_i \cap S_j|=l} \Pr[S_i \wedge S_j] &= \sum_m \sum_{\substack{z_1 + \dots + z_l = m \\ x_1 + \dots + x_{k-l} = n - m \\ y_1 + \dots + y_{k-l} = n - m}} (\Pr[z_1] \cdots \Pr[z_l]) (\Pr[x_1] \cdots \Pr[x_{k-l}]) \\ &\quad \times (\Pr[y_1] \cdots \Pr[y_{k-l}]) \\ &= \sum_m \left( \sum_{z_1 + \dots + z_l = m} \Pr[z_1] \cdots \Pr[z_l] \right) \\ &\quad \times \left( \sum_{x_1 + \dots + x_{k-l} = n - m} \Pr[x_1] \cdots \Pr[x_{k-l}] \right)^2 \\ &= \sum_m \mu_l(m) [\mu_{k-l}(n - m)]^2. \end{aligned}$$

Fortunately, we already made (in Lemma 8) the estimates  $\mu_l(n) < n^{-1+l/k+o(1)}$ , for  $1 \leq l \leq k - 1$ . Fix  $\epsilon < l/2k$ . Then we can pick  $m_0$  such that

$$\mu_l(m) < m^{-1+l/k+o(1)}, \text{ for } m > m_0.$$

Since  $m_0$  is a constant,  $\mu_l(m) < M$  (some constant), for  $m \leq m_0$ . We estimate the above summation in four parts ( $\Delta_1 + \Delta_2 + \Delta_3 + \Delta_4$ ): (i)  $m \leq m_0$ , (ii)  $m_0 < m \leq n/2$ , (iii)  $n/2 < m \leq n - m_0$ , and (iv)  $n - m_0 < m$ .

Two of the following four estimates are easy; we do the easy ones first:

$$\begin{aligned} \Delta_1 &= \sum_{m \leq m_0} \mu_l(m) [\mu_{k-l}(n-m)]^2 \\ &< (n^{-1+(k-l)/k+o(1)})^2 \sum_{m \leq m_0} M \\ &= n^{-2+2(k-l)/k+o(1)} = o(1). \end{aligned}$$

Thus,

The other extreme ( $m > n - m_0$ ) can be estimated similarly:

and he

$$\begin{aligned} \Delta_4 &= \sum_{m > n - m_0} \mu_l(m) [\mu_{k-l}(n-m)]^2 \\ &< (n^{-1+l/k+o(1)}) \sum_{m > n - m_0} M^2 \\ &= n^{-1+l/k+o(1)} = o(1). \end{aligned}$$

Let  
We wo  
hurt  $r_k$   
 $n$ ;  $x_i$  di

Second and third parts of the sum:

Lemma

$$\begin{aligned} \Delta_2 &= \sum_{m_0 < m \leq n/2} \mu_l(m) [\mu_{k-l}(n-m)]^2 \\ &< (n^{-1+(k-l)/k+o(1)})^2 \sum_{m_0 < m \leq n/2} m^{-1+l/k+\epsilon} \\ &= n^{-2l/k+o(1)} \sum_{m_0 < m \leq n/2} m^{-1+l/k+\epsilon} \end{aligned}$$

Proof.

It now suffices to estimate the sum by an integral over the full range  $0 \leq m \leq n$ :

$$\begin{aligned} \Delta_2 &< n^{-2l/k+o(1)} \left( \int_{m=0}^n m^{-1+l/k+\epsilon} + O(1) \right) \\ &= n^{-2l/k+o(1)} (m^{l/k+\epsilon} \Big|_0^n + O(1)) \\ &= n^{-l/k+\epsilon+o(1)} = o(1). \end{aligned}$$

where  $\int$   
strips  $o$

i.  $x$   
ii.  $n$

Similarly,

$$\begin{aligned} \Delta_3 &= \sum_{n/2 < m \leq n - m_0} \mu_l(m) [\mu_{k-l}(n-m)]^2 \\ &< (n^{-1+l/k+o(1)}) \sum_{n/2 < m \leq n - m_0} [(n-m)^{-1+(k-l)/k+\epsilon}]^2 \\ &= (n^{-1+l/k+o(1)}) \sum_{n/2 < m \leq n - m_0} (n-m)^{-2l/k+2\epsilon} \end{aligned}$$

where  $\int$   
We wo

Estimating by integration over the full range,

st:

$$\begin{aligned} \Delta_3 &< n^{-1+l/k+o(1)} \left( \int_{m-0}^n (n-m)^{-2l/k+2\epsilon} + O(1) \right) \\ &< n^{-1+l/k+o(1)} (-(n-m)^{1-2l/k+2\epsilon} \Big|_0^n + O(1)) \\ &= n^{-l/k+2\epsilon+o(1)} = o(1). \end{aligned}$$

Thus,

$$\sum_{|S_i \cap S_j|=l} \Pr[S_i \wedge S_j] = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 = o(1)$$

and hence the claim,

$$\sum_{i \sim j} \Pr[S_i \wedge S_j] = \sum_{i=1}^{k-2} o(1) = o(1). \quad \blacksquare$$

Let  $s = \{S_1, \dots, S_t\}$  be a collection of pairwise disjoint representations of  $n$ . We would like to show that, when  $|s| < C_1 \log n$ , deleting sets  $S_i$  from  $\mathcal{S}$  does not hurt  $r_k(n)$  much. For this purpose, set  $r_k(n; s) = |\{(x_1, \dots, x_k) : x_1 + \dots + x_k = n; x_i \text{ distinct and } x_i \in \mathcal{S} \setminus s\}|$  and define  $\mu' = E[r_k(n; s)]$ .

**Lemma 12.**

$$|s| < C_1 \log n \Rightarrow \mu' = \Theta(\log n).$$

*Proof.* By definition,

$$\begin{aligned} \mu' &= \sum_{\substack{x_1 + \dots + x_k = n \\ 1 < x_1 < \dots < x_k < n \\ x_1, \dots, x_k \notin \cup_i S_i}} \Pr[x_1] \cdots \Pr[x_k] \\ &> \sum_{x_1, \dots, x_k \notin \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k] \end{aligned}$$

$\leq m \leq n$ :

where  $\Sigma^{**}$  denotes the special summation (of Lemma 5) with the  $x_i$ 's restricted to *strips* of width  $\alpha n$ ; i.e., the summation is over  $x_1, \dots, x_k$  such that

- i.  $x_1 + \dots + x_k = n$ .
- ii.  $n\delta_n < x_1 < \alpha n, \alpha n < x_2 < 2\alpha n, \dots, (k-2)\alpha n < x_{k-1} < (k-1)\alpha n$ .

$$\begin{aligned} \mu' &\geq \sum^{**} \Pr[x_1] \cdots \Pr[x_k] - \sum_j \sum_{x_j \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k], \\ &> [b_1 + o(1)] \log n - \sum_j \sum_{x_j \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k], \end{aligned}$$

where the above estimate follows from the estimate (from below) for  $\mu$  in Lemma 5. We would now like to show that for  $j = 1, \dots, k$

$$\sum_{x_j \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k] = o(\log n).$$

Without loss of generality, we show that for  $x_k$  (i.e.,  $j = k$ ). As  $S_1, \dots, S_{C_1 \log n}$  are pairwise disjoint

We also

$$\sum_{x_k \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k] = (C_1 \log n) \sum_{x_k \in S_1}^{**} \Pr[x_1] \cdots \Pr[x_k] < (C_1 \log n) \mu_{k-1}(n - x_k)$$

We estimate that a se

But our choice of the range for  $\Sigma^{**}$  was such that  $x_k < n - (k - 1)\alpha n$ . Thus

$$\sum_{x_k \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k] = (C_1 \log n) (n^{-1+(k-1)/k+o(1)}) = o(\log n).$$

By the c

[In the general case, the range for  $x_j$  ensures that  $(n - x_j) > (1 - j\alpha)n$ , allowing us the estimate  $\mu_{k-1}(n - x_j) < n^{-1+(k-1)/k+o(1)}$ .] Thus

$$\sum_j \sum_{x_j \in \cup_i S_i}^{**} \Pr[x_1] \cdots \Pr[x_k] = o(\log n)$$

and hence

$$\mu' > [b_1 + o(1)] \log n - o(\log n).$$

With preceding  $B_j$ . Thus

On the other hand, we have, trivially,  $\mu' < \mu < [b_2 + o(1)] \log n$ , completing the proof of the lemma. ■

We are now set to complete the second part of the main theorem.

By Lemm

**Theorem 3.** *A.a.*  $\exists n_1$  s.t.  $r_k(n) > C_1 \log n$ ,  $n > n_1$ .

*Proof.* We would first want to show the probability that  $r_k^*(n) \leq C_1 \log n$  is small, for all  $n$ . Then we could complete the proof by using the Borel–Cantelli lemma. Recall that  $s$  denotes a collection of pairwise disjoint representations of  $n$ . We want to estimate

Together

$$\Pr[\exists s: |s| \leq C_1 \log n, \text{ } s \text{ maximal}]$$

Plugging

$$= \sum_{i=0}^{C_1 \log n} \Pr[\exists s: |s| = i, \text{ } s \text{ maximal}]$$

$\Pr[\exists s:$

$$= \sum_{i=0}^{C_1 \log n} \sum_{\substack{S_1, \dots, S_i \\ \text{pairwise disjoint}}} (\Pr[S_1 \wedge \dots \wedge S_i] \times \Pr[\{S_1, \dots, S_i\} \text{ is maximal}])$$

$$= \sum_{i=0}^{C_1 \log n} \left( \sum_{\substack{S_1, \dots, S_i \\ \text{pairwise disjoint}}} \Pr[S_1 \wedge \dots \wedge S_i] \right) \times \Pr[\{S_1, \dots, S_i\} \text{ is maximal}] \quad (**)$$

We already know how to estimate the first factor. By the *disjointness lemma*,

$$\sum_{\substack{S_1, \dots, S_i \\ \text{pairwise} \\ \text{disjoint}}} \Pr[S_1 \wedge \dots \wedge S_i] \leq \mu^i / i!$$

We estimate the second factor by the *correlation inequality*! Let  $B_j$  be the event that a set  $S_j' \subset \mathcal{S}$  of  $k$  distinct numbers is a representation of  $n$ . Then,

$$\Pr[\{S_1, \dots, S_i\} \text{ is maximal}] = \Pr[\cap_j \bar{B}_j]$$

By the correlation inequality (Lemma 3),

$$\begin{aligned} \Pr[\cap_j \bar{B}_j] &\leq \left[ \prod_j \Pr[\bar{B}_j] \right] e^{2\sum_{j \neq j'} \Pr[B_j \wedge B_{j'}]} \\ &= \left[ \prod_j (1 - \Pr[B_j]) \right] e^{2\sum_{j \neq j'} \Pr[B_j \wedge B_{j'}]} \\ &\leq e^{-\sum_j \Pr[B_j]} e^{2\sum_{j \neq j'} \Pr[B_j \wedge B_{j'}]} \end{aligned}$$

With foresight, we have computed the two exponents in Lemmas 11 and 12 preceding this theorem!  $r_k(n; s)$  is precisely the number of the probabilistic events  $B_j$ . Thus by Lemma 12,

$$\sum_j \Pr[B_j] = E[r_k(n; s)] = \mu' > [b_1 - o(1)] \log n.$$

By Lemma 11, the correlation

$$2 \sum_{j \neq j'} \Pr[B_j \wedge B_{j'}] \leq 2 \sum_{i \neq j} \Pr[S_i \wedge S_j] = o(1).$$

Together these facts imply

$$\Pr[\{S_1, \dots, S_i\} \text{ is maximal}] < e^{-[b_1 - o(1)] \log n}.$$

Plugging this in (\*\*\*) yields

$$\begin{aligned} \Pr[\exists s: |s| \leq C_1 \log n, s \text{ maximal}] &< \sum_{i=0}^{C_1 \log n} \left( \frac{\mu^i}{i!} \right) e^{-[b_1 - o(1)] \log n} \\ &= e^{-[b_1 - o(1)] \log n} \sum_{i=0}^{C_1 \log n} \frac{\mu^i}{i!} \\ &< e^{-[b_1 - o(1)] \log n} \left( \frac{e}{C_1} \right)^{C_1 \mu}, \text{ provided that} \\ &\quad 0 < C_1 < 1, \\ &< e^{-[b_1 - o(1)] \log n} \left( \frac{e}{C_1} \right)^{C_1 [b_2 + o(1)] \log n} \end{aligned}$$

(\*\*\*)

We choose  $C_1$  small enough so that  $(e/C_1)^{C_1[b_2+o(1)]\log n} < e^{\log n}$ . This is possible, since  $(e/C_1)^{C_1} \rightarrow 1$  as  $C_1 \rightarrow 0^+$ .

$$\Pr[\exists s: |s| \leq C_1 \log n, s \text{ maximal}] < e^{-[b_1-o(1)] \log n} e^{\log n} = n^{-b_1+1+o(1)}$$

Recall that  $b_1 = D_k C^k$ . So by choosing  $C > (3/D_k)^{1/k}$ , we can make  $b_1 > 3$ . [Note that this choice makes the condition  $C > (1/3D_k)^{1/k}$  in Lemma 7 superfluous.] Thus,

$$\Pr[\exists s: |s| \leq C_1 \log n, s \text{ maximal}] < n^{-2+o(1)}.$$

But this implies

$$\sum_{n=1}^{\infty} \Pr[\exists s: |s| \leq C_1 \log n, s \text{ maximal}] = \sum_{n=1}^{\infty} n^{-2+o(1)} < \infty,$$

allowing us to conclude, by the Borel-Cantelli lemma,

$$\text{a.a. } \exists n_1 \text{ s.t. } r_k(n) > C_1 \log n, \text{ for } n > n_1.$$

Theorems 2 and 3 together imply the main theorem (Theorem 1).

*Alternative Proof of Theorem 3.* We have realized Theorem 3 can be proved with much less effort with a very recent result of Svante Janson. (In particular, Lemma 12 becomes unnecessary.) Janson's result [7] is a generalization of his original correlation inequality and can be described as follows. Let  $A_1, \dots, A_n$  be the events as defined in the correlation inequality lemma. Further let  $N$  denote the (random) number of events in the family  $\{A_i\}$  which occur. Define

$$\mu = E[N] = \sum_i \Pr[A_i] \text{ and } \delta = \frac{1}{\mu} \sum_{i \sim j} \Pr[A_i \wedge A_j]$$

**Theorem (S. Janson).** If  $0 \leq \epsilon \leq 1$ , then

$$\Pr[N \leq (1 - \epsilon)\mu] \leq e^{-[1/2(1+\delta)]\epsilon^2\mu}.$$

(Note that correlation inequality is the special case when  $\epsilon = 1$ .)

To prove Theorem 3 we first identify  $A_i$  to be the event that a valid representation  $S_i$  of  $n$  is present in our random set  $\mathcal{S}$ . Then  $N$  is precisely  $r_k(n)$ . By Janson's theorem it would follow, for  $0 \leq C_1 \leq 1$

$$\Pr[r_k(n) \leq C_1\mu] \leq e^{-[1/2(1+\delta)](1-C_1)^2\mu}$$

where  $\delta = (1/\mu) \sum_{i \sim j} \Pr[A_i \wedge A_j] = o(1)$ , by Lemma 11. We also know,  $\mu > [b_1 + o(1)]\log n$ . Thus

$$\Pr[r_k(n) \leq C_1\mu] \leq e^{-\{(1/2[1+o(1)])(1-C_1)^2[b_1+o(1)]\}\log n}$$

REPRESENT

Once agreed

Theorem

ACKNOWLEDGMENT

We wish

REFERENCES

[1] R. B. *Comb*  
 [2] P. Erdős *des N*  
 [3] P. Erdős *Soc.*  
 [4] P. Erdős *Acta*  
 [5] H. Ha  
 [6] S. Janson *nonexi Graph*  
 [7] S. Janson 221-2.

Received C

Once again, we observe that  $C_1$  can be chosen arbitrarily small, and  $b_1$  can be made greater than 4, so that

$$\Pr[r_k(n) \leq C_1 \mu] \leq n^{-2+o(1)}.$$

Theorem 3 now follows by simply applying the Borel–Cantelli lemma.

#### ACKNOWLEDGMENT

We wish to thank Joel Spencer for many useful discussions.

#### REFERENCES

- [1] R. B. Boppana and J. H. Spencer, A useful elementary correlation inequality, *J. Combinat. Theory Ser. A*, **50**, 305–307 (1989).
- [2] P. Erdős, Problems and results in additive number theory, in *Colloque sur la Théorie des Nombres (CBRM)*, Bruxelles, 1956, pp. 127–137.
- [3] P. Erdős and R. Rado, Intersection theorems for systems of sets, *J. London Math. Soc.*, **35**, 85–90 (1960).
- [4] P. Erdős and A. Rényi, Additive properties of random sequences of positive integers, *Acta Arith.*, **6**, 83–110 (1960).
- [5] H. Halberstam and K. F. Roth, *Sequences*, Springer-Verlag, New York, 1983.
- [6] S. Janson, T. Łuczak, and A. Ruciński, An exponential bound for the probability of nonexistence of a specified subgraph in a random graph, *Proceedings of Random Graphs '87 (Poznań)*, (M. Karoński, ed.) to appear.
- [7] S. Janson, Poisson approximation for large deviations, *Random Struct. Alg.*, **1**, 221–229 (1990).

Received October 30, 1989