

NON-PLANAR EXTENSIONS OF SUBDIVISIONS OF PLANAR GRAPHS

Sergey Norin

Department of Mathematics and Statistics
McGill University
Montreal, Quebec H3A 2K6, Canada

and

Robin Thomas¹

School of Mathematics
Georgia Institute of Technology
Atlanta, Georgia 30332-0160, USA

5 November 1998, Revised 19 December 2013.

¹ Partially supported by NSF under Grants No. DMS-9623031, DMS-0200595 and DMS-1202640, and by NSA under Grant No. MDA904-98-1-0517.

ABSTRACT

A graph G is *almost 4-connected* if it is simple, 3-connected, has at least five vertices, and $V(G)$ cannot be partitioned into three sets A, B, C in such a way that $|C| = 3$, $|A| \geq 2$, $|B| \geq 2$, and no edge of G has one end in A and the other end in B . A graph K is a *subdivision* of a graph G if K is obtained from G by replacing its edges by internally disjoint nonzero length paths with the same ends, called *segments*. Let G, H be almost 4-connected graphs such that G is planar and has at least seven vertices, H is non-planar, and H has a subgraph isomorphic to a subdivision of G . If K is a subgraph of H , then a *K -path* in H is a path with at least one edge, both ends in K , and otherwise disjoint from K . We prove that there exists a subgraph K of H isomorphic to a subdivision of G such that one of the following conditions holds.

- (i) There exists a K -path in H such that no face boundary of K contains both ends of the path.
- (ii) There exist two disjoint K -paths with ends s_1, t_1 and s_2, t_2 , respectively, such that the vertices s_1, s_2, t_1, t_2 belong to some face boundary of K in the order listed. Moreover, for $i = 1, 2$ the vertices s_i and t_i do not belong to the same segment of K , and if two segments of K include all of s_1, t_1, s_2, t_2 , then those segments are vertex-disjoint.

This is a lemma to be used in other papers. In fact, we prove a more general theorem, where we relax the connectivity assumptions and do not assume that G is planar. Instead of face boundaries we work with a collection of cycles that cover every edge twice and have pairwise connected intersection. Finally, we prove a version of this result that applies when $G \setminus X$ is planar for some set $X \subseteq V(G)$ of size at most k , but $H \setminus Y$ is non-planar for every set $Y \subseteq V(H)$ of size at most k .

1. INTRODUCTION

In this paper graphs are finite and simple (i.e., they have no loops or multiple edges). *Paths* and *cycles* have no “repeated” vertices or edges. A graph is a *subdivision* of another if the first can be obtained from the second by replacing each edge by a non-zero length path with the same ends, where the paths are disjoint, except possibly for shared ends. The replacement paths are called *segments*, and their ends are called *branch-vertices*. For later convenience a one-vertex component of a graph is also regarded as a segment, and its unique vertex as a branch-vertex. Let G, S, H be graphs such that S is a subgraph of H and is isomorphic to a subdivision of G . In that case we say that S is a G -*subdivision* in H . If G has no vertices of degree two (which will be the case in our applications), then the segments and branch-vertices of S are uniquely determined by S . An S -*path* is a path of length at least one with both ends in S and otherwise disjoint from S . A graph G is *almost 4-connected* if it is simple, 3-connected, has at least five vertices, and $V(G)$ cannot be partitioned into three sets A, B, C in such a way that $|C| = 3$, $|A| \geq 2$, $|B| \geq 2$, and no edge of G has one end in A and the other end in B .

Let a non-planar graph H have a subgraph S isomorphic to a subdivision of a planar graph G . For various problems in structural graph theory it is useful to know the minimal subgraphs of H that have a subgraph isomorphic to a subdivision of G and are non-planar. We show that under some mild connectivity assumptions these “minimal non-planar extensions” of G are quite nice:

(1.1) *Let G be an almost 4-connected planar graph on at least seven vertices, let H be an almost 4-connected non-planar graph, and let there exist a G -subdivision in H . Then there exists a G -subdivision S in H such that one of the following conditions holds:*

(i) *there exists an S -path in H joining two vertices of S not incident with the same face,*

or

(ii) *there exist two disjoint S -paths with ends s_1, t_1 and s_2, t_2 , respectively, such that the vertices s_1, s_2, t_1, t_2 belong to some face boundary of S in the order listed. Moreover, for $i = 1, 2$ the vertices s_i and t_i do not belong to the same segment of S , and if two segments of S include all of s_1, t_1, s_2, t_2 , then those segments are vertex-disjoint.*

The connectivity assumptions guarantee that the face boundaries in a planar embedding of S are uniquely determined, and hence it makes sense to speak about incidence with faces. The result is related to, but independent of [9].

In Section 6 we deduce the following corollary, stated there as (6.6). A graph is a *minor* of another if the first can be obtained from a subgraph of the second by contracting edges. If G is a graph and $u, v \in V(G)$ are not adjacent, then by $G + uv$ we denote the graph obtained from G by adding an edge with ends u and v .

(1.2) *Let G be an almost 4-connected triangle-free planar graph, and let H be an almost 4-connected non-planar graph such that H has a subgraph isomorphic to a subdivision of G . Then there exists a graph G' such that G' is isomorphic to a minor of H , and either*

- (i) $G' = G + uv$ for some vertices $u, v \in V(G)$ such that no facial cycle of G contains both u and v , or*
- (ii) $G' = G + u_1v_1 + u_2v_2$ for some distinct vertices $u_1, u_2, v_1, v_2 \in V(G)$ such that u_1, u_2, v_1, v_2 appear on some facial cycle of G in the order listed.*

While the statement of (1.2) is nicer, it has the drawback that we assume that H has a subgraph isomorphic to a *subdivision* of G , and deduce that it has only a *minor* isomorphic to G' . That raises the question whether there is a similar theory that applies when H has a minor isomorphic to G . Such a theory indeed exists and is developed in [2], using (4.6). Informally, there is an analogue of (1.1), where either of the two outcomes may be preceded by up to two vertex splits (inverse operations to edge contraction).

In the applications of (1.1) the graph G is known explicitly, but H is not, and we are trying to deduce some information about H . Since it is possible to generate all graphs that can be obtained from subdivision of G by means of (i) or (ii) above, we thus obtain a list of specific non-planar graphs such that H has a subgraph isomorphic to a subdivision of one of the graphs in the list. The graphs G of interest in applications tend to possess a lot of symmetry, and so the generation process is usually less daunting than it may seem.

A sample application of our result is presented in Section 7, but let us informally describe the applications from [1] and [12]. Theorem (6.2), a close relative of (1.1), is used in [1] to show that for every positive integer k , there is an integer N such that

every 4-connected non-planar graph with at least N vertices has a minor isomorphic to the complete bipartite graph $K_{4,k}$, or the graph obtained from a cycle of length $2k + 1$ by adding an edge joining every pair of vertices at distance exactly k , or the graph obtained from a cycle of length k by adding two vertices adjacent to each other and to every vertex on the cycle. Using this it can be shown that, except for one well-defined infinite family, there are only finitely many graphs of crossing number at least two that are minimal in a specified sense.

In [12] it is shown that an almost 4-connected non-planar graph of girth at least five has a subgraph isomorphic to a subdivision of P_{10}^- , the Petersen graph with one edge deleted. (It follows from this that Tutte's 4-flow conjecture [15] holds for graphs with no subdivision isomorphic to P_{10}^- .) The way this is done is that first it is shown that if G is a graph of girth at least five and minimum degree at least three, then it has a subgraph isomorphic to a subdivision of the Dodecahedron or P_{10}^- . Corollary (1.2) is then used to show that if G is an almost 4-connected non-planar graph with a subgraph isomorphic to a subdivision of the Dodecahedron, then G has a subgraph isomorphic to a subdivision of P_{10}^- .

We actually prove several results that are more general than (1.1). It turns out that global planarity is not needed for the proof to go through; thus we formulate most of our results in terms of not necessarily planar graphs with a specified set \mathcal{C} of cycles that cover every edge twice, have pairwise empty or connected intersection, and satisfy another natural condition. We call such sets of cycles disk systems. To deduce (1.1) we let \mathcal{C} be the disk system of facial cycles in S . This greater generality allows us to prove an analogue of (1.1) for graphs on higher surfaces.

We also investigate an extension of our original problem to apex graphs. What can we say when G has a set $X \subseteq V(G)$ of size at most k such that $G \setminus X$ is planar, but H has no such set? Is there still an analogue of (1.1)? To prove an exact analogue seems to be a difficult problem that will require a complicated answer. Luckily, for our applications we can assume that G is triangle-free, and we can afford to "sacrifice" a few edges from X to $G \setminus X$. With those two simplifying assumptions we were able to prove (9.9), a result along the lines of (1.1), that is simple enough to allow a concise statement and yet strong

enough to allow us to deduce the desired applications. One such application can be found in [3].

The paper is organized as follows. Throughout the paper we will have to transform one G -subdivision to another, and it will be useful to keep track of the changes we have (or have not) made. There are four kinds of such transformation, called reroutings, and we introduce them in Section 2. In Section 3 we prove a useful and well-known lemma which says that if a graph H has a subgraph isomorphic to a subdivision of a graph G and H is 3-connected, then H has a subgraph isomorphic to a subdivision of G such that all “bridges” are “rigid”. In fact, we need a version of this for graphs that are not necessarily 3-connected. We also review several basic results about planar graphs in Section 3. In Section 4 we introduce disk systems and prove a version of our main result without assuming any connectivity of G or H . In Section 5 we eliminate one of the outcomes by assuming that H is almost 4-connected, and in Section 6 we prove (1.1) and a couple of closely related theorems. In Section 7 we illustrate the use of (1.1). Section 8 contains a technical improvement of one of the earlier lemmas for use in Section 9, where we prove a version of our result when G is at most k vertices away from being planar and H is not.

2. REROUTINGS

We will need a fair amount of different kinds of reroutings that transform one G -subdivision into another, and in order to avoid confusion it seems best to collect them all in one place for easy reference. If P is a path and $x, y \in V(P)$, then xPy denotes the subpath of P with ends x and y .

First we recall the classical notion of a bridge. Let S be a subgraph of a graph H . An S -bridge in H is a connected subgraph B of H such that $E(B) \cap E(S) = \emptyset$ and either $E(B)$ consists of a unique edge with both ends in S , or for some component C of $H \setminus V(S)$ the set $E(B)$ consists of all edges of H with at least one end in $V(C)$. The vertices in $V(B) \cap V(S)$ are called the *attachments* of B .

Let G, H be graphs, let G have no vertices of degree two, let S be a G -subdivision in H , let v be a vertex of S of degree k , let P_1, P_2, \dots, P_k be the segments of S incident with v , and let their other ends be v_1, v_2, \dots, v_k , respectively. Let $x, y \in V(P_1 \cup P_2 \cup \dots \cup P_k)$

be distinct vertices, and let Q be an S -path with ends x and y . Furthermore, let P be a suitable subpath of S , to be specified later. We wish to define a new G -subdivision S' by removing all edges and internal vertices of P from $S \cup Q$. If $x, y \in V(P_1)$, P_1 has length at least two and $P = xP_1y$, then we say that S' is obtained from S by an *I-rerouting*. If, in addition, the S -bridge containing Q has all attachments in P_1 , then we say that S' is obtained from S by a *proper I-rerouting*. See Figure 1. We emphasize that we indeed require that P_1 have at least two edges.




Figure 1. Proper I-rerouting.

Let $k = 3$, let $x \in V(P_1) - \{v\}$, let y be an internal vertex of P_2 , and let $y \in V(xP_1v)$. In those circumstances we say that S' is obtained from S by a *T-rerouting*. See Figure 2.




Figure 2. T-rerouting.

If $k \geq 4$ and there exists an integer $i \in \{1, 2\}$ such that P_i has length at least two, $x, y \in V(P_i)$ and $P = xP_iy$, then we say that S' is obtained from S by a *V-rerouting*, and

we say that it is obtained by a *proper V-rerouting* if all the attachments of the S -bridge containing Q belong to $P_1 \cup P_2$. In that case we say that S' is obtained from S by a *proper V-rerouting based at P_1 and P_2* . Thus a V-rerouting is also an I-rerouting, but not so for proper reroutings.

Let $k \geq 4$, let $x_1, x_2 \in V(P_1) - \{v\}$ and $y_1, y_2 \in V(P_2) - \{v\}$ be distinct vertices such that the vertices x_1, x_2, v, y_1, y_2 appear on the path $P_1 \cup P_2$ in the order listed, and for $i = 1, 2$ let Q_i be an S -path in H with ends x_i and y_i such that Q_1 and Q_2 are disjoint. Let S' be obtained from $S \cup Q_1 \cup Q_2$ by deleting the edges and internal vertices of the paths $x_1 P_1 x_2$ and $y_1 P_2 y_2$. Then S' is a G -subdivision in H , and we say that S' is obtained from S by an *X-rerouting of S* . See Figure 3. It is obtained by a *proper X-rerouting* if the bridges containing Q_1 and Q_2 have all their attachments in $P_1 \cup P_2$. In that case we say that S' is obtained from S by a *proper X-rerouting based at P_1 and P_2* . We say that S' is obtained from S by a *rerouting* if it is obtained from S by an I-rerouting, a V-rerouting, a T-rerouting or an X-rerouting. This relation is not symmetric, because in an I-rerouting and V-rerouting we require that the path that is being changed have length at least two. The distinction among different kinds of rerouting as well as proper reroutings will not be needed until the last two sections and may be safely ignored until then.




Figure 3. X-rerouting.

3. RIGID BRIDGES AND PLANAR GRAPH LEMMAS

Let G be a graph with no vertices of degree two, and let S be a G -subdivision in a graph H . If B is an S -bridge of H , then we say that B is *unstable* if it has at least

one attachment and some segment of S includes all the attachments of B ; otherwise we say that B is *rigid*. Our next lemma, essentially due to Tutte, says that in a 3-connected graph it is possible to make all bridges rigid by changing S using proper I-rerouting only. A *separation* of a graph G is a pair (A, B) of subsets of $V(G)$ such that $A \cup B = V(G)$, and there is no edge between $A - B$ and $B - A$. The *order* of (A, B) is $|A \cap B|$. We say that an S -bridge J is *2-separated* from S if there exists a segment Z of S , two (not necessarily distinct) vertices $u, v \in V(Z)$ and a separation (A, B) of H such that A includes all branch-vertices of S , $V(J \cup uZv) \subseteq B$ and $A \cap B = \{u, v\}$.

(3.1) *Let G be a graph with no vertices of degree two, let H be a graph, and let S be a G -subdivision in H . Then there exists a G -subdivision S' in H obtained from S by a sequence of proper I-reroutings such that every unstable S' -bridge is 2-separated from S' .*

Proof. We use the same argument as in [3, Lemma 2.1], but we give the proof for completeness. We may choose a G -subdivision S' obtained from S by a sequence of proper I-reroutings such that the number of edges that belong to rigid S' -bridges is maximum. We will show that S' is as desired. To that end we may assume that S' has a segment Z such that some S' -bridge that has at least one attachment has all its attachments in Z .

Let v_0, v_1, \dots, v_k be distinct vertices of Z , listed in order of occurrence on Z such that v_0 and v_k are the ends of Z and $\{v_1, \dots, v_{k-1}\}$ is the set of all internal vertices of Z that are attachments of a rigid S' -bridge. We may assume that $k \geq 1$, for otherwise every S' -bridge with all attachments in Z is 2-separated from S' . Now let J be an S' -bridge with at least one attachment and all attachments contained in Z , and let $x, y \in V(Z)$ be the two (not necessarily distinct) attachments of J that maximize xZy . We claim that for $i = 1, 2, \dots, k - 1$ the vertex v_i does not belong to the interior of xZy . To prove this claim suppose to the contrary that v_i belongs to the interior of xZy . Then replacing the path xZy by a subpath of J with ends x and y is a proper I-rerouting that produces a G -subdivision S'' with strictly more edges belonging to rigid S'' -bridges, because every edge that belongs to a rigid S' -bridge belongs to a rigid S'' -bridge, and both edges of S' incident with v_i belong to a rigid S'' -bridge, contrary to the choice of S' . This proves our claim that v_i does not belong to the interior of xZy . Thus there exists an integer

$i = 1, 2, \dots, k$ such that xZy is a subpath of $v_{i-1}Zv_i$. Let B be the union of the vertex-set of $v_{i-1}Zv_i$ and the vertex-sets of all unstable S' -bridges whose attachments are contained in $v_{i-1}Zv_i$, and let $A := V(H) - (B - \{v_{i-1}, v_i\})$. Then the earlier claim implies that (A, B) is a separation, witnessing that J is 2-separated from S , as desired. \square

We will need the following result, a relative of [5, 8, 10, 11, 13]. If G is a graph and $X \subseteq V(G)$, then $G[X]$ denotes the graph $G \setminus (V(G) - X)$.

(3.2) *Let G be a graph, and let C be a cycle in G . Then one of the following conditions holds:*

- (i) *the graph G has a planar embedding in which C bounds a face,*
- (ii) *there exists a separation (A, B) of G of order at most three such that $V(C) \subseteq A$ and $G[B]$ does not have a drawing in a disk with the vertices in $A \cap B$ drawn on the boundary of the disk,*
- (iii) *there exist two disjoint paths in G with ends $s_1, t_1 \in V(C)$ and $s_2, t_2 \in V(C)$, respectively, and otherwise disjoint from C such that the vertices s_1, s_2, t_1, t_2 occur on C in the order listed.*

Proof. The lemma is vacuously true for graphs on at most two vertices. Let G be a graph on at least three vertices, let C be a cycle in G , and assume that the lemma holds for graphs on fewer than $|V(G)|$ vertices. We may assume that G is 3-connected, for otherwise the lemma follows by induction. By [5, Theorem 3.2] either the lemma holds, or there exists a separation (A, B) of G of order at most three such that $V(C) \subseteq A$ and $|B - A| \geq 2$. By moving components of $G \setminus (A \cap B)$ from A to B we may assume that every component of $G \setminus B$ includes at least one vertex of C . We may assume that $G[B]$ can be drawn in a disk with $A \cap B$ drawn on the boundary of the disk, for otherwise the lemma holds. Let G' be obtained from $G[A]$ by adding an edge joining every pair of nonadjacent vertices in $A \cap B$. Then G' satisfies one of (i)–(iii) by the minimality of G . However, since $G[B]$ can be drawn in a disk as specified above, it follows that G satisfies the same conclusion. \square

If G is a subdivision of a 3-connected planar graph, then it has a unique planar embedding by Whitney's theorem [16], and the cycles that bound faces can be characterized

combinatorially. A cycle C in a graph G is called *peripheral* if it is an induced subgraph of G , and $G \setminus V(C)$ is connected. The following three results are well-known [14, 16].

(3.3) *Let G be a subdivision of a 3-connected planar graph, and let C be a cycle in G . Then the following conditions are equivalent:*

- (i) *the cycle C bounds a face in some planar embedding of G ,*
- (ii) *the cycle C bounds a face in every planar embedding of G ,*
- (iii) *the cycle C is peripheral.*

(3.4) *Let G be a subdivision of a 3-connected planar graph, and let C_1, C_2 be two distinct peripheral cycles in G . Then the intersection of C_1 and C_2 is either null, or a one-vertex graph, or a segment.*

(3.5) *Let G be a subdivision of a 3-connected planar graph, let $v \in V(G)$ and let e_1, e_2, e_3 be three distinct edges of G incident with v . If there exist peripheral cycles C_1, C_2, C_3 in G such that $e_i \in E(C_j)$ for all distinct indices $i, j \in \{1, 2, 3\}$, then v has degree three.*

4. DISK SYSTEMS

The preceding theorems summarize all the properties of peripheral cycles that we will require. However, for the sake of greater generality we will be working with sets of cycles satisfying only those axioms that will be needed. Thus we define a *weak disk system* in a graph G to be a set \mathcal{C} of distinct cycles of G such that

- (X0) every edge of G belongs to exactly two members of \mathcal{C} , and
- (X1) the intersection of any two distinct members of \mathcal{C} is either null, or a one-vertex graph, or a segment.

A weak disk system is a *disk system* if it satisfies (X0), (X1) and

- (X2) if e_1, e_2, e_3 are three distinct edges incident with a vertex v of G and there exist disks C_1, C_2, C_3 such that $e_i \in E(C_j)$ for all distinct integers $i, j \in \{1, 2, 3\}$, then v has degree three.

We will refer to the members of a (weak) disk system as *disks*. Thus by (3.3), (3.4) and (3.5) the peripheral cycles of a subdivision of a 3-connected planar graph form a disk system. If G' is obtained from G by (repeated) rerouting, then a weak disk system \mathcal{C} in G induces a weak disk system \mathcal{C}' in G' in the obvious way. We say that \mathcal{C}' is the *weak disk system induced in G' by \mathcal{C}* . If \mathcal{C} is a disk system, then so is \mathcal{C}' .

Let S be a subgraph of a graph H . Let us recall that a path P in H is an *S -path* if it has at least one edge, and its ends and only its ends belong to S . Now let \mathcal{C} be a weak disk system in S . An S -path P is an *S -jump* if no disk includes both ends of P . Let $x_1, x_2, x_3 \in V(S)$, let $x \in V(H) - V(S)$, and let P_1, P_2, P_3 be three paths in H such that P_i has ends x and x_i , they are pairwise disjoint except for x , and each is disjoint from $V(S) - \{x_1, x_2, x_3\}$. Assume further that for each pair x_i, x_j there exists a disk containing both x_i and x_j , but no disk contains all of x_1, x_2, x_3 . In those circumstances we say that the triple P_1, P_2, P_3 is an *S -triad*. The vertices x_1, x_2, x_3 are its *feet*.

Let S be a graph, and let \mathcal{C} be a weak disk system in S . We say that a subgraph J of S is a *detached K_4 -subdivision* if J is isomorphic to a subdivision of K_4 , every segment of J is a segment of S , and each of the four cycles of J consisting of precisely three segments is a disk.

(4.1) *Let G be a graph with no vertices of degree two, let S be a G -subdivision in a graph H , let \mathcal{C} be a weak disk system in S , and let B be an S -bridge with at least two attachments such that no disk includes all attachments of B . Then one of the following conditions holds:*

- (i) *there exists an S -jump, or*
- (ii) *there exists an S -triad, or*
- (iii) *S has a detached K_4 -subdivision J such that the attachments of B are precisely the branch-vertices of J .*

Proof. We may assume that (i) and (ii) do not hold. Let S and B be as stated, and let A be the set of all attachments of B . Thus $|A| \geq 2$. Since G has at least one edge Since (i) does not hold, we deduce that for every pair of elements $a_1, a_2 \in A$ there exists a disk $C \in \mathcal{C}$ such that $a_1, a_2 \in V(C)$. Since (ii) does not hold, we deduce that the same holds

for every triple of elements of A .

Now let $k \geq 3$ be the maximum integer such that for every k -element subset A' of A there exists a disk $C \in \mathcal{C}$ such that $A' \subseteq V(C)$. By hypothesis $k < |A|$, and hence there exist distinct vertices $a_1, a_2, \dots, a_{k+1} \in A$ such that $a_1, a_2, \dots, a_{k+1} \in V(C)$ for no disk $C \in \mathcal{C}$. For $i = 1, 2, \dots, k+1$ let $C_i \in \mathcal{C}$ be a disk in S such that $V(C_i)$ includes all of a_1, a_2, \dots, a_{k+1} except a_i . Then these disks are pairwise distinct. Since a_1 and a_2 belong to both C_3 and C_4 and \mathcal{C} satisfies (X1), there exists a segment P_{12} of S that is a subgraph of $C_3 \cap C_4$ and contains a_1 and a_2 . Similarly, for all distinct integers $i, j = 1, 2, \dots, k+1$, there is a segment P_{ij} of S such that $a_i, a_j \in V(P_{ij})$ and P_{ij} is a subgraph of C_ℓ for all $\ell \in \{1, 2, \dots, k+1\} - \{i, j\}$. Now for all $i = 1, 2, \dots, k+1$ the vertex a_i is an end of P_{ij} , for otherwise the segments P_{ij} ($j \in \{1, 2, \dots, k+1\} - \{i\}$) would be all equal, implying that a_1, a_2, \dots, a_{k+1} all belong to $V(C_t)$ for all $t = 1, 2, \dots, k+1$, a contradiction. Thus a_1, a_2, \dots, a_{k+1} are branch-vertices of S . It follows that $\bigcup P_{ij}$ is a subdivision of a complete graph J . Since $P_{23} \cup P_{24} \cup P_{34}$ is a cycle and it is a subgraph of C_1 , it is equal to C_1 . Similarly for C_2, C_3, C_4 . Hence $k = 3$, and since (i) does not hold we deduce from (X1) that $A = \{a_1, a_2, a_3, a_4\}$. Thus (iii) holds, as desired. \square

In the following definitions let S be a subgraph of a graph H and let \mathcal{C} be a weak disk system in a graph S . Let $C \in \mathcal{C}$, and let P_1 and P_2 be two disjoint S -paths with ends u_1, v_1 and u_2, v_2 , respectively, such that u_1, u_2, v_1, v_2 belong to $V(C)$ and occur on C in the order listed. In those circumstances we say that the pair P_1, P_2 is an S -cross. We also say that it is an S -cross on C . We say that u_1, v_1, u_2, v_2 are the *feet* of the cross. We say that the cross P_1, P_2 is *weakly free* if

(F1) for $i = 1, 2$ no segment of S includes both ends of P_i .

We say that a cross P_1, P_2 is *free* if it satisfies (F1) and

(F2) no two segments of S that share a vertex include all the feet of the cross.

The intent of freedom is that the feet of the cross are not separated from “most of S ” by a separation of order at most three, but it does not quite work that way for our definition. If C is a cycle in S consisting of three segments, then every weakly free cross on C is free. That should be regarded as a drawback of our definition. However, it turns out

that it is not a problem in any of our applications, because in all applications the graph G has girth at least four. On the other hand, there does not seem to be an easy way to eliminate crosses on cycles consisting of three segments, and since we do not need to do it, we chose to avoid it. It should be noted, however, that the “right” definition of freedom should avoid crosses on cycles consisting of three segments.

A separation (X, Y) of H is called an S -separation if the order of (X, Y) is at most three, $X - Y$ includes at most one branch-vertex of S , and the graph $H[X]$ does not have a drawing in a disk with $X \cap Y$ drawn on the boundary of the disk.

We say that \mathcal{C} is *locally planar* in H if for every S -bridge B of H with at least two attachments there exists a disk $C_B \in \mathcal{C}$ such that C_B includes all attachments of B and for every disk $C \in \mathcal{C}$ the graph $C \cup \bigcup B$ has a planar drawing with C bounding the outer face, where the union is taken over all S -bridges B of H with $C_B = C$.

Let Z be a segment of S , let z, w be the ends of Z , and let P_1, P_2 be two disjoint S -paths in H with ends x_1, y_1 and x_2, y_2 , respectively, such that $z, x_1, x_2, y_1, w \in V(Z)$ occur on Z in the order listed, and $y_2 \notin V(Z)$. Let P_3 be a path disjoint from $V(S) - \{y_2\}$ with one end $x_3 \in V(P_1)$ and the other $y_3 \in V(P_2)$ and otherwise disjoint from $P_1 \cup P_2$. We say that the triple P_1, P_2, P_3 is an S -tripod based at Z , and that x_1, y_1, x_2, y_2 (in that order) are its *feet*. We say that zZx_1, y_1Zw and $y_3P_2y_2$ are the *legs* of the tripod. See Figure 4.




Figure 4. An S -tripod.

(4.2) Let G be a graph with no vertices of degree two, let H be a connected graph, and let S be a G -subdivision in H with a weak disk system \mathcal{C} . Then H has a G -subdivision S' obtained from S by repeated I -reroutings such that S' and the weak disk system \mathcal{C}' in S' induced by \mathcal{C} satisfy one of the following conditions:

- (i) there exists an S' -jump, or
- (ii) there exists a weakly free S' -cross in H on some member of \mathcal{C}' , or
- (iii) H has an S' -separation (X, Y) such that $X - Y$ includes no branch-vertex of S' , or
- (iv) S' has a detached K_4 -subdivision J and H has an S' -bridge B such that the attachments of B are precisely the branch-vertices of J , or
- (v) there exists an S' -triad, or
- (vi) the weak disk system \mathcal{C}' is locally planar in H .

Proof. We proceed by induction on $|V(H)|$. Suppose for a contradiction that none of (i)–(vi) holds. We start with the following claim.

- (1) Let S' be a G -subdivision in H obtained from S by repeated I -reroutings, and let \mathcal{C}' be the weak disk system induced in S' by \mathcal{C} . Then for every S' -bridge B of H with at least two attachments there exists a disk $C \in \mathcal{C}'$ such that $V(C)$ includes all attachments of B .

Claim (1) follows from (4.1), for otherwise one of the outcomes (i), (iv), (v) holds, a contradiction. This proves (1).

- (2) There exists a G -subdivision S' in H obtained from S by repeated I -reroutings such that every S' -bridge is rigid.

To prove (2) let S' be as in (3.1). We may assume that there exists an unstable S' -bridge B , for otherwise (2) holds. Let Z be a segment of S that includes all attachments of B . By (3.1) there exist a separation (X, Y) and vertices $x, y \in V(Z)$ such that Y includes every branch-vertex of S , $V(B \cup xZy) \subseteq X$ and $X \cap Y = \{x, y\}$. Since (X, Y) does not satisfy (iii), the graph $H[X]$ has a drawing in a disk with x, y on the boundary of the disk. Let H' be obtained from $H \setminus (X - Y)$ by adding an edge joining x, y if x and y are distinct and not adjacent in H , and let $H' := H \setminus (X - Y)$ otherwise. By induction applied to G , H' and a suitable modification of the graph S we conclude that H' satisfies one

of the conclusions of the lemma. But then H also satisfies the conclusion of the lemma, because if H' satisfies (vi) it follows from the planarity of $H[X]$ that so does H . The other conditions are straightforward. This proves (2).

(3) *There exist a G -subdivision S' in H obtained from S by repeated I -reroutings and an S' -tripod.*

To prove (3) we choose S' as in (2); hence every S' -bridge is rigid. It now follows from (1) that for every S' -bridge B there exists a unique disk C in the weak disk system \mathcal{C}' induced in S' by \mathcal{C} such that all attachments of B belong to $V(C)$. For every disk C of S' let H_C be the union of C and all S' -bridges B whose attachments are included in $V(C)$. Since (vi) does not hold, there exists a disk C of G such that H_C does not have a planar drawing with C bounding the infinite face.

By (3.2) and the fact that (iii) does not hold there exists an S' -cross P_1, P_2 in C . For $i = 1, 2$, let x_i, y_i be the ends of P_i and let B_i be the S' -bridge that includes P_i . We may assume that there is a segment Z of S' such that $x_1, x_2, y_1 \in V(Z)$, for otherwise the S' -cross P_1, P_2 satisfies (F1). We claim that we may assume that $y_2 \notin V(Z)$. Indeed, if $y_2 \in V(Z)$, then since B_1 and B_2 are rigid, there exists a path from $P_1 \cup P_2$, say from P_2 , to a vertex $v \in V(S') - V(Z)$, disjoint from $V(P_1 \cup P_2 \cup S') - \{v\}$. It follows that $P_1 \cup P_2 \cup P$ includes a S' -cross with at least one foot outside Z . Thus we may assume that $y_2 \notin V(Z)$.

If $B_1 = B_2$, then there exists a path P_3 as in the definition of S' -tripod, and hence the claim holds. Thus we may assume that $B_1 \neq B_2$. Since B_1 is rigid there exists a path P_3 in B_1 with one end in $V(P_1) - \{x_1, y_1\}$ and the other end $z \in V(S') - V(Z)$. If $z = y_2$, then P_1, P_2, P_3 is an S -tripod, as desired, and so we may assume that $z \neq y_2$. Then $P_1 \cup P_2 \cup P_3$ includes a weakly free S' -cross, unless some segment Z' of S' includes either z, y_2, y_1 in the order listed, or z, y_2, x_1 in the order listed. By symmetry we may assume the former. Then y_1 is a common end of Z and Z' . Let S'' be obtained from S' by replacing $x_1 Z y_1$ by P_1 ; then $P_3, P_2 \cup x_1 Z x_2$ is a weakly free S'' -cross, as desired. This proves (3).

To complete the proof of the theorem we may select a G -subdivision R in H obtained

from S by repeated I-reroutings and an R -tripod P_1, P_2, P_3 such that the sum of the lengths of the tripod's legs is minimum. Let $Z, z, w, x_1, y_1, x_2, y_2, x_3, y_3$ be as in the definition of tripod.

Let R' be obtained from R by rerouting x_1Zy_1 along P_1 ; then $x_1Zy_1, P_3 \cup y_3P_2y_2, x_2P_2y_3$ is an R' -tripod with the same legs. Thus there is symmetry between $x_1Zy_1 \cup x_2P_2y_3$ and $P_1 \cup P_3$.

Let X' be the vertex-set of $x_1Zy_1 \cup x_2P_2y_3 \cup P_1 \cup P_3$, and let $Y' = V(R) - (X' - \{x_1, y_1, y_3\})$. If there is no path between X' and Y' in $H \setminus \{x_1, y_1, y_3\}$, then there exists a separation (X, Y) of order three in H with $X' \subseteq X$ and $Y' \subseteq Y$ (and hence $X \cap Y = \{x_1, y_1, y_3\}$). Then (X, Y) is an R -separation, and hence (iii) holds, a contradiction. Thus there exists a path P in $H \setminus \{x_1, y_1, y_3\}$ with ends $x \in X'$ and $y \in Y'$. From the symmetry established in the previous paragraph we may assume that $x \in V(P_1) \cup V(P_3) - \{x_1, y_1, y_3\}$. It follows from the minimality of legs that $y \notin V(Z) \cup V(P_2)$.

Let C_1, C_2 be the two disks of R that include Z . Then $y_2 \in V(C_i)$ for some $i = 1, 2$, say $i = 1$, for otherwise P_2 is an R -path satisfying (i). Since $y_2 \notin V(Z)$, (X1) implies that $y_2 \notin V(C_2)$. Since the vertices x_1, y_1, y_2, y are attachments of an R -bridge, by (1) there exists a disk C in G such that $x_1, y_1, y_2, y \in V(C)$. Since $x_1, y_1 \in V(C)$, (X1) implies that $C = C_1$ or $C = C_2$, but $y_2 \notin V(C_2)$, and so $C = C_1$. In particular, $y, y_2 \in V(C_1)$. Since $y \neq y_2$ (because $y \notin V(P_2)$), $P_1 \cup P_2 \cup P_3 \cup P$ includes an R -cross in C_1 satisfying (F1), unless either $z = x_1$ and z, y, x, w appear on C_1 in the order listed, or $y_1 = w$ and w, y, y_2, x appear on C_1 in the order listed. We may therefore assume by symmetry that the former case holds. Let R'' be obtained from R by replacing x_1Zy_1 by P_1 ; then $P_3 \cup y_3P_2y_2, P \cup xZy_1$ is an R'' -cross satisfying (F1), as desired. \square

Our next objective is to improve outcome (ii) of the previous lemma. Let S be a subgraph of H , let C be a cycle in S , and let P_1, P_2 be a weakly free S -cross on C . If the cross P_1, P_2 is not free, then there exist two distinct segments Z_1, Z_2 of S , both subgraphs of C and both incident with a branch-vertex v of S such that $Z_1 \cup Z_2$ includes all the feet of P_1, P_2 . In those circumstances we say that the cross P_1, P_2 is *centered at* v and that it is *based at* Z_1 and Z_2 . We will treat the cases when v has degree three and when it has

degree at least four separately.

We say that an S -triad in a graph H is *local* if there exists a vertex v of S of degree three in S such that each of the three segments of S incident with v includes exactly one foot of the triad. We say that the local S -triad is *centered* at v .

(4.3) *Let G be a graph with no vertices of degree two, let H be a graph, let S be a G -subdivision in H with a weak disk system \mathcal{C} , let $C \in \mathcal{C}$, let $v \in V(C)$ have degree in S exactly three, and let P_1, P_2 be a weakly free S -cross in H on C centered at v . Then there exist a G -subdivision S' obtained from S by exactly one T -rerouting centered at v and a local S' -triad.*

Proof. For $i = 1, 2$ let x_i, y_i be the ends of P_i , and let P_1, P_2 be based at Z_1 and Z_2 . Then we may assume that $x_1, x_2, v \in V(Z_1)$ occur on Z_1 in the order listed; then $y_2, y_1, v \in V(Z_2)$ occur on Z_2 in the order listed. Let S' be the G -subdivision obtained from S by rerouting vZ_2y_2 along P_2 . Then $P_1, y_1Z_2y_2, vZ_1y_1$ is a desired S' -triad. \square

Converting weakly free crosses centered at vertices of degree at least four into free crosses is best done by splitting vertices, but we are concerned with subdivisions, and therefore we take a different route. In the next lemma we need \mathcal{C} to be a disk system (not merely a weak one).

(4.4) *Let G be a graph with no vertices of degree two, let H be a graph, let S be a G -subdivision in H with a disk system \mathcal{C} , and assume that H has a weakly free S -cross centered at a vertex of degree at least four. Then there exists a G -subdivision S' obtained from S by repeated rerouting such that S' and the disk system \mathcal{C}' induced in S' by \mathcal{C} satisfy one the following conditions:*

- (i) H has an S' -jump,
- (ii) H has a free S' -cross on some disk in \mathcal{C}' , or
- (iii) H has an S' -separation (X, Y) such that $X - Y$ includes no branch-vertex of S' .

Proof. Let P_1, P_2 be a weakly free S -cross in H centered at a vertex v of degree at least four. Thus there exist two segments Z_1, Z_2 of S , both incident with v , such that Z_1, Z_2

include all the feet of the cross. For $i = 1, 2$ let x_i, y_i be the ends of P_i . We may assume that $x_1, x_2, v \in V(Z_1)$ occur on Z_1 in the order listed; then $y_2, y_1, v \in V(Z_2)$ and occur on Z_2 in the order listed. For $i = 1, 2$ let v_i be the other end of Z_i and let $L_1 = x_1 Z_1 v_1$ and $L_2 = y_2 Z_2 v_2$.

We may assume that among all triples (S', P'_1, P'_2) , where S' is a G -subdivision obtained from S by repeated rerouting and P'_1, P'_2 is a weakly free S' -cross based at Z'_1, Z'_2 (where Z'_1, Z'_2 are the branches of S' corresponding to Z_1, Z_2), the triple (S, P_1, P_2) is chosen so that $|V(L_1)| + |V(L_2)|$ is minimum.

Let X' be the vertex-set of $P_1 \cup P_2 \cup v Z_1 x_1 \cup v Z_2 y_2$ and let $Y' = V(S) - (X' - \{v, x_1, y_2\})$. If there is no path in $H \setminus \{v, x_1, y_2\}$ with one end in X' and the other in Y' , then there exists a separation (X, Y) of order three with $X' \subseteq X$ and $Y' \subseteq Y$. This separation satisfies (iii), and so we may assume that there exists a path P in $H \setminus \{v, x_1, y_2\}$ with one end $x \in X'$ and the other end $y \in Y'$. From the symmetry we may assume that x belongs to the vertex-set of $P_1 \cup v Z_2 y_2$.

If $y \in V(L_1)$, then replacing P_1 by P if $x \notin V(P_1)$ and by $P \cup x P_1 y_1$ otherwise produces a cross that contradicts the choice of the triple (S, P_1, P_2) . If $y \in V(L_2)$, then replacing $y Z_2 x$ by P if $x \notin V(P_1)$ and replacing $y Z_2 y_1$ by $P \cup x P_1 y_1$ results in a G -subdivision S' obtained from S by a rerouting, and P_1, P_2 can be modified to give a cross P'_1, P'_2 such that the triple (S', P'_1, P'_2) contradicts the choice of (S, P_1, P_2) . Thus $y \notin V(Z_1 \cup Z_2)$.

Let C be the disk that includes both Z_1 and Z_2 (it exists, because P_1, P_2 is a cross), and for $i = 1, 2$ let C_i be the other disk that includes Z_i . If $y \in V(C)$, then $P_1 \cup P_2 \cup P$ includes a free cross, and so (ii) holds. Thus we may assume that $y \notin V(C)$. Similarly, if $y \notin V(C_2)$, then $P_1 \cup P$ includes an S -jump with one end y and the other end x or y_1 , and so we may assume that $y \in V(C_2)$. Since v has degree at least four, (X1) and (X2) imply that $V(C_1) \cap V(C_2) = \{v\}$. It follows that $y \notin V(C_1)$. Now let S' be obtained from S by an X-rerouting using the cross P_1, P_2 , and let Z'_1, Z'_2 be the segments of S' corresponding to Z_1, Z_2 , respectively. Thus $Z'_1 = v_1 Z_1 x_1 \cup P_1 \cup y_1 Z_2 v$ and $Z'_2 = v_2 Z_2 y_2 \cup P_2 \cup x_2 Z_1 v$. Now $P \cup x Z_2 y_1$ includes an S' -jump with one end y and the other end in the interior of Z'_1 , and so (i) holds. \square

We can summarize some of the lemmas of this section as follows.

(4.5) *Let G be a connected graph with no vertices of degree two that is not the complete graph on four vertices, let H be a graph, and let S be a G -subdivision in H with a disk system \mathcal{C} . Then H has a G -subdivision S' obtained from S by repeated reroutings such that S' and the weak disk system \mathcal{C}' induced in S' by \mathcal{C} satisfy one of the following conditions:*

- (i) *there exists an S' -jump in H , or*
- (ii) *there exists a free S' -cross in H on some disk of S' , or*
- (iii) *H has an S' -separation (X, Y) such that $X - Y$ includes no branch-vertex of S' , or*
- (iv) *there exists an S' -triad, or*
- (v) *the disk system \mathcal{C}' is locally planar in H .*

Proof. By (4.2) there exists a G -subdivision S_1 obtained from S by a sequence of reroutings such that one of the outcomes of that lemma holds. But (4.2)(iv) does not hold, because \mathcal{C} satisfies (X2) and G is not K_4 . We may assume therefore that (4.2)(ii) holds, for otherwise S_1 and the weak disk system induced in S_1 by \mathcal{C} satisfy (4.5). Thus S_1 has a disk C and a weakly free cross P_1, P_2 on C . We may assume that P_1, P_2 is not free, for otherwise (4.5)(ii) holds. Thus there exists a branch-vertex v of S_1 that belongs to C and two distinct segments Z_1, Z_2 of S_1 , both subgraphs of C and both incident with v such that the cross P_1, P_2 is centered at v and based at Z_1, Z_2 . If v has degree three in S_1 , then the lemma holds by (4.3) and if v has degree at least four, then the lemma holds by (4.4). \square

The following theorem will be used in [2]. Recall that a graph G is almost 4-connected if it is 3-connected, has at least five vertices, and, for every separation (A, B) of G of order 3, one of $A - B, B - A$ contains at most one vertex.

(4.6) *Let G be a graph with no vertices of degree two that is not the complete graph on four vertices, let H be an almost 4-connected graph, and let S be a G -subdivision in H with a disk system \mathcal{C} . Then H has a G -subdivision S' obtained from S by repeated reroutings such that S' and the disk system \mathcal{C}' induced in S' by \mathcal{C} satisfy one of the following conditions:*

- (i) there exists an S' -jump in H , or
- (ii) there exists a free S' -cross in H on some disk of S' , or
- (iii) there exists an S' -triad, or
- (iv) the disk system \mathcal{C}' is locally planar in H .

Proof. By (4.5) we may assume that there exists a G -subdivision S' obtained from S by repeated reroutings such that S' and the weak disk system \mathcal{C}' induced in S' by \mathcal{C} satisfy (4.5)(iii), for otherwise the theorem holds. Thus H has an S' -separation (X, Y) such that $X - Y$ includes no branch-vertex of S' . Since S has at least five branch-vertices, it follows that $|Y - X| \geq 2$. But $|X - Y| \geq 2$, because $H[X]$ cannot be drawn in a disk with $X \cap Y$ drawn on the boundary of the disk. This contradicts the almost 4-connectivity of H . \square

5. TRIADS

In this section we improve outcome (iv) of (4.5). A graph G is *internally 4-connected* if it is 3-connected and for every separation (A, B) of order three one of $G[A], G[B]$ has at most three edges. (Thus every 4-connected graph is internally 4-connected and every internally 4-connected graph is almost 4-connected.) If S is a G -subdivision in a graph H , then there is a mapping η that assigns to each $v \in V(G)$ the corresponding vertex $\eta(v) \in V(S)$, and to every edge $e \in E(G)$ the corresponding path $\eta(e)$ of S . We say that η is a *homeomorphic embedding*, and we write $\eta : G \hookrightarrow S \subseteq H$ to denote that fact that η is a homeomorphic embedding that maps G onto the subgraph S of H .

(5.1) *Let G be an almost 4-connected graph, let H be a graph, let S be a G -subdivision in H with a weak disk system \mathcal{C} , and assume that there exists an S -triad in H that is not local and has set of feet F . Assume also that if $|V(G)| \leq 6$, then G is internally 4-connected. Then*

- (i) there exists a segment of S with both ends in F , or
- (ii) $S \setminus F$ is connected and F is an independent set in S .

Proof. Let the S -triad be Q_1, Q_2, Q_3 , and let $F = \{x_1, x_2, x_3\}$ be labeled so that x_i is an end of Q_i . Let J be the subgraph of S with vertex-set F and no edges. We may assume that S has at least two J -bridges, for otherwise (ii) holds. Assume first that some J -bridge of S includes no branch-vertex of S , except possibly as an attachment. Then that J -bridge is a subgraph of a segment Z that includes two members of F , say x_1 and x_2 . It follows that x_1 and x_2 are the ends of Z , for if x_1 is an internal vertex of Z , then the disk containing x_1 and x_3 contains x_2 as well, a contradiction. Hence (i) holds.

Thus we may assume that every J -bridge of S includes a branch-vertex of S that is not an attachment of J , and since there are at least two J -bridges of S , it follows that S has separation (X, Y) with $X \cap Y = \{x_1, x_2, x_3\}$ such that both $X - Y$ and $Y - X$ include a branch-vertex of S .

We claim that one of $X - Y, Y - X$ includes at most one branch-vertex of S . To prove this claim, suppose the contrary and let $\eta : G \hookrightarrow S \subseteq H$ be a homeomorphic embedding. Let $z_1, z_2, z_3 \in V(G) \cup E(G)$ be defined as follows. Let $i \in \{1, 2, 3\}$. If x_i is a branch-vertex of S , then let $z_i \in V(G)$ be such that $\eta(z_i) = x_i$; otherwise x_i is the interior vertex of a unique segment $\eta(z_i)$ of S , and we define z_i that way. Let X' be the set of all vertices x of G such that $\eta(x) \in X$, and let Y' be defined analogously. Then $X' \cup Y' = V(G)$, and there are exactly $3 - |X' \cap Y'|$ edges of G with one end in $X' - Y'$ and the other in $Y' - X'$. Note that $X' \cap Y' = \{z_1, z_2, z_3\} \cap V(G)$. If $z_1, z_2, z_3 \in V(G)$, then our claim follows from the almost 4-connectivity of G . For the next case assume that $z_1 \in E(G)$ and $z_2, z_3 \in V(G)$, and let u_1, v_1 be the ends of z_1 with $u_1 \in X'$ and $v_1 \in Y'$. By the almost 4-connectivity of G applied to the separation $(X' \cup \{v_1\}, Y')$ we deduce that $|Y' - X' - \{v_1\}| \leq 1$, and, by symmetry, $|X' - Y' - \{u_1\}| \leq 1$. Thus $|V(G)| \leq 6$, and hence G is internally 4-connected. Since G has at least five vertices we may assume that $X' - Y' - \{u_1\}$ is not empty, say $x \in X' - Y' - \{u_1\}$. Then x has neighbors u_1, z_2, z_3 . Since u_1 has degree at least three, it is adjacent to z_2 or z_3 , and hence x has degree three and belongs to a triangle, contrary to the internal 4-connectivity of G . Thus, the claim holds when at most one of z_1, z_2, z_3 is an edge. The other two cases are similar, and are omitted. This completes the proof of our claim that one of $X - Y, Y - X$ includes exactly one branch-vertex of G . From the symmetry we may assume that $X - Y$ includes exactly one branch-vertex of S , say v . It

follows that v has degree three and that Q_1, Q_2, Q_3 is a local triad, a contradiction. \square

If G is internally 4-connected and planar we have the following corollary.

(5.2) *Let G be an internally 4-connected planar graph, let H be a graph, let S be a G -subdivision in H , and let \mathcal{C} be the disk system in S consisting of peripheral cycles of S . Then every S -triad is local.*

Proof. Let F be the set of feet of an S -triad, and let us assume for a contradiction that the S -triad is not local. Let us fix a drawing of S in the sphere. Since every pair of vertices in F are cofacial by (3.3), there exists a simple closed curve ϕ intersecting S precisely in the set F and such that both disks bounded by ϕ include a branch-vertex of S . Thus (5.1)(ii) does not hold, and (5.1)(i) does not hold by the internal 4-connectivity of G and the fact that the S -triad is not local. That contradicts (5.1). \square

(5.3) *Let G, H be graphs, where G has no vertices of degree two, let S be a G -subdivision in H , let \mathcal{C} be a weak disk system in S , and let Q_1, Q_2, Q_3 be an S -triad in H such that two of its feet are the ends of a segment Z of S . Then there exist a G -subdivision S' in H obtained from S by I-rerouting the segment Z and an S' -jump.*

Proof. Let the feet of the S -triad be x_1, x_2, x_3 be numbered so that x_i is an end of Q_i , and let Z have ends x_1 and x_2 . Let S' be obtained from S by replacing Z by $Q_1 \cup Q_2$. Then Q_3 is an S' -jump, for if its ends belong to a disk of S' , then the corresponding disk of S would include all of x_1, x_2, x_3 , contrary to the definition of an S -triad. This proves the lemma. \square

Let G, H be graphs, let G have no vertices of degree two, let S be a G -subdivision in H , let $v \in V(S)$ have degree three in S , let Z_1, Z_2, Z_3 be the three segments of S incident with v , and let Q_1, Q_2, Q_3 be a local S -triad centered at v with feet x_1, x_2, x_3 , where $x_i \in V(Z_i)$. Let S' be the G -subdivision obtained from $S \cup Q_1 \cup Q_2 \cup Q_3$ by deleting v and all the edges and internal vertices of the paths $x_i Z_i v$ for $i = 1, 2, 3$. We say that S' was obtained from S by a *triad exchange*. It follows that $x_1 Z_1 v, x_2 Z_2 v, x_3 Z_3 v$ is an S' -triad.

(5.4) Let G, H be graphs, where G has no vertices of degree two, let S be a G -subdivision in H , let \mathcal{C} be a weak disk system in S , and let Q_1, Q_2, Q_3 be a local S -triad in H . Then there exists a G -subdivision S' obtained from S by repeated rerouting and at most one triad exchange such that S' and the weak disk system \mathcal{C}' in S' induced by \mathcal{C} satisfy one of the following conditions:

- (i) there exists an S' -jump in H , or
- (ii) there exists a free S' -cross on some member of \mathcal{C}' , or
- (iii) there exists an S' -separation in H .

Proof. Let the triad Q_1, Q_2, Q_3 be centered at v , let its feet be x_1, x_2, x_3 , let Z_1, Z_2, Z_3 be the three segments of S incident with v numbered so that $x_i \in V(Z_i)$, and let v_i be the other end of Z_i . Let L_i be the subpath of Z_i with ends v_i and x_i , and let P_i be the subpath of Z_i with ends v and x_i . We say that the paths L_1, L_2, L_3 are the legs of the S -triad Q_1, Q_2, Q_3 . We may assume that S and Q_1, Q_2, Q_3 are chosen so that there is no G -subdivision of H with a triad as above obtained from S by a rerouting such that the sum of the lengths of its legs is strictly smaller than $|E(L_1)| + |E(L_2)| + |E(L_3)|$. Let $X_1 = V(P_1 \cup P_2 \cup P_3 \cup Q_1 \cup Q_2 \cup Q_3)$ and $Y_1 = V(S) - (X_1 - \{x_1, x_2, x_3\})$. If $H \setminus \{x_1, x_2, x_3\}$ has no path between X_1 and Y_1 , then H has a separation (X, Y) such that $X \cap Y = \{x_1, x_2, x_3\}$, $X_1 \subseteq X$, and $Y_1 \subseteq Y$. Then (X, Y) satisfies (iii), as desired.

We may therefore assume that there exists a path P as above. Let the ends of P be $x \in X_1 - \{x_1, x_2, x_3\}$ and $y \in Y_1 - \{x_1, x_2, x_3\}$. We may assume that P has no internal vertex in $X_1 \cup Y_1$. We claim that $y \notin V(L_1 \cup L_2 \cup L_3)$. Indeed, if $x \in V(Q_1 \cup Q_2 \cup Q_3)$, then $y \notin V(L_1 \cup L_2 \cup L_3)$ by the choice of Q_1, Q_2, Q_3 (no change of S needed). So we may assume that $x \in V(P_1 \cup P_2 \cup P_3)$ and $y \in V(L_1 \cup L_2 \cup L_3)$. Now replacing a path of $P_1 \cup P_2 \cup P_3$ by P is an I-rerouting or T-rerouting, and the resulting G -subdivision S' has an S' -triad that contradicts the choice of S and Q_1, Q_2, Q_3 . Thus $y \notin V(Z_1 \cup Z_2 \cup Z_3)$.

The operation of triad exchange exchanges the roles of $P_1 \cup P_2 \cup P_3$ and $Q_1 \cup Q_2 \cup Q_3$. Thus by applying the triad exchange operation if needed we gain symmetry between $P_1 \cup P_2 \cup P_3$ and $Q_1 \cup Q_2 \cup Q_3$. Thus may assume that $x \in V(P_1 \cup P_2 \cup P_3)$. We may assume that S and P do not satisfy (i), and hence there exists a disk C in S such that

$x, y \in V(C)$. It follows that C includes two of the segments incident with v , say Z_1 and Z_2 . We may assume that $Q_1 \cup Q_2, P$ is not a free S -cross in C for otherwise (ii) holds, and hence $v_1 = x_1, v_2 = x_2$ and there is a segment Z of S with ends v_1 and v_2 . Let S' be the G -subdivision obtained from S by the triad exchange that replaces Q_1, Q_2, Q_3 by P_1, P_2, P_3 . Then $P \cup P_1 \cup P_2 \cup P_3$ includes an S' -path with ends x_3 and y . We may assume that this path is not an S' -jump, for otherwise (i) holds. Thus there exists a disk C' in S' that includes Z and x_3 , and hence includes all of x_1, x_2, x_3 , contrary to the fact that Q_1, Q_2, Q_3 is a triad. \square

The results thus far can be summarized as follows.

(5.5) *Let G be an almost 4-connected graph, let H be a graph, and let S be a G -subdivision in H with a disk system \mathcal{C} . Assume that if G has at most six vertices, then it is internally 4-connected. Then H has a G -subdivision S' obtained from S by repeated reroutings and possibly one triad exchange such that S' and the disk system \mathcal{C}' induced in S' by \mathcal{C} satisfy one of the following conditions:*

- (i) *there exists an S' -jump in H , or*
- (ii) *there exists a free S' -cross in H on some disk of S' , or*
- (iii) *H has an S' -separation, or*
- (iv) *there exists an S' -triad with set of feet F such that $S' \setminus F$ is connected and F is an independent set in S' .*
- (v) *the disk system \mathcal{C}' is locally planar in H .*

Proof. Let S' be as in (4.5), and let \mathcal{C}' be the corresponding disk system in S' . We may assume that (4.5)(iv) holds, for otherwise the lemma holds. Let t be an S' -triad. If t is local, then the result holds by (5.4). Otherwise by (5.1) and (5.3)(i) either outcome (i) or outcome (iv) holds. \square

6. WHEN G IS PLANAR

We are now ready to reformulate the above results in terms of embedded graphs. By a *surface* we mean a compact connected 2-dimensional manifold with no boundary. A graph S embedded in a surface Σ is *polyhedrally embedded* if S is a subdivision of a 3-connected graph and every homotopically non-trivial simple closed curve intersects the graph at least three times. It follows that the face boundaries of S form a disk system, say \mathcal{C} . Now if S' is another G -subdivision obtained from S by rerouting or triad exchange then S uniquely determines an embedding of S' in Σ (up to a homotopic shift) and the disk system induced in S' by \mathcal{C} consists of the face boundaries in S' .

(6.1) *Let G be an almost 4-connected graph, let H be a graph, let S be a G -subdivision in H , polyhedrally embedded in a surface Σ , and assume that S does not extend to an embedding of H . Assume also that if G has at most six vertices, then it is internally 4-connected. Then there exists a G -subdivision S' in H obtained from S by repeated reroutings and at most one triad exchange such that one of the following conditions holds for the induced embedding of S' into Σ :*

- (i) *there exists an S' -path in H such that no face boundary of S' includes both ends of the path,*
- (ii) *there exists a free S' -cross on some face boundary of S' , or*
- (iii) *H has an S' -separation, or*
- (iv) *there exist an independent set $F \subseteq V(S')$ of size three, a non-separating simple closed curve in Σ intersecting S' precisely in F , and an S' -triad in H with set of feet F such that $S' \setminus F$ is connected.*

Proof. Let \mathcal{C} be the disk system described prior to the statement of (6.1). By (5.5) there exists a G -subdivision S' in H obtained from S by repeated reroutings and at most one triad exchange that satisfies one of (i)–(vi) of that lemma. If (i), (ii) or (iii) holds, then our lemma holds. Condition (5.5)(v) does not hold, because S does not extend to an embedding of H . Thus we may assume that (5.5)(iv) holds. Let x_1, x_2, x_3 be the feet of the triad; since every pair of x_1, x_2, x_3 belong to a common face boundary, there exists a simple closed curve ϕ passing through x_1, x_2, x_3 and those faces. Since no face boundary

of S' includes all of x_1, x_2, x_3 and $S' \setminus \{x_1, x_2, x_3\}$ is connected, it follows that ϕ does not separate Σ . Thus (iv) holds. \square

From now on we will be working exclusively with disk systems consisting of peripheral cycles in subdivisions of 3-connected planar graphs, and so the notions such as S -jump or S -cross will refer to the disk system consisting of all peripheral cycles. If G is planar, then there is no non-separating closed curve, and hence condition (iv) from the above theorem cannot hold. Thus we have the following corollary for planar graphs. The corollary is used in [1].

(6.2) *Let G be an almost 4-connected planar graph, let H be a non-planar graph, and let S be a G -subdivision in H . Assume also that if G has at most six vertices, then G is internally 4-connected. Then there exists a G -subdivision S' in H obtained from S by repeated reroutings and at most one triad exchange such that S' and the disk system of peripheral cycles in S' satisfy one of the following conditions:*

- (i) *there exists an S' -path in H such that no peripheral cycle of S' includes both ends of the path,*
- (ii) *there exists a free S' -cross on some peripheral cycle of S' , or*
- (iii) *H has an S' -separation.*

Finally, we prove (1.1), which we restate in a slightly stronger form.

(6.3) *Let G be an almost 4-connected planar graph, let H be an almost 4-connected non-planar graph, and let S be a G -subdivision in H . Assume that if $|V(G)| \leq 6$, then G is internally 4-connected. Then there exists a G -subdivision S' in H obtained from S by repeated reroutings and at most one triad exchange such that S' and the disk system of peripheral cycles in S' satisfy one of the following conditions:*

- (i) *there exists an S' -jump in H , or*
- (ii) *there exists a free S' -cross in H on some peripheral cycle of S' .*

Proof. Let G, H, S be as stated. By (6.2) there exists a G -subdivision S' in H obtained from S by repeated reroutings and at most one triad exchange such that one of the conclusions

of (6.2) holds. We may assume that H has an S' -separation (X, Y) , for otherwise the lemma holds. Then $|X - Y| \geq 2$, because $H[X]$ cannot be drawn in a disk with $X \cap Y$ drawn on the boundary of the disk. The set $X - Y$ includes at most one branch-vertex of S' by the definition of S' -separation. We claim that $|Y - X| \geq 2$. This is clear if S' has at least six branch-vertices; otherwise G has exactly five vertices and hence is internally 4-connected. It follows that X cannot include four branch-vertices of S' , and so $|Y - X| \geq 2$, as claimed. But that contradicts the almost 4-connectivity of H . \square

We need the following lemma. Let G be a subdivision of a 3-connected planar graph, and let x, y be vertices or edges of G . We say that x, y are *cofacial* if some peripheral cycle in G includes both x and y .

(6.4) *Let G be an internally 4-connected planar graph, and let $e \in E(G)$ and $v \in V(G)$ be not cofacial. Then at least one end of e is not cofacial with v .*

Proof. Let us fix a planar drawing of G , and suppose for a contradiction that both ends of e are cofacial with v . By (3.3) there exists a simple closed curve in the plane that passes through v , the two ends of e , and is otherwise disjoint from G . Since v and e are not cofacial this curve disconnects G , contrary to the internal 4-connectivity of G . \square

We also need the following analogue of (6.4).

(6.5) *Let G be an internally 4-connected planar graph, and let $e, f \in E(G)$ be not cofacial. Then some end of e is not cofacial with some end of f .*

Proof. Let u_1, u_2 be the ends of e . By (6.4) it suffices to show that one of u_1, u_2 is not cofacial with f . Thus we may assume for a contradiction that there exist peripheral cycles C_1, C_2 in G , both containing f and such that $u_i \in V(C_i)$. Let us fix a drawing of G in the plane. By (3.3) there exists a simple closed curve intersecting the graph G three times: in u_1, u_2 and in an internal point of f . However, that contradicts the internal 4-connectivity of G . \square

If we allow contracting edges and G has no peripheral cycles of length three, then (6.3) can be further simplified. The next result is a restatement of (1.2), because every triangle-free almost 4-connected graph is internally 4-connected.

(6.6) *Let G be a triangle-free internally 4-connected planar graph, and let H be an almost 4-connected non-planar graph such that H has a subgraph isomorphic to a subdivision of G . Then there exists a graph G' such that G' is isomorphic to a minor of H , and either*

- (i) $G' = G + uv$ for some vertices $u, v \in V(G)$ such that no peripheral cycle of G contains both u and v , or
- (ii) $G' = G + u_1v_1 + u_2v_2$ for some distinct vertices $u_1, u_2, v_1, v_2 \in V(G)$ such that u_1, u_2, v_1, v_2 appear on some peripheral cycle of G in the order listed.

Proof. By (6.3) there exist a homeomorphic embedding $\eta : G \hookrightarrow S \subseteq H$ and either an S -jump or a free S -cross. Assume first that P is an S -jump with ends a and b . If both a and b are branch-vertices, then (i) holds. Let us assume that a is a branch-vertex, say $a = \eta(v)$ and that b belongs to the interior of $\eta(e)$ for some edge $e \in E(G)$. Since P is an S -jump it follows that v and e are not cofacial. By (6.4) there exists an end u of e such that u and v are not cofacial. Then $G + (u, v)$ satisfies (i). We may therefore assume that neither a nor b is a branch-vertex. Let a be an internal vertex of $\eta(f)$ and let b be an internal vertex of $\eta(e)$, where $e, f \in E(G)$ are not cofacial. By (6.5) there is an end u of e that is not cofacial with an end v of f . It follows that $G + (u, v)$ satisfies (i).

We may therefore assume that P_1, P_2 is a free S -cross in H on some peripheral cycle C . Let F be the set of feet of this cross, and let B be the set of branch-vertices of H that belong to C . We define a bipartite graph L with bipartition (F, B) by saying that $f \in F$ is adjacent to $b \in B$ if some subpath of C has ends f and b and includes no vertex of $F \cup B$ in its interior. Since C has at least four branch-vertices and P_1, P_2 is a free cross, Hall's theorem implies that L has a complete matching from F to B . Let F be matched into $\{\eta(u_1), \eta(u_2), \eta(v_1), \eta(v_2)\}$, where $\eta(C_0) = C$ and u_1, u_2, v_1, v_2 occur on C_0 in the order listed. Then $G + u_1v_1 + u_2v_2$ satisfies (ii), as desired. \square

7. AN APPLICATION

By the *cube* we mean the graph of the 1-skeleton of the 3-dimensional cube. As an application of the results of this paper we examine non-planar graphs that have a subgraph isomorphic to a subdivision of the cube. Other applications appeared in [1, 12]. Let W denote the graph obtained from the cube by adding an edge joining two vertices at distance three, and let V_8 be the graph obtained from a cycle of length eight by adding edges joining every pair of diagonally opposite vertices. See Figure 5. The following result is a relative of [6].




Figure 5. The graphs W and V_8 .

(7.1) *Let H be an almost 4-connected non-planar graph that has a subgraph isomorphic to a subdivision of the cube. Then H has a subgraph isomorphic to a subdivision of V_8 or W .*

Proof. Let K denote the cube. By (6.3) there exists a homeomorphic embedding $\eta : K \hookrightarrow S \subseteq H$ such that (i) or (ii) of (6.3) holds. Suppose first that (i) holds, and let P be a path as in (i) with ends x and y . If $\eta(u) = x$ and $\eta(v) = y$, where $u, v \in V(K)$ are at distance three in K , then η can be extended to yield a W -subdivision in H , and the theorem holds. Otherwise it is easy to see that η can be extended to produce a V_8 -subdivision in H .

If (ii) holds, then there exists a free η -cross in some cycle $\eta(C)$ of S , where C is a cycle of K of length four. Let the vertices of C be v_1, v_2, v_3, v_4 , in order. Let K' be obtained from K by deleting the edges v_1v_2 and v_3v_4 , and adding the edges v_1v_3 and v_2v_4 . The existence of the free cross implies that H has a subgraph isomorphic to a subdivision of K' . But K' is isomorphic to V_8 , and so the result holds. \square

8. IMPROVING LEMMA (4.2)

In the rest of the paper all disk systems will consist of peripheral cycles of subdivisions of 3-connected planar graphs. The following is a version of (5.4), but without rerouting or a triad exchange.

(8.1) *Let G, H be graphs, where G is internally 4-connected and planar and is not isomorphic to the cube, let S be a G -subdivision in H , let \mathcal{C} be the disk system of peripheral cycles in S , and let Q_1, Q_2, Q_3 be a local S -triad in H centered at $v \in V(S)$ such that the S -bridge containing $Q_1 \cup Q_2 \cup Q_3$ has an attachment y that does not belong to any of the segments incident with v . Then there exists an S -jump in H with one end y .*

Proof. Let Z_1, Z_2, Z_3 be the three segments incident with v , let v_i be the other end of Z_i , and let x_1, x_2, x_3 be the feet of the triad Q_1, Q_2, Q_3 numbered so that $v_i \in V(Z_i)$. Let us fix a drawing of S in the sphere. For distinct integers $i, j, k \in \{1, 2, 3\}$ let f_i be the face of S incident with Z_j and Z_k . By hypothesis there exists a path P with ends $x \in V(Q_1 \cup Q_2 \cup Q_3) - \{x_1, x_2, x_3\}$ and $y \in V(S) - V(Z_1 \cup Z_2 \cup Z_3)$, disjoint from $S \setminus y$. We may assume that for all $i = 1, 2, 3$ the vertices y and x_i are incident with the same face of S , for otherwise the lemma holds; let g_i denote that face. Let $i, j, k \in \{1, 2, 3\}$ be distinct. There exists a simple closed curve ϕ_k that intersects S in $\{y, v_i, v_j\}$ and is otherwise contained in $g_i \cup g_j \cup f_k$. By the internal 4-connectivity of G one of the disks bounded by ϕ_k includes at most one branch-vertex of S . Let u_k denote that branch-vertex if it exists; otherwise u_k is undefined and $g_i = g_j = f_k$. It follows that G has at most eight vertices; the corresponding branch-vertices of S are v, v_1, v_2, v_3, y and a subset of $\{u_1, u_2, u_3\}$. Since v_1 has degree at least three, we deduce that at least one of u_1, u_2, u_3

exists, say u_3 does. Then no segment of S has ends y and v_1 , or y and v_2 , or v_1 and v_3 , or v_2 and v_3 , by the internal 4-connectivity of G . Since v_1 and v_2 have degree at least three, it follows that u_1 and u_2 also exist. Thus G is isomorphic to the cube, a contradiction. \square

We need to prove a variant of (4.2), where rerouting is not used. First we need a definition. Let S be a subdivision of a 3-connected planar graph, let W be a segment of S , let z, w be the ends of W , and let P_1, P_2 be two disjoint S -paths in H with ends x_1, y_1 and x_2, y_2 , respectively, such that $z, x_1, x_2, y_1, w \in V(W)$ occur on W in the order listed, and $y_2 \notin V(W)$. Let P_3 be a path disjoint from $V(S) - \{y_2\}$ with one end $x_3 \in V(P_1)$ and the other $y_3 \in V(P_2)$ and otherwise disjoint from $P_1 \cup P_2$. Thus P_1, P_2, P_3 is an S -tripod based at W . Let \mathcal{C} be the disk system in S consisting of peripheral cycles, and let C, C' be the two disks that contain W . Let $y_2 \in V(C) - V(C')$, and let P_4 be an S -path with ends x_4 and y_4 , where x_4 belongs to the interior of x_1Wy_1 and $y_4 \in V(C') - V(C)$, such that no internal vertex of P_4 belongs to $P_1 \cup P_2 \cup P_3$. For $i = 1, 2$ let B_i be the S -bridge of H that includes P_i . Let us assume further that

- all attachments of B_1 and B_2 belong to C ,
- every S -bridge other than B_1 or B_2 that has an attachment in the interior of x_1Wy_1 has all its attachments in $V(C') \cup \{y_2\}$, and
- if $B_1 \neq B_2$, then for $i = 1, 2$ the vertex y_2 is the only attachment of B_i that does not belong to W .

In those circumstances we say that the quadruple P_1, P_2, P_3, P_4 is an S -tunnel. It is worth noting that if $B_1 \neq B_2$, then $y_2 = y_3$. See Figure 6.

(8.2) *Let G be an internally 4-connected planar graph, let H be a graph, and let S be a G -subdivision in H such that every unstable S -bridge is 2-separated from S . Then one of the following conditions holds:*

- (i) *there exists an S -jump, or*
- (ii) *there exists a weakly free S -cross in H , or*
- (iii) *H has an S -separation (X, Y) such that $X - Y$ includes no branch-vertex of S , or*
- (iv) *there exists an S -triad, or*




Figure 6. An S -tunnel.

(v) *there exists an S -tunnel, or*

(vi) *the graph H is planar.*

Proof. We proceed by induction on $|V(G)|$. Suppose for a contradiction that none of (i)–(vi) holds. As in Claim (2) of (4.2) we may assume that every S -bridge is rigid, for otherwise the lemma follows by induction. Now since every S -bridge is rigid, it follows from (4.1) and the fact that (i) and (iv) do not hold that for every S -bridge B there exists a unique disk C such that all attachments of B belong to $V(C)$. For every disk C of S let H_C be the union of C and all S -bridges B whose attachments are included in $V(C)$. Since (vi) does not hold, there exists a disk C of S such that H_C does not have a planar drawing with C bounding the infinite face. The same argument as in the proof of Claim (3) of (4.2) shows that there exists an S -tripod.

Let us select a segment Z and vertex $y_2 \notin V(Z)$ such that there exists an S -tripod P_1, P_2, P_3 based at Z with feet x_1, y_1, x_2, y_2 . Let $x_3 \in V(P_1)$ and $y_3 \in V(P_2)$ be the ends of P_3 ; we say that $y_3 P_2 y_2$ is the *leg* of the S -tripod. Let us, in addition, select an S -tripod based at Z so that its leg is minimal. Let the leg be L . We say that a vertex $z \in Z$ is *sheltered* if z is an internal vertex of $x_1 Z y_1$ for some S -tripod based at Z with

feet x_1, y_1, x_2, y_2 and leg L , and we say that the tripod *shelters* the vertex z . Now let $x'_1, y'_1 \in V(Z)$ be not sheltered but such that every internal vertex of $x'_1 Z y'_1$ is sheltered, and let X' be the union of $x'_1 Z y'_1$ and $V(P_1 \cup x_2 P_2 y_3 \cup P_3)$, over all S -tripods P_1, P_2, P_3 with leg L that shelter an internal vertex of $x'_1 Z y'_1$.

Let $Y' = V(S \cup L) - (X' - \{x'_1, y'_1, y_3\})$. If there is no path between X' and Y' in $H \setminus \{x'_1, y'_1, y_3\}$, then there exists a separation (X, Y) of order three in H with $X' \subseteq X$ and $Y' \subseteq Y$ (and hence $X \cap Y = \{x'_1, y'_1, y_3\}$). Then (X, Y) is an S -separation, and hence (iii) holds, a contradiction. Thus there exists a path P in $H \setminus \{x_1, y_1, y_3\}$ with ends $x \in X'$ and $y \in Y'$. We may assume that P has no internal vertex in $X' \cup Y'$; thus P has no internal vertex in S . If $x \in V(Z)$ let P_1, P_2, P_3 be an S -tripod that shelters x ; otherwise let P_1, P_2, P_3 be an S -tripod that shelters some vertex of $x'_1 Z y'_1$ such that $x \in V(P_1 \cup P_2 \cup P_3)$.

We claim that P may be chosen so that $y \notin V(Z) \cup V(L)$. It is clear that $y \notin V(L)$ by the choice of L , and so we may assume that $y \in V(Z)$. Let B be the S -bridge that includes P . If B includes at least one of P_1, P_2, P_3 , then $B \cup P_1 \cup P_2 \cup P_3$ includes an S -tripod that shelters x'_1 or y'_1 , a contradiction. The same conclusion holds (or we obtain contradiction to the minimality of L) if the only attachment of B outside Z is y_2 . Thus we may assume that B has an attachment in $V(S) - V(Z) - \{y_2\}$, and so P may be replaced by a path with an end not in $V(Z) \cup V(L)$. This proves our claim that we may assume that $y \notin V(Z) \cup V(L)$.

Let C_1, C_2 be the two disks of S that include Z . Then $y_2 \in V(C_1 \cup C_2)$, for otherwise P_2 is an S -jump and (i) holds. From the symmetry we may assume that $y_2 \in V(C_1)$. Thus $y_2 \notin V(C_2)$ by (X1). Assume first that $x \notin V(Z)$. If $y \in V(C_1)$, then $P_1 \cup P_2 \cup P_3 \cup P$ includes a weakly free cross on C_1 , a contradiction. Thus $y \notin V(C_1)$. Since for every S -bridge there is a disk that includes all the attachments of the S -bridge, there exists a disk $C_3 \in \mathcal{C}$ such that either $x_2, y_2, y \in V(C_3)$ or $x_1, y_1, y_2, y \in V(C_3)$. But $y \notin V(C_1)$, and hence $C_3 \neq C_1$. But C_1, C_2 are the only two disks that contain x_2 and the only two disks that contain both x_1 and y_1 . Thus $C_3 = C_2$, contrary to $y_2 \notin V(C_2)$, a contradiction which completes the case $x \notin V(Z)$. We may therefore assume that $x \in V(Z)$, and that P cannot be chosen with $x \notin V(Z)$. If $y \notin V(C_1) \cup V(C_2)$, then P is an S -jump, contrary to the fact that (i) does not hold, and if $y \in V(C_1)$, then $P \cup P_1 \cup P_2 \cup P_3$ includes a weakly

free S -cross, contrary to the fact that (ii) does not hold. Thus $y \in V(C_2) - V(C_1)$. We claim that P_1, P_2, P_3, P is an S -tunnel. To this end let B_i be the S -bridge containing P_i for $i = 1, 2$. The fact that the case $x \notin Z$ and $y \notin V(C_1)$ earlier in this paragraph led to a contradiction implies that all attachments of B_1 and B_2 belong to C_1 . Furthermore, if $B_1 \neq B_2$ and one of them has an attachment in $V(C_1) - V(Z) - \{y_2\}$, then $B_1 \cup B_2$ includes a weakly-free cross, contrary to the fact that (ii) does not hold. Finally, let B be an S -bridge other than B_1 or B_2 with an attachment in the interior of $x_1 Z y_1$. The argument in this paragraph for the case $x \in V(Z)$ shows that every attachment of B belongs to $V(C_2) \cup \{y_2\}$. This proves that P_1, P_2, P_3, P is an S -tunnel, as desired. \square

9. APEX GRAPHS

Let G be a graph. By a *mold* for G we mean a collection $Z = (Z_e : e \in F)$ of (not necessarily disjoint) sets, where $F \subseteq E(G)$ and each Z_e is disjoint from $V(G)$. Given a mold Z for G we define a new graph L as follows. We add the elements of $\bigcup_{e \in F} Z_e$ as new vertices. We subdivide each edge $e \in F$ exactly once, denoting the new vertex by \hat{e} . Finally, for every $e \in F$ and every $z \in Z_e$ we add an edge between z and \hat{e} . We say that L is the graph *determined by G and Z* .

Assume now that there exists a homeomorphic embedding of L into a graph H , assume that G is planar, but that the graph obtained from H by deleting the vertices that correspond to $\bigcup_{e \in F} Z_e$ is not. Can the results obtained thus far be extended to this scenario? We will study this question in this section, and we will find that under some simplifying assumptions the answer is yes. The main technical lemma is (9.8), from which we derive (9.9), the main result of this section. When $|\bigcup_{e \in F} Z_e| = 1$ the main result has a simpler form, stated as (9.10).

Actually, we will not be given a homeomorphic embedding of L into H , but some hybrid between a homeomorphic embedding and a minor containment instead. We now introduce this hybrid. Let us recall that $\eta : G \hookrightarrow S \subseteq H$ means that S is a G -subdivision in H and η maps vertices of G to vertices of S and edges of G to the corresponding paths of S . Let $\eta : G \hookrightarrow S \subseteq H$. Let $Z = (Z_e : e \in F)$ be a mold for G . We say that Z is a *mold*

for G in H if $Z_e \subseteq V(H)$ for every $e \in F$. By abusing notation slightly we will regard Z as a graph with vertex-set $\bigcup_{e \in F} Z_e$ and no edges. Thus we can speak of $(S \cup Z)$ -bridges. By an $S \cup Z$ -link we mean a subgraph B of H such that either B is isomorphic to K_2 and its vertices but not its edge belong to $S \cup Z$, or B consists of a connected subgraph K of $H \setminus V(S \cup Z)$ together with some edges from K to $S \cup Z$ and their ends. Thus every $S \cup Z$ -bridge is an $S \cup Z$ -link, but not the other way around. We say that a mold Z is *feasible* for η if for every $e \in F$ and every $z \in Z_e$ there exists an $S \cup Z$ -link B_{ez} of H such that the following conditions hold for all $e \in F$ and all $z \in Z_e$:

- (i) $Z_e \subseteq V(H) - V(S)$,
- (ii) $z \in V(B_{ez})$,
- (iii) $V(B_{ez}) \cap V(B_{e'z'}) \subseteq V(S \cup Z)$ for all distinct $e, e' \in F$ and all $z \in Z_e$ and $z' \in Z_{e'}$,
- (iv) either some internal vertex of $\eta(e)$ belongs to B_{ez} , or both ends of $\eta(e)$ belong to B_{ez} and some $S \cup Z$ -bridge includes all $B_{ez'}$ for $z' \in Z_e$.

If the mold Z is feasible and the graphs B_{ez} are as above, then we say that the collection $(B_{ez} : e \in F, z \in Z_e)$ of graphs is a *cast* for Z and η in H . Thus feasibility is the promised hybrid between homeomorphic embeddings and minors, as the next lemma explains.

(9.1) *Let G, H be graphs, let $Z = (Z_e : e \in F)$ be a mold for G in H , and let L be the graph determined by G and Z . If $\eta : G \hookrightarrow S \subseteq H \setminus V(Z)$ and Z is feasible for η , then L is isomorphic to a minor of H . Conversely, if $\eta_0 : L \hookrightarrow S_0 \subseteq H$ satisfies $\eta_0(z) = z$ for every $z \in V(Z)$ and η is the restriction of η_0 to G , then Z is feasible for η .*

Proof. Let $\eta : G \hookrightarrow S \subseteq H \setminus V(Z)$ and let Z be feasible for η . Thus there exists a cast $\Gamma = (B_{ez} : e \in F, z \in Z_e)$ for Z and η in H . For $e \in F$ we define a connected graph K_e as follows. If there exists $z \in Z_e$ such that B_{ez} has no attachment in the interior of $\eta(e)$, then let B be the $S \cup Z$ -bridge that includes $B_{ez'}$ for all $z' \in Z_e$ (this exists by the last axiom in the definition of a feasible mold), and let $K_e := B \setminus V(S \cup Z)$. Otherwise let K_e be the union of the interior of $\eta(e)$ and $B_{ez} \setminus V(S \cup Z)$ over all $z \in Z_e$, and all edges from the latter sets to the interior of $\eta(e)$. By contracting all but one edge of each path $\eta(e)$ for every $e \in E(G) - F$ we obtain an L -minor, where each $v \in V(G)$ is represented by $\eta(v)$, each $z \in V(Z)$ is represented by itself, and for $e \in F$ the vertex \hat{e} of L is represented by

K_e . Thus H has an L -minor.

Conversely, if $\eta_0 : L \hookrightarrow S_0 \subseteq H$ satisfies $\eta_0(z) = z$ for every $z \in V(Z)$ and η is the restriction of η_0 to G , then a cast for Z and η in H is constructed by letting $B_{ez} := \eta_0(z\hat{e})$. \square

A cast $(B_{ez} : e \in F, z \in Z_e)$ is *united* if there exist distinct edges $e, e' \in F$ and (not necessarily distinct) vertices $z \in Z_e$ and $z' \in Z_{e'}$ such that B_{ez} and $B_{e'z'}$ are subgraphs of the same $S \cup Z$ -bridge. A cast $(B_{ez} : e \in F, z \in Z_e)$ is *full* if each B_{ez} is an $S \cup Z$ -bridge.

(9.2) *Let G, S, H be graphs, let $\eta : G \hookrightarrow S \subseteq H$, and let $Z = (Z_e : e \in F)$ be a feasible mold for G in H . Then there exists a cast for Z and η in H that is either united or full.*

Proof. Let $\Gamma = (B_{ez} : e \in F, z \in Z_e)$ be a cast for Z and η in H . If Γ is not united, then we may replace each B_{ez} by the $S \cup Z$ -bridge it is contained in, thereby producing a full cast. \square

Let G, S, H, η and Z be as above. As in earlier sections of the paper we will produce S -jumps and S -crosses. However, in order for them to be useful we need them to behave well with respect to a cast. That leads to the following definitions. An S -path P is *compatible* with a full cast $(B_{ez} : e \in F, z \in Z_e)$ if P is disjoint from Z and it is the case that if P is a subgraph of B_{ez} for some $e \in F$ and $z \in Z_e$, then either one of the ends of P belongs to the interior of $\eta(e)$, or B_{ez} has no attachment in the interior of $\eta(e)$ (in which case both ends of $\eta(e)$ are attachments of B_{ez} by the last axiom in the definition of cast) and one end of P is an end of $\eta(e)$. We say that a cross P_1, P_2 is *compatible* with a full cast $(B_{ez} : e \in F, z \in Z_e)$ if both S -paths P_1, P_2 are compatible with the cast, and if P_1, P_2 are subgraphs of the same S -bridge B , then $B = B_{ez}$ for no $e \in F$ and $z \in Z_e$.

(9.3) *Let G, H be graphs, let $\eta : G \hookrightarrow S \subseteq H'$ be a homeomorphic embedding, let $Z = (Z_e : e \in F)$ be a feasible mold for G in H , let $(B_{ez} : e \in F, z \in Z_e)$ be a full cast for Z and η in H , and let P be an S -path in $H \setminus Z$. Let F' be the set of all edges $e \in F$ such that if P is a subgraph of B_{ez} for some $z \in Z_e$, then either one end of P is an internal vertex of the path $\eta(e)$, or B_{ez} has no attachment in the interior of $\eta(e)$ and one end of P is an end of $\eta(e)$. Then P is compatible with the cast $(B_{ez} : e \in F', z \in Z_e)$.*

The proof is clear.

The following lemma shows how to use S -jumps compatible with a full cast.

(9.4) *Let G be an internally 4-connected planar graph, let H be a graph, let $Z = (Z_e : e \in F)$ be a mold for G in H , and let L be the graph determined by G and Z . If $\eta : G \hookrightarrow S \subseteq H \setminus V(Z)$ is a homeomorphic embedding, Γ is a full cast for Z and η in H and there exists an S -jump compatible with Γ , then there exist vertices $u, v \in V(G)$ that are not cofacial in G such that $L + uv$ is isomorphic to a minor of H .*

Proof. Let $\Gamma = (B_{ez} : e \in F, z \in Z_e)$ and let P be an S -jump compatible with Γ . The proof of (9.1) and the definition of compatible path imply that there exists a graph L' obtained from L by subdividing at most two edges of $E(G) - F$ such that $L' + xy$ is isomorphic to a minor of H for some two vertices $x, y \in V(L')$ that are not cofacial in $L' \setminus V(Z)$. This is straightforward, except for the case when P is a subgraph of B_{ez} for some $e \in F$ and $z \in Z_e$, and B_{ez} has no attachment in the interior of $\eta(e)$. Then one end of P , say $\eta(x)$, is an end of $\eta(e)$ by the definition of compatible path. If the other end of P is $\eta(y)$ for some $y \in V(G)$, then $L + y\hat{e}$ is isomorphic to a minor of H , and y and \hat{e} are not cofacial in $L \setminus V(Z)$, because y and x are not. If the other end of P belongs to the interior of $\eta(f)$ for some $f \in E(G)$, then similarly $L' + w\hat{e}$ is isomorphic to a minor of H , where L' is obtained from L by subdividing the edge f and w denotes the new vertex, and w and \hat{e} are not cofacial in $L' \setminus V(Z)$. This completes the argument that $L' + xy$ is isomorphic to a minor of H .

The conclusion now follows from (6.4) and (6.5) in the same way as (6.6). □

Next we shows how to use S -crosses compatible with a full cast.

(9.5) *Let G be an internally 4-connected triangle-free planar graph, let H be a graph, let $Z = (Z_e : e \in F)$ be a mold for G in H , and let L be the graph determined by G and Z . If $\eta : G \hookrightarrow S \subseteq H \setminus V(Z)$ is a homeomorphic embedding, Γ is a full cast for Z and η in H and there exists a free S -cross compatible with Γ , then either*

- (i) *there exist vertices $u, v \in V(G)$ that are not cofacial in G such that $L + uv$ is isomorphic to a minor of H , or*

(ii) there exists a facial cycle C of G and distinct vertices $u_1, u_2, v_1, v_2 \in V(C)$ appearing on C in the order listed such that $L + u_1v_1 + u_2v_2$ is isomorphic to a minor of H .

Proof. The proof is similar to the proof of (9.4). Let us omit the tedious details, but instead clarify one subtlety. Let C be a disk in S , and let P be an H -path compatible with Γ with ends $x, y \in V(C)$, where x, y do not belong to the same segment of S . Let $\Gamma = (B_{ez} : e \in F, z \in Z_e)$, and let P be a subgraph of B_{ez} for some $e \in F$ and $z \in Z_e$. If $\eta(e)$ is not a subgraph of C , then the definition of compatible path implies that B_{ez} has no attachment in the interior of $\eta(e)$ and one of x, y is an end of $\eta(e)$, say y is its end. If x is a branch vertex of S , then let $u \in V(G)$ be such that $\eta(u) = x$; otherwise let $u \in V(G)$ be such that $\eta(u) \neq y$ is an end of the segment of S containing x . Since x, y do not belong to the same segment, (3.3) implies that u and \hat{e} are not cofacial in $L \setminus V(Z)$. The presence of P guarantees that $L + u\hat{e}$ is isomorphic to a minor of H , and so (i) holds.

Let P_1, P_2 be a free S -cross compatible with Γ . The previous paragraph shows that we may assume that if P_i is a subgraph of B_{ez} for some $e \in F$ and $z \in Z_e$, then $\eta(e)$ is a subgraph of C . The rest of the proof is similar to (9.4), using the argument of (6.6). \square

Let $\eta : G \hookrightarrow S \subseteq H$, and let $\eta' : G \hookrightarrow S' \subseteq H$ be obtained from η by a rerouting. If H' is a subgraph of H and both S and S' are subgraphs of H' , then we say that the rerouting is *within* H' .

Our next objective is to give a sufficient condition for a rerouting to preserve feasibility. To that end we need to discuss the effect of reroutings on casts. Let G, H be graphs, let $\eta : G \hookrightarrow S \subseteq H$ be a homeomorphic embedding, let $F \subseteq E(G)$, and let $\eta' : G \hookrightarrow S' \subseteq H$ be obtained from η by a rerouting. We say that the rerouting is *F-safe* if the following conditions are satisfied:

- (i) if the rerouting replaces a subpath of S by an S -path Q and Q is a subgraph of an S -bridge B , and $e \in F$ is such that either B has an attachment in the interior of $\eta(e)$, or both ends of $\eta(e)$ are attachments of B , then the rerouting is an I-rerouting based at $\eta(e)$,
- (ii) if the rerouting is a T-rerouting centered at $\eta(v) \in V(S)$, then no edge of G incident with v belongs to F , and

(iii) if the rerouting is a V- or X-rerouting based at $\eta(e_1)$ and $\eta(e_2)$, then $e_1, e_2 \notin F$.

Thus every proper I-rerouting is F -safe.

(9.6) *Let G, H be graphs, let $\eta : G \hookrightarrow S \subseteq H$ be a homeomorphic embedding, let Z be a mold for G in H that is feasible for η , let there be a full cast for Z and η in H , and let $\eta' : G \hookrightarrow S' \subseteq H$ be obtained from η by an F -safe rerouting within $H \setminus V(Z)$. Then Z is feasible for η' .*

Proof. Let $Z = (Z_e : e \in F)$ and let $\Gamma = (B_{ez} : e \in F, z \in Z_e)$ be a full cast for Z and η in H . Let $e \in F$ and $z \in Z_e$. We wish to define an $S' \cup Z$ -link B'_{ez} . If the rerouting is an I-rerouting, then let W be a segment of S such that the rerouting is based at W ; otherwise let W be the null graph. The construction will be such that $V(B'_{ez}) \subseteq V(B_{ez} \cup W)$. That will guarantee that the links thus defined will satisfy the third axiom in the definition of feasibility. The other axioms will be clear from the construction.

Assume first that B_{ez} includes an S -path Q that replaced a subpath P of S during the rerouting. Since Γ is a cast, the S -bridge B_{ez} either has an attachment in the interior of $\eta(e)$, or both ends of $\eta(e)$ are attachments of B_{ez} . The first axiom in the definition of F -safety implies that the rerouting is an I-rerouting based at $\eta(e)$. The $S \cup Z$ -bridge B_{ez} includes a path from z to the interior of Q ; let B'_{ez} be such a path with no internal vertex in $S' \cup Z$. This completes the construction when B'_{ez} includes an S -path that replaced a subpath P of S during the rerouting.

Thus we may assume that B_{ez} includes no such S -path. If no attachment of B_{ez} belongs to $\eta(e)$ and to the interior of a subpath of S that got replaced by an S -path during the rerouting, then we let $B'_{ez} := B_{ez}$. We may therefore assume that an attachment x of B_{ez} belongs to $\eta(e)$ and to the interior of a subpath P of S that got replaced by an S -path Q during the rerouting. The second and third axiom in the definition of safety imply that the rerouting is an I-rerouting and that x belongs to the interior of $\eta(e)$. Thus the I-rerouting is based at $\eta(e)$. We define $B'_{ez} := P \cup B_{ez}$. If the ends of Q are not equal to the ends of $\eta(e)$, then the $S' \cup Z$ -link B'_{ez} has an attachment in the interior of $\eta'(e)$. Thus we may assume that Q and $\eta(e)$ have the same ends. But then for every $z' \in Z_e$ the $S \cup Z$ -link $B_{ez'}$ has an attachment in the interior of $P = \eta(e)$, and hence $B'_{ez'}$ is a

subgraph of the $S' \cup Z$ -bridge containing $\eta(e)$. Hence the $S' \cup Z$ -links B'_{ez} satisfy the last feasibility axiom. The other axioms are clear.

Thus $(B'_{ez} : e \in F, z \in Z_e)$ is a cast for Z and η' in H , as required. \square

(9.7) *Let G, H be graphs, let $Z = (Z_e : e \in F)$ be a mold for G in H , and let $\eta : G \hookrightarrow S \subseteq H$ be a homeomorphic embedding. If Z is feasible for η and there exists a full cast for Z and η in H , and $\eta' : G \hookrightarrow S' \subseteq H$ is obtained from η by a proper I-rerouting within $H \setminus V(Z)$, then Z is feasible for η' .*

Proof. This follows immediately from (9.6), because a proper I-rerouting is F -safe. \square

The following is the main technical lemma of this section.

(9.8) *Let G be an internally 4-connected planar graph not isomorphic to the cube, let H be a graph, and let $Z = (Z_e : e \in F)$ be a mold for G in H . Let $H' := H \setminus \bigcup_{e \in F} Z_e$, and let $\eta_0 : G \hookrightarrow S_0 \subseteq H'$ be a homeomorphic embedding such that the mold Z is feasible for η_0 . Then there exist a homeomorphic embedding $\eta : G \hookrightarrow S \subseteq H'$ obtained from η_0 by repeated reroutings within H' and a set $F' \subseteq F$ such that every two edges in $F - F'$ are cofacial in G and letting Z' denote the mold $(Z_e : e \in F')$ one of the following conditions holds:*

- (i) *there is a united cast for Z' and η , or*
- (ii) *there exists an S -jump compatible with some full cast for Z' and η , or*
- (iii) *there exists a free S -cross compatible with some full cast for Z' and η , or*
- (iv) *H' has an S -separation, or*
- (v) *H' is planar.*

Proof. Let $\eta_0 : G \hookrightarrow S_0 \subseteq H'$ and Z be as stated. We may assume that (i) does not hold. We start with the following claim.

- (1) *Let $\eta : G \hookrightarrow S \subseteq H'$ be obtained from η_0 by repeated reroutings within H' , let $F' \subseteq F$ be such that every two edges in $F - F'$ are cofacial in G , let $Z' := (Z_e :$*

$e \in F'$), and let there exist a full cast for Z' and η in H' . If η' is obtained from η by an F' -safe rerouting, then there is a full cast for Z' and η' in H' .

To prove (1) we first notice that (9.6) implies that Z' is feasible for η' in H' . By (9.2) there is a cast for Z' and η' in H' that is united or full. The former does not hold by our assumption that (i) does not hold, and hence the latter holds. This proves (1).

By (3.1) applied to the graphs G, S_0 and H' there exists a homeomorphic embedding $\eta : G \hookrightarrow S \subseteq H'$ obtained from η_0 by repeated proper I-reroutings such that every unstable S -bridge is 2-separated from S . Since every proper I-rerouting is F -safe, it follows from (1) that there is a full cast for Z and η in H' . Let $\Gamma := (B_{ez} : e \in F, z \in Z_e)$ be such a full cast.

(2) *If there exists an S -jump in H' , then the theorem holds.*

To prove (2) let P be an S -jump. If the S -bridge containing P is equal to B_{ez} for some $e \in F$ and $z \in Z_e$, then there is exactly one such edge e , and we define $F' := F - \{e\}$; otherwise we let $F' := F$. Then P is compatible with $(Z_e : e \in F')$, and hence F' and η satisfy (ii). This proves (2).

(3) *Let u be a vertex of G of degree three, let F' be obtained from F by removing all edges incident with u , let $\eta' : G \hookrightarrow S' \subseteq H'$ be obtained from η by a sequence of F' -safe reroutings, and let there exist a local S' -triad centered at $\eta'(u)$. Then F' satisfies the conclusion of the theorem.*

To prove (3) we first deduce from (1) that there exists a full cast $\Gamma' = (B'_{ez}; e \in F', z \in Z_e)$ for Z' and η' . Let Z_1, Z_2, Z_3 be the three segments of S' incident with $v := \eta'(u)$, let v_i be the other end of Z_i , and let the local S' -triad be Q_1, Q_2, Q_3 , where Q_i has ends x and $x_i \in V(Z_i)$. Let $L_i := v_i Z_i x_i$ and $P_i := v Z_i x_i$. We may assume that η' and the triad Q_1, Q_2, Q_3 are chosen so that $|V(L_1)| + |V(L_2)| + |V(L_3)|$ is minimum.

Let $X_1 = V(P_1 \cup P_2 \cup P_3 \cup Q_1 \cup Q_2 \cup Q_3)$ and $Y_1 = V(S) - (X_1 - \{x_1, x_2, x_3\})$. If $H' \setminus \{x_1, x_2, x_3\}$ has no path between X_1 and Y_1 , then H' has a separation (X, Y) such that $X \cap Y = \{x_1, x_2, x_3\}$, $X_1 \subseteq X$, and $Y_1 \subseteq Y$. Then (X, Y) satisfies outcome (iv) of the theorem.

We may therefore assume that there exists a path P in H' as above. Let the ends of P be $x \in X_1 - \{x_1, x_2, x_3\}$ and $y \in Y_1 - \{x_1, x_2, x_3\}$. We may assume that P has no internal vertex in $X_1 \cup Y_1$. If P is a subgraph of the S' -bridge B'_{ez} for some $e \in F'$ and $z \in Z_e$, then we may assume that y satisfies the following specifications. If B'_{ez} has an attachment in the interior of $\eta'(e)$, then we may assume that y belongs to the interior of $\eta'(e)$; otherwise we may assume that y is an end of $\eta'(e)$ (because both ends of $\eta'(e)$ are attachments of B'_{ez} by the last axiom in the definition of cast, and at least one end of $\eta'(e)$ does not belong to $Z_1 \cup Z_2 \cup Z_3$, because G is internally 4-connected).

Assume first that $x \in V(Q_1 \cup Q_2 \cup Q_3)$. Then $y \notin V(L_1 \cup L_2 \cup L_3)$ by the choice of Q_1, Q_2, Q_3 . Since G is not isomorphic to a cube we deduce from (8.1) that there is an S' -jump with one end y . The choice of y implies that the S' -jump is compatible with Γ' , and hence outcome (ii) holds. This completes the case that $x \in V(Q_1 \cup Q_2 \cup Q_3)$. Furthermore, it implies that we may assume that the S' -bridge containing $Q_1 \cup Q_2 \cup Q_3$ has all attachments in $Z_1 \cup Z_2 \cup Z_3$.

Thus $x \in V(P_1 \cup P_2 \cup P_3)$. Let B be the S' -bridge containing P . If B has an attachment outside $Z_1 \cup Z_2 \cup Z_3$, then P may be replaced by a path with an end not in $Z_1 \cup Z_2 \cup Z_3$; otherwise replacing a path of $P_1 \cup P_2 \cup P_3$ by P is a T-rerouting centered at v , and it is F' -safe. The resulting homeomorphic embedding has a triad that contradicts the choice of η' and Q_1, Q_2, Q_3 . Thus we may assume that $y \notin V(Z_1 \cup Z_2 \cup Z_3)$.

We may assume that P is not an S' -jump, for otherwise (ii) holds, because P is compatible with Γ' by the choice of y . Thus there exists a disk C in S' such that $x, y \in V(C)$. It follows that C includes two of the segments incident with v , say Z_1 and Z_2 . Now both S' -paths $Q_1 \cup Q_2$ and P are compatible with Γ , the former because for $e \in F'$ the S' -bridge B'_{ez} does not include $Q_1 \cup Q_2$, which in turn follows from the fact that the S' -bridge containing $Q_1 \cup Q_2 \cup Q_3$ has all attachments in $Z_1 \cup Z_2 \cup Z_3$. Thus $Q_1 \cup Q_2, P$ is an S' -cross compatible with Γ , and it is free by the internal 4-connectivity of G . This proves (3).

(4) *If there exists an S -triad in H' , then the theorem holds.*

To prove (4) assume that there exists an S -triad in H' . The triad is local by (5.2), and hence

the claim follows from (3) applied to the homeomorphic embedding η . This proves (4).

(5) *If there exists a weakly free S -cross in H' , then the theorem holds.*

To prove (5) let P_1, P_2 be a weakly-free S -cross in H' on a disk C , and assume for a moment that the cross is free. Let F' be obtained from F by removing the at most two edges $e \in F$ such that P_1 or P_2 is a subgraph of B_{ez} for some $z \in Z_e$. If each removed edge belongs to $E(C)$, then F' satisfies outcome (iii) of the present theorem, and so we may assume that say P_1 is a subgraph of B_{ez} for some $e \in F - E(C)$ and $z \in Z_e$. But then the bridge B_{ez} includes an S -jump or an S -triad in H' , and hence the theorem holds by (2) and (4). This concludes the case when P_1, P_2 is a free cross, and so we may assume that it is not.

Thus there exist segments Z_1, Z_2 in S with common end v and the other ends v_1, v_2 , respectively, such that both are subgraphs of C and such that the ends of P_i can be labeled x_i, y_i in such a way that $v_1, x_1, x_2, v, y_1, y_2, v_2$ occur on $Z_1 \cup Z_2$ in the order listed. There are two cases depending on the degree of v . Assume first that the degree of v is three. If the S -bridge of H' that includes the path P_1 has no attachment outside of the three segments incident with v , then replacing $x_1 Z_1 v$ by P_1 is a T-rerouting that is F' -safe, where F' is as in (3), and $P_2, x_1 P_1 x_2, x_2 P_1 v$ is a local triad. Thus in this case the theorem holds by (3). We may therefore assume that there exists a path P with ends $x \in V(P_1) - \{x_1, y_1\}$ and $y \in V(S)$ such that P has no internal vertex in $S \cup P_1 \cup P_2$ and y does not belong to any of the segments of S incident with v . If $y \in V(C)$, then there exists a free S -cross, a case we already handled, and so we may assume not. For $i = 1, 2$ let C_i be the disk other than C that includes Z_i . Since y_1 belongs to the interior of Z_2 , the only two disks it belongs to are C and C_2 . We may assume that $y \in V(C_2)$, for otherwise $P_1 \cup P$ includes an S -jump (with ends y_1 and y), in which case the theorem holds by (2). But $C_1 \cap C_2$ is equal to the third segment incident with v , and hence $x_1 \notin V(C_1 \cap C_2)$, and therefore $x_1 \notin V(C_2)$. Thus $P_1 \cup P$ includes an S -jump with ends x_1 and y , and so the theorem holds by (2). This completes the case when v has degree three.

We may therefore assume that v has degree at least four. Let us define the *height* of the cross P_1, P_2 to be $|E(L_1)| + |E(L_2)|$, where $L_1 := v_1 Z_1 x_1$ and $L_2 := v_2 Z_2 y_2$. We

proceed similarly as in the proof of (4.4), but with extra care. Let F' be obtained from F by deleting all edges $e \in F$ such that $\eta(e)$ is a subgraph of C , and let $Z' := (Z_e : e \in F')$. We claim that F' satisfies the conclusion of the theorem. Let $e_1, e_2 \in E(G)$ be such that $\eta(e_i) = Z_i$. Let $\eta_1 : G \hookrightarrow S_1 \subseteq H'$ be a homeomorphic embedding obtained from η by a sequence of proper V -reroutings based at e_1, e_2 and let Q_1, Q_2 be a weakly free S_1 -cross based at $\eta_1(e_1), \eta_1(e_2)$ such that among all such triples (η_1, Q_1, Q_2) this one minimizes the height of the cross Q_1, Q_2 . Since every proper V -rerouting is F' -safe, it follows from (1) that there is full cast for Z' and η_1 in H' . In order to prevent the introduction of unnecessary notation we now make the assumption that $\eta = \eta_1$, $P_1 = Q_1$ and $P_2 = Q_2$. This can be done with the proviso that for the remainder of the proof of (5) Γ is a full cast for Z' and η (as opposed to a full cast for Z).

Let X' be the vertex-set of $P_1 \cup P_2 \cup vZ_1x_1 \cup vZ_2y_2$ and let $Y' = V(S) - (X' - \{v, x_1, y_2\})$. If there is no path in $H' \setminus \{v, x_1, y_2\}$ with one end in X' and the other in Y' , then there exists a separation (X, Y) of order three with $X' \subseteq X$ and $Y' \subseteq Y$. This separation satisfies (iv), as required, and so we may assume that there exists a path P in $H' \setminus \{v, x_1, y_2\}$ with one end $x \in X'$ and the other end $y \in Y'$.

We first complete the proof of (5) assuming that $y \notin V(Z_1 \cup Z_2)$, that at least one of x, y is not in $V(C_1)$, and that at least one of x, y is not in $V(C_2)$. From the symmetry we may assume that $x \in V(P_1 \cup y_2Z_2v)$. If P_1 is a subgraph of B_{ez} for some $e \in F'$ and $z \in Z_e$, then we may assume that either y belongs to the interior of $\eta(e)$, or y is an end of $\eta(e)$ and $y \notin V(C)$. (This is indeed possible—by the choice of F' at least one end of $\eta(e)$ does not belong to $Z_1 \cup Z_2$.) If $y \notin V(C \cup C_2)$, then $P_1 \cup P$ includes an S -jump with ends y_1 and y , which is compatible with Γ . Thus (ii) holds. Next let us assume that $y \in V(C_2)$. Then $y \notin V(C)$, because $C \cap C_2 = Z_2$. Since v has degree at least four, (X2) implies that $V(C_1) \cap V(C_2) = \{v\}$. It follows that $y \notin V(C_1)$, and so $P_1 \cup P$ includes an S -jump, which is compatible with Γ . Thus, again, (ii) holds. We may therefore assume that $y \in V(C)$. That implies that P_1 is a subgraph of B_{ez} for no $e \in F'$ and $z \in Z_e$, and so from the symmetry we may assume the same about P_2 . Since $y \in V(C)$ we deduce that $P_1 \cup P_2 \cup P$ includes a free cross. Since P_1, P_2 are not subgraphs of any B_{ez} for $e \in F'$ it follows that the cross is compatible with Γ , and so (iii) holds. This completes the case

that $y \notin V(Z_1 \cup Z_2)$, at least one of x, y is not in $V(C_1)$, and at least one of x, y is not in $V(C_2)$. Thus we may assume that the S -bridge that contains P_1 has all its attachments in $Z_1 \cup Z_2$, and from the symmetry we may assume the same about the S -bridge containing P_2 . In particular, the X -rerouting of Z_1, Z_2 that makes use of P_1, P_2 is proper, and hence is F' -safe.

Next we handle the case that $y \in V(Z_1 \cup Z_2)$. Again, from the symmetry we may assume that $x \in V(P_1 \cup y_2 Z_2 v)$. If $y \in V(L_1)$, then replacing P_1 by P if $x \notin V(P_1)$ and by $P \cup x P_1 y_1$ otherwise produces a cross of smaller height, contrary to the choice of the triple (η_1, Q_1, Q_2) . If $y \in V(L_2)$, then replacing $y Z_2 x$ by P if $x \notin V(P_1)$ and replacing $y Z_2 y_1$ by $P \cup x P_1 y_1$ results in a homeomorphic embedding η' obtained from η by a proper V -rerouting, and P_1, P_2 can be modified to give a cross P'_1, P'_2 such that the triple (η', P'_1, P'_2) contradicts the choice of (η_1, Q_1, Q_2) . Thus $y \notin V(Z_1 \cup Z_2)$.

Finally, from the symmetry we may assume that $x, y \in V(C_2)$. Let B be the S -bridge containing P . Since we may assume that x, y cannot be chosen to satisfy any of the cases already handled, it follows that every attachment of B belongs to C_2 . Thus we may assume that if $B = B_{ez}$ for some $e \in F'$ and $z \in Z_e$, then either y is an internal vertex of $\eta(e)$, or B has no attachment in the interior of $\eta(e)$ and y is an end of $\eta(e)$. Since $V(C_1) \cap V(C_2) = \{v\}$, it follows that $y \notin V(C_1)$. Now let $\eta' : G \hookrightarrow S' \subseteq H'$ be obtained from η by the X -rerouting using the cross P_1, P_2 , and let Z'_1, Z'_2 be the segments of S' corresponding to Z_1, Z_2 , respectively. Thus $Z'_1 = v_1 Z_1 x_1 \cup P_1 \cup y_1 Z_2 v$ and $Z'_2 = v_2 Z_2 y_2 \cup P_2 \cup x_2 Z_1 v$. As pointed out earlier, this rerouting is F' -safe. It follows that Γ is a cast for Z' and η' . Now P is an S' -jump, and is compatible with Γ by the choice of y . Thus (ii) holds. This completes the proof of (5).

Since every unstable S -bridge is 2-separated from S we may apply (8.2) to the graphs G, S and H' to deduce that one of the outcomes (i)–(vi) of that lemma holds. But we may assume that (i) does not hold by (2), we may assume that (ii) does not hold by (5), we may assume that (iii) does not hold, because otherwise outcome (iv) of the present theorem holds, we may assume that (8.2)(iv) does not hold by (4), and we may assume that (8.2)(vi) does not hold, for otherwise outcome (v) holds. Thus we may assume that (8.2)(v) holds.

Let the notation be as in the definition of S -tunnel as introduced prior to (8.2), and

let $e_0 \in E(G)$ be such that $\eta(e_0) = W$. Let D and D' be cycles in G such that $\eta(D) = C$ and $\eta(D') = C'$. If one of B_1, B_2 is equal to B_{ez} for some $e \in E(D) - \{e_0\}$, then such an e is unique (because if $B_1 \neq B_2$, then both have the same unique attachment outside W), and we denote it by e_1 ; otherwise e_1 is undefined. If e_1 is well-defined we define $F' := F - \{e_0, e_1\}$; otherwise we define $F' := F - \{e_0\}$. Let $Z' := (Z_e : e \in F')$. Let $\eta' : G \hookrightarrow S' \subseteq H'$ be obtained from η by replacing $x_1 W y_1$ by P_1 . Then this rerouting is F' -safe, and so Z' is feasible for η' by (9.6). By (9.2) we may assume that there is a full cast for Z' and η' in H' , for otherwise the theorem holds. Let $\Gamma' = (B'_{ez} : e \in F', z \in Z_e)$ be such a cast as constructed in the proofs of (9.6) and (9.2). Let B' be the S' -bridge containing P_2 and P_4 . Then B' is a subgraph of the union of $B_1, B_2, x_1 W y_1$ and all S -bridges that have an attachment in the interior of $x_1 W y_1$. It follows from the definition of S -tunnel by analyzing the proof of (9.6) that if $B' = B'_{ez}$ for some $e \in F'$ and $z \in Z_e$, then $e \in E(D') - \{e_0\}$. The S' -bridge B' includes an S' -path P with one end say $x \in V(C) - V(W)$ and the other end say $y \in V(C') - V(W)$. We may assume that $x \in V(\eta(e_1))$ if e_1 is well-defined. In a manner similar as before, by replacing y by a different vertex if necessary, we may choose P to be compatible with Γ' . If P is an S' -jump, then outcome (ii) holds, and so we may assume that it is not. Thus some disk C'' of S' includes both x and y . It follows that B' includes an S' -triad. By (5.2) the triad is local; let it be centered at $v \in V(S)$. It follows that $\eta(e_0)$ is incident with v , and so is $\eta(e_1)$ if e_1 is well-defined (by the choice of x). Thus the theorem holds by (3). \square

We deduce the following corollary.

(9.9) *Let G be an internally 4-connected triangle-free planar graph not isomorphic to the cube, and let $F \subseteq E(G)$ be such that no two elements of F belong to the same facial cycle of G . Let H be a graph, and let $Z = (Z_e : e \in F)$ be a mold for G in H . Let $H' := H \setminus \bigcup_{e \in F} Z_e$, and let $\eta_0 : G \hookrightarrow S_0 \subseteq H'$ be a homeomorphic embedding such that the mold Z is feasible for η_0 . If H' is internally 4-connected and non-planar, then there exists a set $F' \subseteq F$ with $|F - F'| \leq 1$ such that the graph L determined by G and $(Z_e : e \in F')$ satisfies one of the following conditions:*

- (i) there exist vertices $u, v \in V(L) - V(Z)$ that do not belong to the same facial cycle of $L \setminus V(Z)$ such that $L + uv$ is isomorphic to a minor of H ,
- (ii) there exists a facial cycle C of $L \setminus V(Z)$ and distinct vertices $u_1, u_2, v_1, v_2 \in V(C)$ appearing on C in the order listed such that $L + u_1v_1 + u_2v_2$ is isomorphic to a minor of H .

Proof. Let $\eta : G \hookrightarrow S \subseteq H', F'$ and $Z' = (Z_e : e \in F')$ be as in (9.8). Then $|F - F'| \leq 1$, because no two edges of F are cofacial. By (9.8) one of (i)–(v) of that theorem holds. But (iv) does not hold, because H' is internally 4-connected, and (v) does not hold, because H' is not planar. Let $\Gamma = (B_{ez} : e \in F', z \in Z_e)$ be a cast satisfying (i), (ii), or (iii) of (9.8). If (9.8)(i) holds, then let $e, f \in F$ be distinct edges such that B_{ez} and B_{fw} are subgraphs of the same $S \cup Z$ -bridge for some $z \in Z_e$ and $w \in Z_f$. It follows from the proof of (9.1) that $L + \hat{e}\hat{f}$ is isomorphic to a minor of H , as required for (i). If (9.8)(ii) holds, then (i) holds by (9.4), and if (9.8)(iii) holds, then (ii) holds by (9.5). \square

When $V(Z)$ has size one we get the following explicit version, which is used in [3].

(9.10) *Let G be an internally 4-connected triangle-free planar graph not isomorphic to the cube, and let $F \subseteq E(G)$ be a non-empty set such that no two edges of F are incident with the same face of G . Let G' be obtained from G by subdividing each edge in F exactly once, and let L be the graph obtained from G' by adding a new vertex $v \notin V(G')$ and joining it by an edge to all the new vertices of G' . Let a subdivision of L be isomorphic to a subgraph of H , and let $u \in V(H)$ correspond to the vertex v . If $H \setminus u$ is internally 4-connected and non-planar, then there exists an edge $e \in E(L)$ incident with v such that either*

- (i) there exist vertices $x, y \in V(G')$ not belonging to the same face of G' such that $(L \setminus e) + xy$ is isomorphic to a minor of H , or
- (ii) there exist vertices $x_1, x_2, x_3, x_4 \in V(G')$ appearing on some face of G' in order such that $(L \setminus e) + x_1x_3 + x_2x_4$ is isomorphic to a minor of H .

Proof. For $e \in F$ let $Z_e := \{u\}$. Then L is the graph determined by G and Z . Since a subdivision of L is isomorphic to a subgraph of H , the second half of (9.1) implies that

Z is feasible for a homeomorphic embedding $\eta : G \hookrightarrow S \subseteq H \setminus u$. By (9.9) the corollary holds. \square

Acknowledgement

We thank Alexander Kelmans for carefully reading an earlier version of this manuscript, providing extensive comments and pointing out several errors. We would like to acknowledge that this paper grew out of ideas developed in the work of the second author with Neil Robertson and Paul Seymour.

References

1. G. Ding, B. Oporowski, R. Thomas and D. Vertigan, Large non-planar graphs and an application to crossing-critical graphs, *J. Combin. Theory Ser. B* **101** (2011), 111–121.
2. R. Hegde and R. Thomas, Non-embeddable extensions of embedded minors, manuscript.
3. K. Kawarabayashi, S. Norine, R. Thomas and P. Wollan, K_6 minors in 6-connected graphs of bounded tree-width, [arXiv:1203.2171](https://arxiv.org/abs/1203.2171).
4. A. K. Kelmans, The concept of a vertex in a matroid, the non-separating cycles and a new criterion for graph planarity, In *Algebraic Methods in Graph Theory*, **1**, Colloq. Math. Soc. János Bolyai (Szeged, Hungary), North-Holland **25** (1978), 345–388.
5. A. K. Kelmans, Finding special subdivisions of K_4 in a graph, In *Finite and infinite sets*, Colloq. Math. Soc. János Bolyai (Eger, Hungary), North-Holland **37** (1981), 487–508.
6. A. K. Kelmans, On 3-connected graphs without essential 3-cuts or triangles, *Soviet Math. Dokl.* **33** (1986), 698–703.
7. A. K. Kelmans, Graph planarity and related topics, *Contemporary Math.* **147** (1993), 635–667.
8. N. Robertson and P. D. Seymour, Graph Minors IX. Disjoint crossed paths, *J. Combin. Theory Ser. B* **49** (1990), 40–77.
9. N. Robertson, P. D. Seymour and R. Thomas, Excluded minors in cubic graphs, manuscript.

10. P. D. Seymour, Disjoint paths in graphs, *Discrete Math.* **29** (1980), 293-309.
11. Y. Shiloach, A polynomial solution to the undirected two paths problem, *J. Assoc. Comp. Machinery*, **27**, (1980), 445-456.
12. R. Thomas and J. Thomson, Excluding minors in nonplanar graphs of girth five, *Combinatorics, Probability and Computing* **9** (2000), 573-585.
13. C. Thomassen, 2-linked graphs, *Europ. J. Combinatorics* **1** (1980), 371-378.
14. W. T. Tutte, How to draw a graph, *Proc. London Math. Soc.* **13** (1963), 743-768.
15. W. T. Tutte, On the algebraic theory of graph colorings, *J. Combin. Theory Ser. B* **1** (1966), 15-50.
16. H. Whitney, Congruent graphs and the connectivity of graphs, *Amer. J. Math.* **54** (1932), 150-168.

This material is based upon work supported by the National Science Foundation. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.