

October 27, 2015



Math 3012 - Applied Combinatorics Lecture 18


William T. Trotter
trotter@math.gatech.edu

Reminder

Test 3 Tuesday, November 24, 2015. Details on material for which you will be responsible were sent by email after class the preceding Thursday. Again, I ask all of you to study hard. Experience shows that the closing portion of this course has most content. The concepts and techniques will have lasting value.


Inclusion/Exclusion - Prelude

Question In the "Venn Diagram" shown below, the universe X contains 23 elements. There are 8 in the set A and 11 in B . If there are 5 in $A \cap B$, then how many elements of X belong to neither A nor B ?



Inclusion/Exclusion - Prelude (2)

Question In the "Venn Diagram" shown below, the universe X contains 2307 elements. We want to determine the number of elements of X that don't belong to any of A , B and C . If we know the number of elements in the following sets, can we do this? A , B , C , $A \cap B$, $B \cap C$, $A \cap C$, $A \cap B \cap C$.



Inclusion/Exclusion (1)

Notation Let X be a set of objects and suppose that for every element i in $\{1, 2, \dots, n\}$, we have a property P_i so that for all x in X , the statement “ x satisfies property P_i ” is either true or false ... but never ambiguous. Then for a subset S of $\{1, 2, \dots, n\}$, let $N(S)$ be the subset of X consisting of all x in X which satisfy property P_i for all i in S . Note that $N(\emptyset) = X$.

Notation Let N_0 be the subset of X consisting of those objects which satisfy **none** of the properties.

Inclusion/Exclusion (2)

Theorem Let X be a set of objects and let P_i be a property for X for each $i = 1, 2, \dots, n$.
Then:

$$N_0 = \sum_{S \subseteq \{1, 2, \dots, n\}} (-1)^{|S|} N(S)$$

Example When $n = 2$,

$$N_0 = N(\emptyset) - N(1) - N(2) + N(1).$$

Inclusion/Exclusion (3)

Theorem Let X be a set of objects and let P_i be a property for X for each $i = 1, 2, \dots, n$.
Then:

$$N_0 = \sum_{S \subseteq \{1, 2, \dots, n\}} (-1)^{|S|} N(S)$$

Example When $n = 3$,

$$\begin{aligned} N_0 &= N(\emptyset) \\ &\quad - N(1) - N(2) - N(3) \\ &\quad + N(12) + N(13) + N(23) \\ &\quad - N(123). \end{aligned}$$

Inclusion/Exclusion (4)

Example When $n = 4$,

$$N_0 = N(\emptyset)$$

$$- N(1) - N(2) - N(3) - N(4)$$

$$+ N(12) + N(13) + N(14) + N(23) + N(24) + N(34)$$

$$- N(123) - N(124) - N(134) - N(234)$$

$$+ N(1234).$$

Inclusion/Exclusion (5)

Observation In general, there are 2^n terms in the inclusion/exclusion formula. How can this possibly be of use?

Derangements

Definition A permutation σ of $\{1, 2, \dots, n\}$ is called a derangement if $\sigma(i) \neq i$ for all $i = 1, 2, \dots, n$.

Example 38754126 and 21436587 are derangements but 57314682 and 75318642 are not.

Exercise Write all derangements of $\{1, 2, 3, 4, 5\}$.

Notation Let d_n denote the number of derangements of $\{1, 2, \dots, n\}$.

Derangements (2)

Inclusion/Exclusion Formula for Derangements

$$\begin{aligned}d_n &= \sum_{S \subseteq \{1, 2, \dots, n\}} (-1)^{|S|} N(S) \\ &= \sum_{0 \leq k \leq n} (-1)^k C(n, k) (n - k)!\end{aligned}$$

Explanation When S is a subset of $\{1, 2, \dots, n\}$ and $|S| = k$, $|N(S)| = (n - k)!$ To see this, note that if σ satisfies P_i and i belongs to S , then $\sigma(i) = i$. So the positions corresponding to elements of S are determined, and the other $n - k$ positions are an arbitrary permutation of the remaining elements.

Surjections (1)

Notation For an integer n , let $[n]$ denote $\{1, 2, \dots, n\}$. Also, let $S(n, m)$ denote the number of surjections from $[n]$ to $[m]$.

Exercise Determine $S(5, 3)$ by hand.

Surjections (2)

Inclusion/Exclusion Formula for Surjections

$$\begin{aligned}d_n &= \sum_{S \subseteq \{1, 2, \dots, n\}} (-1)^{|S|} N(S) \\ &= \sum_{0 \leq k \leq m} (-1)^k C(m, k) (m - k)^n\end{aligned}$$

Explanation When S is a subset of $\{1, 2, \dots, m\}$ and $|S| = k$, $|N(S)| = (m - k)^n$. To see this, note that if f satisfies P_i and i belongs to S , then i is not in the range of f . In other words, f is an function whose domain is $[n]$ and whose range is a set of size $m - k$.

The Euler φ -function

Notation For an integer $n \geq 2$, let $\varphi(n)$ denote the number of elements in $[n]$ which are relatively prime to n .

Example $\varphi(12) = 4$ since 1, 5, 7 and 11 are relatively prime to 12.

Exercise Compute $\varphi(144)$.

Exercise Compute $\varphi(324481700624)$.

The Euler φ -function

Inclusion/Exclusion Formula for Euler φ -Function

Suppose the prime factors of n are: p_1, p_2, \dots, p_k .

Then

$$\varphi(n) = n \left(1 - \frac{1}{p_1}\right) \left(1 - \frac{1}{p_2}\right) \dots \left(1 - \frac{1}{p_k}\right)$$

Explanation When m has the common prime factors p_3, p_7 and p_8 with n , then the number of such m is $n/p_3p_7p_8$.

The Euler φ -function

Example Compute $\varphi(324481700624)$

Maple reports that

$$324481700624 = 2^4(109)(727)(255923)$$

Therefore

$$\begin{aligned}\varphi(324481700624) &= 324481700624(1-1/2)(1-1/109) \\ &\quad (1 - 1/727)(1 - 1/255923) \\ &= 2^3(108)(726)(255922) \\ &= 160530657408.\end{aligned}$$