

Name: _____

Recitation Section: _____

Math 1553 Lecture A, Quiz 5: 2.1, 2.2, 2.3 (10 points, 10 minutes)**Solutions**

Show your work and justify answers where appropriate. If you write the correct answer without sufficient work or justification, you will receive little or no credit.

1. (1 point each) True or False. Circle TRUE if the statement is *always* true. Otherwise, circle FALSE. You do not need to justify your answer on this problem.

a) If A is an $n \times n$ matrix and the equation $Ax = 0$ has only the trivial solution, then every vector in \mathbf{R}^n can be written as a linear combination of the columns of A .

 TRUE

FALSE

(Invertible Matrix Theorem)

b) If A is an invertible $n \times n$ matrix, then $(A^{-1})^{-1} = A$.

 TRUE

FALSE

c) If $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is a linear transformation that is not invertible, then there are vectors x and y in \mathbf{R}^n so that $x \neq y$ but $T(x) = T(y)$.

 TRUE

FALSE

 $(T \text{ is not invertible } \implies T \text{ is not one-to-one})$

d) If A is a 3×4 matrix and B is a 4×2 matrix, then the linear transformation Z defined by $Z(x) = ABx$ has domain \mathbf{R}^2 and codomain \mathbf{R}^3 .

 TRUE

FALSE

 $(AB \text{ is a } 3 \times 2 \text{ matrix})$

2. (3 points) Let $A = \begin{pmatrix} 4 & 5 \\ -1 & 1 \end{pmatrix}$. Find A^{-1} .

Solution.

If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ then $A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$, so

$$A^{-1} = \frac{1}{4 - (-5)} \begin{pmatrix} 1 & -5 \\ 1 & 4 \end{pmatrix} = \begin{pmatrix} \frac{1}{9} & -\frac{5}{9} \\ \frac{1}{9} & \frac{4}{9} \end{pmatrix}.$$

(Turn over to the back side for problem 3!)

3. (3 points) Write two square matrices A and B so that $AB \neq BA$. Demonstrate that $AB \neq BA$ by computing AB and BA .

Solution.

Many examples. We did one in the handwritten notes in class and there is a different example on the PDF slides. Here is a third:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}.$$

$$AB = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \quad BA = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$