

# Section 2.6

## Subspaces

## Outline of Section 2.6

- Definition of subspace
- Examples and non-examples of subspaces
- Spoiler alert: Subspaces are the same as spans
- Spanning sets for subspaces
- Two important subspaces for a matrix:  $\text{Col}(A)$  and  $\text{Nul}(A)$

## Subspaces

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  with:

1. The zero vector is in  $V$ .
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .
3. If  $u$  is in  $V$  and  $c$  is a scalar, then  $cu$  is in  $V$ .

The second and third properties are called “closure under addition” and “closure under scalar multiplication.”

Together, the second and third properties could together be rephrased as: closure under linear combinations.

# Which are subspaces?

A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  of  $\mathbb{R}^n$  with:

1. The zero vector is in  $V$ .
2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .
3. If  $u$  is in  $V$  and  $c$  is a scalar, then  $cu$  is in  $V$ .

Which are subspaces? Which of the 3 properties are satisfied:    1    2    3

1. The unit circle in  $\mathbb{R}^2$

2. The point  $(1, 2, 3)$  in  $\mathbb{R}^3$

3. The  $xy$ -plane in  $\mathbb{R}^3$

4. The  $xy$ -plane together with the  $z$ -axis in  $\mathbb{R}^3$

## Which are subspaces?

Poll

Is the first quadrant of  $\mathbb{R}^2$  a subspace?

1. yes
2. no

## Which are subspaces?

1.  $\left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid a + b = 0 \right\}$

2.  $\left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid a + b = 1 \right\}$

3.  $\left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid ab \neq 0 \right\}$

4.  $\left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbb{R}^2 \mid a, b \text{ rational} \right\}$

## Spans and subspaces

**Fact.** Any  $\text{Span}\{v_1, \dots, v_k\}$  is a subspace.

Why?

**Fact.** Every subspace  $V$  is a span.

Why?

So now we know that three things are the same:

- subspaces
- spans
- planes through 0

So why bother with the word “subspace”? Sometimes easier to check a subset is a subspace than to check it is a span (see null spaces, eigenspaces). Also, it makes sense (and is often useful) to think of a subspace *without a particular spanning set in mind*. Try thinking of other examples where it is useful to have two names for the same thing, like: water /  $\text{H}_2\text{O}$  or free throw / foul shot.

## Column Space and Null Space

$A = m \times n$  matrix.

$\text{Col}(A) =$  **column space** of  $A =$  span of the columns of  $A$

$\text{Nul}(A) =$  **null space** of  $A =$  (set of solutions to  $Ax = 0$ )

Example.  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$

$\text{Col}(A) =$  subspace of  $\mathbb{R}^m$

$\text{Nul}(A) =$  subspace of  $\mathbb{R}^n$

We have already been interested in both. We have been computing null spaces all semester. Also, we have seen that  $Ax = b$  is consistent exactly when  $b$  is in the span of the columns of  $A$ , or,  $b$  is in  $\text{Col}(A)$ .



## Spanning sets for $\text{Nul}(A)$ and $\text{Col}(A)$

Find spanning sets for  $\text{Nul}(A)$  and  $\text{Col}(A)$

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$

## Spanning sets for $\text{Nul}(A)$ and $\text{Col}(A)$

In general:

- our usual parametric solution for  $Ax = 0$  gives a spanning set for  $\text{Nul}(A)$
- the pivot columns of  $A$  form a spanning set for  $\text{Col}(A)$

**Warning!** Not the pivot columns of the reduced matrix.

Notice that the columns of  $A$  form a (possibly larger) spanning set. We'll see later that the above recipe is the smallest spanning set.

## Spanning sets

Find a spanning set for the plane  $2x + 3y + z = 0$  in  $\mathbb{R}^3$ .

## Subspaces and Null spaces

**Fact.** Every subspace is a null space.

Why? Given a spanning set, you can reverse engineer the  $A$ ...

It's actually a little tricky to do this. Given the spanning set, you make those vectors the rows of a matrix, then row reduce and find vector parametric form, and then make those vectors the rows of a new matrix. Try an example! Why does this work? (Stay tuned to Chapter 6 to find out!)

**Example.** Find a matrix  $A$  whose null space is the span of  $(1, 1, 1)$  and  $(1, 2, 3)$ . You should get the matrix  $A = \begin{pmatrix} 1 & -2 & 1 \end{pmatrix}$ .

So now we know that four things are the same:

- subspaces
- spans
- planes through  $0$
- solutions to  $Ax = 0$

(Make sure you understand what we mean when we say these are all the same!)

## So why learn about subspaces?

If subspaces are the same as spans, planes through the origin, and solutions to  $Ax = 0$ , why bother with this new vocabulary word?

The point is that we have been throwing around terms like “3-dimensional plane in  $\mathbb{R}^4$ ” all semester, but we never said what “dimension” and “plane” are. Subspaces give the proper way to define a plane. Soon we will learn the meaning of a dimension of a subspace.

## All the ways

Here are all the ways we know to describe a subspace:

1. As span:

$$\text{Span} \left\{ \left( \begin{array}{c} -1 \\ 1 \\ 0 \end{array} \right), \left( \begin{array}{c} -1 \\ 0 \\ 1 \end{array} \right) \right\}$$

2. As a column space:

$$\text{Col} \left( \begin{array}{cc} -1 & -1 \\ 1 & 0 \\ 0 & 1 \end{array} \right)$$

3. As a null space:

$$\text{Nul} \left( \begin{array}{ccc} 1 & 1 & 1 \end{array} \right)$$

4. As the set of solutions to a homogeneous linear system:

$$x + y + z = 0$$

5. Same, but in set builder notation:

$$\left\{ \left( \begin{array}{c} a \\ b \\ c \end{array} \right) : a + b + c = 0 \right\}$$

## Section 2.6 Summary

- A **subspace** of  $\mathbb{R}^n$  is a subset  $V$  with:
  1. The zero vector is in  $V$ .
  2. If  $u$  and  $v$  are in  $V$ , then  $u + v$  is also in  $V$ .
  3. If  $u$  is in  $V$  and  $c$  is in  $\mathbb{R}$ , then  $cu \in V$ .
- Two important subspaces: **Nul( $A$ )** and **Col( $A$ )**
- Find a spanning set for Nul( $A$ ) by solving  $Ax = 0$  in vector parametric form
- Find a spanning set for Col( $A$ ) by taking pivot columns of  $A$  (not reduced  $A$ )
- Four things are the same: subspaces, spans, planes through 0, null spaces

## Typical exam questions

- Consider the set  $\{(x, y) \in \mathbb{R}^2 \mid xy \geq 0\}$ . Is it a subspace? If not, which properties does it fail?
- Consider the  $x$ -axis in  $\mathbb{R}^3$ . Is it a subspace? If not, which properties does it fail?
- Consider the set  $\{(x, y, z, w) \in \mathbb{R}^4 \mid x + y - z + w = 0\}$ . Is it a subspace? If not, which properties does it fail?
- Find spanning sets for the column space and the null space of

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

- True/False: The set of solutions to a matrix equation is always a subspace.
- True/False: The zero vector is a subspace.