

## Math 6452 - Fall 2017 Homework 2

Work all these problems and talk to me if you have any questions on them, but carefully write up and turn in **only** problems 4, 5, 7, 8, 9, 12. **Due: In class on September 19.**

1. If  $S^2$  is the unit sphere in  $\mathbb{R}^3$  and  $N = (0, 0, 1)$  and  $S = (0, 0, -1)$  then we have the two stereographic coordinate maps  $\pi_N : (S^2 - N) \rightarrow \mathbb{R}^2$  and  $\pi_S : (S^2 - S) \rightarrow \mathbb{R}^2$ . If  $p$  is a point in  $S^2$  not equal to  $N$  or  $S$  then we can use the first to express a tangent vector in  $T_p S^2$  in terms of the basis  $\{\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}\}$  (where we are using Cartesian coordinates  $(x^1, x^2)$  on  $\mathbb{R}^2$ ) as

$$v = v^1 \frac{\partial}{\partial x^1} + v^2 \frac{\partial}{\partial x^2}.$$

Similarly we can use the second to express the same vector in terms of the basis  $\{\frac{\partial}{\partial y^1}, \frac{\partial}{\partial y^2}\}$  (where we are using Cartesian coordinates  $(y^1, y^2)$  on  $\mathbb{R}^2$ ) as

$$v = w^1 \frac{\partial}{\partial y^1} + w^2 \frac{\partial}{\partial y^2}.$$

Write the  $w^i$  in terms of the  $v^i$  (and the coordinate transform  $\pi_S \circ \pi_N^{-1}$ ).

In particular if  $\pi_N(p) = (1, 0)$  and  $v = \frac{\partial}{\partial x^1}$  then express  $v$  in the other coordinate system.

2. Let  $M$  and  $N$  be two smooth manifolds.
  - (a) Show that for  $(p, q) \in M \times N$  we have

$$T_{(p,q)}(M \times N) = (T_p M) \times (T_q N).$$

- (b) If  $\pi : M \times N \rightarrow M : (p, q) \mapsto p$  is the projection map then

$$df_{(p,q)} : T_{(p,q)}(M \times N) \rightarrow T_p M$$

is the projection map  $(v, w) \mapsto v$ .

- (c) Fix a point  $q_0 \in N$  and let  $f : M \rightarrow M \times N : p \mapsto (p, q_0)$  then show that

$$df_p : T_p M \rightarrow T_{(p,q_0)}(M \times N)$$

is given by  $v \mapsto (v, 0)$ .

3. Let  $f : M \rightarrow N$  be a smooth map between smooth manifolds and define  $F : M \rightarrow (M \times N) : p \mapsto (p, f(p))$ . Show that  $dF_p(v) = (v, df_p(v))$ . (Here we are of course using Problem 3 (a) to write the tangent bundle of  $M \times N$  as a product.)
4. If  $f : M \rightarrow N$  is a submersion, then show  $f$  is an open map. (That is show that for any open set  $U$  in  $M$  the image  $f(U)$  is open in  $N$ .)
5. If  $M$  is a compact smooth manifold and  $N$  is a connected smooth manifold, then show that any smooth submersion  $f : M \rightarrow N$  is surjective. Is there a submersion from  $S^2$  to any  $\mathbb{R}^n$ , with  $n > 0$ ?
6. Let  $M$  be a compact smooth manifold and  $N$  a connected smooth manifold. If they both have the same dimension and are non-empty show that any embedding  $f : M \rightarrow N$  is a diffeomorphism.

7. Show that  $\mathbb{C}P^1$  is diffeomorphic to  $S^2$ .

Hint: Using stereographic coordinates on  $S^2$  and our “standard” coordinates on  $\mathbb{C}P^1$  we see both manifolds can be covered by 2 coordinate charts. Study the transition functions between these coordinate charts and see if you can define a map using the coordinate charts.

8. Define the map

$$f : \mathbb{C}P^n \rightarrow \mathbb{C}P^m$$

by

$$f([x^0 : \cdots : x^n]) = [x^0 : \cdots : x^n : 0 : \cdots : 0]$$

where  $n \leq m$ . Show that  $f$  is a smooth embedding. (Notice that this says that  $S^2$  is submanifold of  $\mathbb{C}P^2$ , or any  $\mathbb{C}P^n$  with  $n > 0$  for that matter. Later we will see that this is a “non-trivial”  $S^2$ .)

9. With  $f$  as in the previous problem show that  $\mathbb{C}P^{n+1} - f(\mathbb{C}P^n)$  is diffeomorphic to  $\mathbb{C}^{n+1}$ . (So for example  $\mathbb{C}P^2$  is the union of  $\mathbb{C}P^1 \cong S^2$  and  $\mathbb{C}^2$ . Thus we can think of  $\mathbb{C}P^2$  is the compactification of  $\mathbb{C}^2$  by an “ $S^2$  at infinity”.)

10. A smooth map  $f : (\mathbb{C}^{n+1} - \{(0, \dots, 0)\}) \rightarrow (\mathbb{C}^{k+1} - \{(0, \dots, 0)\})$  is called homogeneous of degree  $k$  if  $f(\lambda p) = \lambda^k f(p)$  for all  $\lambda \neq 0$  and  $p \in (\mathbb{C}^{n+1} - \{(0, \dots, 0)\})$ . Show that  $f$  induces a map

$$\tilde{f} : \mathbb{C}P^n \rightarrow \mathbb{C}P^k.$$

Show this map is smooth.

11. Define the map

$$f : \mathbb{C}P^n \times \mathbb{C}P^m \rightarrow \mathbb{C}P^{nm+n+m}$$

by

$$f([x^0 : \cdots : x^n], [y^0 : \cdots : y^m]) = [x^0 y^0 : x^0 y^1 : \cdots : x^0 y^m : x^1 y^0 : \cdots : x^n y^m].$$

Show  $f$  is a smooth map and that  $f$  it is an embedding. (Notice that this shows, for example, that  $S^2 \times S^2$  is a submanifold of  $\mathbb{C}P^3$ ).

**Note:** The last 4 problems could also have been carried out for real projective spaces.

12. Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a homogeneous polynomial. This implies that there is some integer  $k$  such that

$$f(tx^1, \dots, tx^n) = t^k f(x^1, \dots, x^n)$$

for all  $(x^1, \dots, x^n)$ . Prove that  $f^{-1}(a)$ , for  $a \neq 0$ , is an  $(n - 1)$ -dimensional manifold. Moreover show that if  $a$  and  $b$  are both positive then  $f^{-1}(a)$  and  $f^{-1}(b)$  are diffeomorphic and similarly if  $a$  and  $b$  are both negative. Finally show that if  $a$  and  $b$  have different signs that  $f^{-1}(a)$  and  $f^{-1}(b)$  do not have to be diffeomorphic by considering  $f(x, y, z) = x^2 + y^2 - z^2$ .

Hint: It might be good to use the famous Euler identity for homogeneous functions

$$\sum_{i=1}^n x^i \frac{\partial f}{\partial x^i} = kf,$$

(you don’t need to prove this identity, though feel free to if you like) to prove that 0 is the only critical value of  $f$ . To find the diffeomorphism consider the map  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  obtained by multiplication by an appropriate root of  $a/b$ .