

Math 6452 - Fall 2018 Homework 5

Work all these problems and talk to me if you have any questions on them, but carefully write up and turn in **only** problems 2, 5, 6, 7, 8, 12 **Due: In class on November 9.**

1. For any finite dimensional vector space show that there are canonical isomorphisms $V \otimes \mathbb{R} \cong V \cong \mathbb{R} \otimes V$.
2. For finite dimensional vector spaces V and W show there is a canonical isomorphism $V^* \otimes W \cong \text{Hom}(V, W)$.
3. Let $\omega^1, \dots, \omega^k$ be covectors in V^* . Show they are linearly dependent if and only if $\omega^1 \wedge \dots \wedge \omega^k = 0$.
4. If $\{\omega^1, \dots, \omega^k\}$ and $\{\eta^1, \dots, \eta^k\}$ are linearly independent covectors in V^* , then show they have the same span if and only if $\omega^1 \wedge \dots \wedge \omega^k = c \eta^1 \wedge \dots \wedge \eta^k$ for some real number c . Show that $c = \det(A)$ where A is the matrix $(a_{i,j})$ and the $a_{i,j}$ are determined by $\omega_i = \sum_j a_{i,j} \eta_j$.
5. Let M be a smooth manifold and let $\omega \in \Gamma(T^k M)$ be a tensor field. Consider the map

$$\Psi_\sigma : (\mathcal{X}(M) \times \dots \times \mathcal{X}(M)) \rightarrow C^\infty(M) : (v_1, \dots, v_k) \mapsto \omega(v_1, \dots, v_k)$$

here $\mathcal{X}(M)$ is the set of vector fields. Show that this map is multilinear over $C^\infty(M)$. Moreover show that given any multilinear map over $C^\infty(M)$

$$\Psi : (\mathcal{X}(M) \times \dots \times \mathcal{X}(M)) \rightarrow C^\infty(M)$$

it is induced from some tensor field by the above construction.

6. Define the 1-form ω on $\mathbb{R}^2 - \{(0, 0)\}$ by

$$\omega(x, y) = \left(\frac{-y}{x^2 + y^2} \right) dx + \left(\frac{x}{x^2 + y^2} \right) dy$$

- (a) Compute $\int_C \omega$ where C is a circle of radius r about the origin.
- (b) Is ω the differential of a function on $\mathbb{R}^2 - \{(0, 0)\}$? Explain why or why not.
7. Prove that a 1-form α on S^1 is the differential of a function if and only if $\int_{S^1} \alpha = 0$.
8. Prove that the first De Rham cohomology of S^1 is $H_{DR}^1(S^1) \cong \mathbb{R}$.
Hint: Show that if α is a fixed 1-form on S^1 such that $\int_{S^1} \alpha \neq 0$ then for any other 1-form ω there is a real number c such that $\omega = c\alpha + df$ for some function f .
9. Consider the forms on \mathbb{R}^3

$$f \in \omega^0(\mathbb{R}^3), \quad f dx + g dy + h dz \in \omega^1(\mathbb{R}^3), \quad \text{and}$$

$$f dy \wedge dz + g dz \wedge dy + h dx \wedge dy \in \omega^2(\mathbb{R}^3).$$

Compute their exterior derivatives. Do they look like anything from vector calculus?

10. Given a vector space V and a vector $v \in V$ define the interior product

$$\iota_v : \Lambda^k(V) \rightarrow \Lambda^{k-1}(V)$$

as follows: given $\omega \in \Lambda^k(V)$ define $\iota_v\omega$ to be the $(k-1)$ form:

$$\iota_v\omega(v_1, \dots, v_{k-1}) = \omega(v, v_1, \dots, v_{k-1}).$$

If $\omega \in \Lambda^k(V)$ and $\eta \in \Lambda^l(V)$ then show that

$$\iota_v(\omega \wedge \eta) = (\iota_v\omega) \wedge \eta + (-1)^k\omega \wedge (\iota_v\eta).$$

11. On \mathbb{R}^{2n} with coordinates $(x^1, y^1, \dots, x^n, y^n)$ define the 1-form $\lambda = \frac{1}{2} \sum (x^i dy^i - y^i dx^i)$. Compute $d\lambda$ and $(d\lambda)^n$ (this means take the wedge product of $d\lambda$ with itself n times, for example $(d\lambda)^3 = (d\lambda) \wedge (d\lambda) \wedge (d\lambda)$). The 2-form $d\lambda$ is called the standard symplectic form on \mathbb{R}^{2n} .
12. Suppose V is an n -dimensional vector space. Given a linear map

$$L : V \rightarrow V$$

there is an induced linear map

$$L^* : \wedge^n(V) \rightarrow \wedge^n(V).$$

Since $\wedge^n(V)$ is 1-dimensional this map is simply multiplication by a constant. We will denote this constant $\det(L)$. Prove the following

- (a) If you choose a basis e_1, \dots, e_n for V then L can be written as a matrix A_L , $\det(L) = \det(A_L)$, where the determinant of a matrix has the usual definition.
- (b) $\det(L \circ S) = \det(L) \det(S)$.
- (c) If L is the identity map then $\det(L) = 1$.
- (d) L is an isomorphism if and only if $\det(L) \neq 0$ and in this case $\det(L^{-1}) = (\det(L))^{-1}$.