

Work the following problems and hand in your solutions. You may work together with other people in the class, but you must each write up your solutions independently. A subset of these will be selected for grading. Write LEGIBLY on the FRONT side of the page only, and STAPLE your pages together.

Definition. A set of vectors $\{f_n\}_{n \in \mathbb{N}}$ in a Hilbert space H is a *tight frame* for H if there exists a number $A > 0$ so that

$$\forall f \in H, \quad \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 = A \|f\|^2.$$

The number A is the *frame bound*. The *frame operator* is the mapping $S : H \rightarrow H$ defined by

$$Sf = \sum_{n=1}^{\infty} \langle f, f_n \rangle f_n \quad \text{for } f \in H.$$

We proved in class that this series converges for each f (this is a consequence of the fact that $\sum |\langle f, f_n \rangle|^2 < \infty$). □

1. Assume that $\{f_n\}_{n \in \mathbb{N}}$ is a tight frame with frame bound A .

(a) Show directly that S and $S - AI$ are self-adjoint, where I is the identity operator.

(b) Show that $S = AI$.

Hint: The norm of a self-adjoint operator T on a Hilbert space H can be computed using the formula

$$\|T\| = \sup_{\|f\|=1} |\langle Tf, f \rangle|.$$

(c) Show directly that the following three statements are equivalent.

i. $\|f_n\|^2 = A$ for every n .

ii. $\{f_n\}_{n \in \mathbb{N}}$ is an orthogonal (but not necessarily orthonormal) sequence with no zero elements. That is, $\langle f_m, f_n \rangle = 0$ if $m \neq n$, and every $f_n \neq 0$.

iii. $\{f_n\}_{n \in \mathbb{N}}$ is a Schauder basis for H .

2. (a) Show that if $T > 1$ then $\{e^{2\pi i n T x}\}_{n \in \mathbb{Z}}$ is incomplete in $L^2[0, 1]$.

(b) Show that if $0 < T < 1$, then $\{e^{2\pi i n T x}\}_{n \in \mathbb{Z}}$ is a tight frame for $L^2[0, 1]$. What is the frame bound?

(c) Show that $\{e^{2\pi i n T x}\}_{n \in \mathbb{Z}}$ is not a Schauder basis for $L^2[0, 1]$ when $0 < T < 1$. In particular, find two different ways to write the constant function 1 (on the domain $[0, 1]$) as an infinite linear combination of the exponentials $e^{2\pi i n T x}$.

3. (a) Prove the following *perturbation result* for frames. Suppose that $\{f_n\}_{n \in \mathbb{N}}$ is a frame for a Hilbert space H with frame bounds A, B . This means that $\{f_n\}_{n \in \mathbb{N}}$ is not necessarily tight, instead, we have that

$$\forall f \in H, \quad A \|f\|^2 \leq \sum_{n=1}^{\infty} |\langle f, f_n \rangle|^2 \leq B \|f\|^2.$$

Suppose also that $\{g_n\}$ is such that $\{f_n - g_n\}_{n \in \mathbb{N}}$ satisfies an *upper* frame bound condition of the form

$$\forall f \in H, \quad \sum_{n=1}^{\infty} |\langle f, f_n - g_n \rangle|^2 \leq R \|f\|^2.$$

(We say then that $\{f_n - g_n\}_{n \in \mathbb{N}}$ is a *Bessel sequence* with *Bessel bound* R ; the sequence $\{f_n - g_n\}_{n \in \mathbb{N}}$ need not have a positive lower frame bound).

Show that $\{g_n\}$ is a frame if $R < A$. Hint: use the triangle inequality in ℓ^2 :

$$\left(\sum_{n=1}^{\infty} |c_n + d_n|^2 \right)^{1/2} \leq \left(\sum_{n=1}^{\infty} |c_n|^2 \right)^{1/2} + \left(\sum_{n=1}^{\infty} |d_n|^2 \right)^{1/2}.$$

(b) Show that if $\{h_n\}$ is a sequence in H which satisfies

$$R = \sum_n \|h_n\|^2 < \infty$$

then $\{h_n\}$ is a Bessel sequence with bound R .

(c) The exponentials $\{e^{2\pi i n T x}\}_{n \in \mathbb{Z}}$ are a frame in $L^2[-\frac{1}{2}, \frac{1}{2}]$ when $T < 1$. Use Problem 2, together with parts (a) and (b) of this problem, to formulate and prove a theorem establishing a sufficient condition on numbers $t_n \in \mathbb{R}$ so that $\{e^{2\pi i t_n x}\}_{n \in \mathbb{Z}}$ is a frame for $L^2[-\frac{1}{2}, \frac{1}{2}]$. Do you think your result is optimal?