

LINEAR


ALGEBRA

Week 6

Section 2.9 : Dimension and Rank

Chapter 2 : Matrix Algebra

Math 1554 Linear Algebra



Topics and Objectives

Topics

We will cover these topics in this section.

1. Coordinates, relative to a basis.
2. Dimension of a subspace.
3. The Rank of a matrix

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Calculate the coordinates of a vector in a given basis.
2. Characterize a subspace using the concept of dimension (or cardinality).
3. Characterize a matrix using the concepts of rank, column space, null space.
4. Apply the Rank, Basis, and Matrix Invertibility theorems to describe matrices and subspaces.

Section 2.9 : Dimension and Rank

Chapter 2 : Matrix Algebra
Math 1554 Linear Algebra



Section 2.9 Slide 183

Topics and Objectives

Topics

We will cover these topics in this section.

1. Coordinates, relative to a basis.
2. Dimension of a subspace.
3. The Rank of a matrix.

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Calculate the coordinates of a vector in a given basis.
2. Characterize a subspace using the concept of dimension (or cardinality).
3. Characterize a matrix using the concepts of rank, column space, null space.
4. Apply the Rank, Basis, and Matrix Invertibility theorems to describe matrices and subspaces.

Section 2.9 Slide 184

Course Schedule


Cancellations due to inclement weather will likely result in cancelling review lectures and possibly moving through course material at a

Week Dates	Mon Lecture	Tue Studio	Wed Lecture	Thu Studio	Fri Lecture
1 8/21 - 8/25	1.1	WS1.1	1.2	WS1.2	1.3
2 8/28 - 9/1	1.4	WS1.3,1.4	1.5	WS1.5	1.7
3 9/4 - 9/8	Break	WS1.7	1.8	WS1.8	1.9
4 9/11 - 9/15	2.1	WS1.9,2.1	Exam 1 Review	Cancelled	2.2
5 9/18 - 9/22	2.3,2.4	WS2.2,2.3	2.5	WS2.4,2.5	2.8
6 9/25 - 9/29	2.9	WS2.8,2.9	3.1,3.2	WS3.1,3.2	3.3
7 10/2 - 10/6	4.9	WS3.3,4.9	5.1,5.2	WS5.1,5.2	5.2
8 10/9 - 10/13	Break	Break	Exam 2 Review	Cancelled	5.3
9 10/16 - 10/20	5.3	WS5.3	5.5	WS5.5	6.1
10 10/23 - 10/27	6.1,6.2	WS6.1	6.2	WS6.2	6.3
11 10/30 - 11/3	6.4	WS6.3,6.4	6.6,6.5	WS6.6,6.5	6.5
12 11/6 - 11/10	6.6	WS6.5,6.6	Exam 3 Review	Cancelled	PageRank
13 11/13 - 11/17	7.1	WSPageRank	7.2	WS7.1,7.2	7.3
14 11/20 - 11/24	7.3,7.4	WS7.2,7.3	Break	Break	Break
15 11/27 - 12/1	7.4	WS7.3,7.4	7.4	WS7.4	7.4
16 12/4 - 12/8	Last lecture	Last Studio	Reading Period		
17 12/11 - 12/15	Final Exams	MATH 1554 Common Final Exam Tuesday, December 12th at 6pm			

Choice of Basis

Key idea: There are many possible choices of basis for a subspace. Our choice can give us dramatically different properties.

Example: sketch $b_1 + b_2$ for the two different coordinate systems below.



Section 2.9 Slide 185

FACT: \mathbb{R}^n is a subspace of \mathbb{R}^n

$1b_1 + 1b_2$
coordinates in $\{b_1, b_2\}$ would be "1, 1, 0, 0, ..."

$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$ for \mathbb{R}^3
 e_1, e_2, \dots, e_n for \mathbb{R}^n

Coordinates

Definition

Let $B = \{b_1, \dots, b_p\}$ be a basis for a subspace H . If \vec{x} is in H , then **coordinates of \vec{x} relative to B** are the weights (scalars) c_1, \dots, c_p so that

$$\vec{x} = c_1 b_1 + \dots + c_p b_p$$

And

$$[\vec{x}]_B = \begin{bmatrix} c_1 \\ \vdots \\ c_p \end{bmatrix}$$

is the **coordinate vector of \vec{x} relative to B** , or the **B -coordinate vector of \vec{x}**

Section 2.9 Slide 186

The **coordinates** in the basis $\{e_1, e_2, \dots\}$

$$\begin{bmatrix} -3 \\ 2 \\ 1 \end{bmatrix} = -3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Example 1

Let $v_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, $v_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, and $\vec{x} = \begin{bmatrix} 3 \\ 3 \\ 5 \end{bmatrix}$. Verify that \vec{x} is in the span of $B = \{v_1, v_2\}$, and calculate $[\vec{x}]_B$.


$$[\vec{x}]_{v_1, v_2} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Handwritten notes: "The components of \vec{x} in basis $\{v_1, v_2\}$ "

$$c_1 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 5 \end{bmatrix}$$

Section 2.9 Slide 107

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 0 & 1 & 3 \\ 0 & 0 & 2 \end{array} \right] \sim \dots \sim \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 3 \\ 0 & 0 & 2 \end{array} \right]$$



Section 2.9 Slide 108

Dimension



Definition


The **dimension** (or cardinality) of a non-zero subspace H , $\dim H$, is the number of vectors in a basis of H . We define $\dim\{0\} = 0$.

Theorem

Any two choices of bases B_1 and B_2 of a non-zero subspace H have the same dimension.

Examples:

- $\dim \mathbb{R}^n = n$
- $H = \{(x_1, \dots, x_n) : x_1 + \dots + x_n = 0\}$ has dimension $n-1$.
- $\dim(\text{Null } A)$ is the number of **free vars** (circled in diagram).
- $\dim(\text{Col } A)$ is the number of **pivot cols** (circled in diagram).



Rank

Definition
The rank of a matrix A is the dimension of its column space.

Example 2: Compute $\text{rank}(A)$ and $\dim(\text{Null}(A))$.

$$A = \begin{bmatrix} 4 & 5 & -3 & 8 \\ 7 & -4 & -3 & 9 \\ 8 & -5 & 2 & 4 \\ 9 & -9 & 6 & -6 \end{bmatrix} \sim \dots \sim \begin{bmatrix} 4 & 5 & -3 & 8 \\ 0 & 0 & 2 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\text{rank}(A) = 3$

basis for $\text{Col } A$ is $\left\{ \begin{bmatrix} 4 \\ 7 \\ 8 \\ 9 \end{bmatrix}, \begin{bmatrix} 5 \\ -4 \\ -5 \\ -9 \end{bmatrix}, \begin{bmatrix} -3 \\ -3 \\ 2 \\ 6 \end{bmatrix} \right\}$

Section 2.9 Slide 109

$\dim \text{Null}(A) = 2$

$\text{Null}(A)$ is a plane in \mathbb{R}^4

$\text{Null}(A) = \{ \vec{x} : A\vec{x} = \vec{0} \}$

basis of $\text{Null } A$ is $\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\}$

from parametric we can form for solving for $A\vec{x} = \vec{0}$.

Rank, Basis, and Invertibility Theorems

Theorem (Rank Theorem)

If a matrix A has n columns, then $\text{rank } A + \dim(\text{Null } A) = n$.

Theorem (Basis Theorem)

Any two bases for a subspace have the same cardinality.

Theorem (Invertibility Theorem)

Let A be an $n \times n$ matrix. These conditions are equivalent.

- A is invertible.
- The columns of A are a basis for \mathbb{R}^n .
- $\text{rank } A = n$.
- $\text{rank } A = \dim(\text{Col } A) = n$.
- $\text{Null } A = \{\vec{0}\}$.

Proof: Every col of A is either a pivot col or a free column.

MIT DLC

if they span \mathbb{R}^n & they are lin. ind. pick some random $b \in \mathbb{R}^n$

$Ax = b$ is always consistent
 $[b]$ is in $\text{Col } A = \vec{x}$.

Section 2.9 Slide 110

Example

If possible, give an example of a 2×3 matrix A , in reduced echelon form, with the given properties.

a) $\text{rank}(A) = 3$

NP

$$\begin{pmatrix} \textcircled{1} & * & * \\ * & \textcircled{1} & * \end{pmatrix} = A$$

b) $\text{rank}(A) = 2$

$$\begin{bmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \end{bmatrix}$$

RREF ✓ $\text{rank } A = 2$ ✓

c) $\dim(\text{Null}(A)) = 2$

$$\begin{bmatrix} 0 & \textcircled{1} & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

RREF ✓ $\text{rank } A = 1$ $\dim \text{Null } A = 2$ ✓

d) $\text{Null } A = \{0\}$

NP

EXAMPLE 1 Let $\mathbf{v}_1 = \begin{bmatrix} 3 \\ 6 \\ 2 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} 3 \\ 12 \\ 7 \end{bmatrix}$, and $B = \{\mathbf{v}_1, \mathbf{v}_2\}$. Then

B is a basis for $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ because \mathbf{v}_1 and \mathbf{v}_2 are linearly independent. Determine if \mathbf{x} is in H , and if it is, find the coordinate vector of \mathbf{x} relative to B .




FIGURE 1 A coordinate system on a plane H in \mathbb{R}^3 .

Check that $\begin{bmatrix} 2 \\ 3 \end{bmatrix}$ is the

coordinate vector for \vec{x} in the basis

$\{\mathbf{v}_1, \mathbf{v}_2\}$.

$$2 \begin{bmatrix} 3 \\ 6 \\ 2 \end{bmatrix} + 3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 12 \\ 7 \end{bmatrix}$$

it worked!

THEOREM
The Invertible Matrix Theorem (continued)

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- m. The columns of A form a basis of \mathbb{R}^n .
- n. $\text{Col } A = \mathbb{R}^n$
- o. $\dim \text{Col } A = n$
- p. $\text{rank } A = n$
- q. $\text{Nul } A = \{\mathbf{0}\}$
- r. $\dim \text{Nul } A = 0$

2.9 EXERCISES

In Exercises 1 and 2, find the vector \mathbf{x} determined by the given coordinate vector $[\mathbf{x}]_{\mathcal{B}}$ and the given basis \mathcal{B} . Illustrate your answer with a figure, as in the solution of Practice Problem 2.

$$1. \mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

$$2. \mathcal{B} = \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$

In Exercises 3–6, the vector \mathbf{x} is in a subspace H with a basis $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$. Find the \mathcal{B} -coordinate vector of \mathbf{x} .

$$3. \mathbf{b}_1 = \begin{bmatrix} 1 \\ -4 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -2 \\ 7 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} -3 \\ 7 \end{bmatrix}$$


$$4. \mathbf{b}_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -3 \\ 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} -7 \\ 5 \end{bmatrix}$$

$$5. \mathbf{b}_1 = \begin{bmatrix} 1 \\ 5 \\ -3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -3 \\ -7 \\ 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 4 \\ 10 \\ -7 \end{bmatrix}$$


$$6. \mathbf{b}_1 = \begin{bmatrix} -3 \\ 1 \\ -4 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 7 \\ 5 \\ -6 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 11 \\ 0 \\ 7 \end{bmatrix}$$

160 CHAPTER 2 Matrix Algebra

7. Let $\mathbf{b}_1 = \begin{bmatrix} 3 \\ 0 \end{bmatrix}$, $\mathbf{b}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 7 \\ -2 \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$, and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$. Use the figure to estimate $[\mathbf{w}]_{\mathcal{B}}$ and $[\mathbf{x}]_{\mathcal{B}}$. Confirm your estimate of $[\mathbf{x}]_{\mathcal{B}}$ by using it and $\{\mathbf{b}_1, \mathbf{b}_2\}$ to compute \mathbf{x} .



8. Let $\mathbf{b}_1 = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$, $\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} -2 \\ 3 \end{bmatrix}$, $\mathbf{y} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$, $\mathbf{z} = \begin{bmatrix} -1 \\ -2.5 \end{bmatrix}$, and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$. Use the figure to estimate $[\mathbf{x}]_{\mathcal{B}}$, $[\mathbf{y}]_{\mathcal{B}}$, and $[\mathbf{z}]_{\mathcal{B}}$. Confirm your estimates of $[\mathbf{y}]_{\mathcal{B}}$ and $[\mathbf{z}]_{\mathcal{B}}$ by using them and $\{\mathbf{b}_1, \mathbf{b}_2\}$ to compute \mathbf{y} and \mathbf{z} .



Exercises 9–12 display a matrix A and an echelon form of A . Find bases for $\text{Col } A$ and $\text{Nul } A$, and then state the dimensions of these subspaces.

$$9. A = \begin{bmatrix} 1 & -3 & 2 & -4 \\ -3 & 9 & -1 & 5 \\ 2 & -6 & 4 & -3 \\ -4 & 12 & 2 & -7 \end{bmatrix} \sim \begin{bmatrix} 1 & -3 & 2 & -4 \\ 0 & 0 & 5 & -7 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$10. A = \begin{bmatrix} 1 & -2 & 9 & 5 & 4 \\ 1 & -1 & 6 & 5 & -3 \\ -2 & 0 & -6 & 1 & -2 \\ 4 & 1 & 9 & 1 & -9 \end{bmatrix} \sim \begin{bmatrix} 1 & -2 & 9 & 5 & 4 \\ 0 & 1 & -3 & 0 & -7 \\ 0 & 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$11. A = \begin{bmatrix} 1 & 2 & -5 & 0 & -1 \\ 2 & 5 & -8 & 4 & 3 \\ -3 & -9 & 9 & -7 & -2 \\ 3 & 10 & -7 & 11 & 7 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 2 & -5 & 0 & -1 \\ 0 & 1 & 2 & 4 & 5 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$12. A = \begin{bmatrix} 1 & 2 & -4 & 3 & 3 \\ 5 & 10 & -9 & -7 & 8 \\ 4 & 8 & -9 & -2 & 7 \\ -2 & -4 & 5 & 0 & -6 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 2 & -4 & 3 & 3 \\ 0 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 & -5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

In Exercises 13 and 14, find a basis for the subspace spanned by the given vectors. What is the dimension of the subspace?

$$13. \left\{ \begin{bmatrix} 1 \\ -3 \\ 2 \\ -4 \end{bmatrix}, \begin{bmatrix} -3 \\ 9 \\ -6 \\ 12 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 4 \\ 2 \end{bmatrix}, \begin{bmatrix} -4 \\ 5 \\ -3 \\ 7 \end{bmatrix} \right\}$$

$$14. \left\{ \begin{bmatrix} 1 \\ -1 \\ -2 \\ 5 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ -1 \\ 6 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ -6 \\ 8 \end{bmatrix}, \begin{bmatrix} -1 \\ 4 \\ -7 \\ 7 \end{bmatrix}, \begin{bmatrix} 3 \\ -8 \\ 9 \\ -5 \end{bmatrix} \right\}$$

15. Suppose a 3×5 matrix A has three pivot columns. Is $\text{Col } A = \mathbb{R}^3$? Is $\text{Nul } A = \mathbb{R}^2$? Explain your answers.

16. Suppose a 4×7 matrix A has three pivot columns. Is $\text{Col } A = \mathbb{R}^3$? What is the dimension of $\text{Nul } A$? Explain your answers.

In Exercises 17 and 18, mark each statement True or False. Justify each answer. Here A is an $m \times n$ matrix.

17. a. If $\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a basis for a subspace H and if $\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p$, then c_1, \dots, c_p are the coordinates of \mathbf{x} relative to the basis \mathcal{B} .
 b. Each line in \mathbb{R}^n is a one-dimensional subspace of \mathbb{R}^n .
 c. The dimension of $\text{Col } A$ is the number of pivot columns of A .
 d. The dimensions of $\text{Col } A$ and $\text{Nul } A$ add up to the number of columns of A .
 e. If a set of p vectors spans a p -dimensional subspace H of \mathbb{R}^n , then these vectors form a basis for H .
18. a. If \mathcal{B} is a basis for a subspace H , then each vector in H can be written in only one way as a linear combination of the vectors in \mathcal{B} .
 b. If $\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a basis for a subspace H of \mathbb{R}^n , then the correspondence $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ makes H look and act the same as \mathbb{R}^p .

- c. The dimension of $\text{Nul } A$ is the number of variables in the equation $A\mathbf{x} = \mathbf{0}$.
- d. The dimension of the column space of A is $\text{rank } A$.
- e. If H is a p -dimensional subspace of \mathbb{R}^n , then a linearly independent set of p vectors in H is a basis for H .

In Exercises 19–24, justify each answer or construction.

- 19. If the subspace of all solutions of $A\mathbf{x} = \mathbf{0}$ has a basis consisting of three vectors and if A is a 5×7 matrix, what is the rank of A ?
- 20. What is the rank of a 4×5 matrix whose null space is three-dimensional?
- 21. If the rank of a 7×6 matrix A is 4, what is the dimension of the solution space of $A\mathbf{x} = \mathbf{0}$?
- 22. Show that a set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_5\}$ in \mathbb{R}^n is linearly dependent when $\dim \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_5\} = 4$.
- 23. If possible, construct a 3×4 matrix A such that $\dim \text{Nul } A = 2$ and $\dim \text{Col } A = 2$.
- 24. Construct a 4×3 matrix with rank 1.
- 25. Let A be an $n \times p$ matrix whose column space is p -dimensional. Explain why the columns of A must be linearly independent.
- 26. Suppose columns 1, 3, 5, and 6 of a matrix A are linearly independent (but are not necessarily pivot columns) and the rank of A is 4. Explain why the four columns mentioned must be a basis for the column space of A .

Section 3.1 : Introduction to Determinants

Chapter 3 : Determinants

Math 1554 Linear Algebra

Topics and Objectives

Topics

We will cover these topics in this section.

1. The definition and computation of a determinant
2. The determinant of triangular matrices

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Compute determinants of $n \times n$ matrices using a cofactor expansion.
2. Apply theorems to compute determinants of matrices that have particular structures.

Topics and Objectives

Topics

We will cover these topics in this section.

1. The definition and computation of a determinant
2. The determinant of triangular matrices

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Compute determinants of $n \times n$ matrices using a cofactor expansion.
2. Apply theorems to compute determinants of matrices that have particular structures.

Section 3.1: Introduction to Determinants

Chapter 3 : Determinants

Math 1554 Linear Algebra

Section 3.1 Slide 172

$$\det: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$$

eg. $\det A = 4$

A Definition of the Determinant

Suppose A is $n \times n$ and has elements a_{ij} .

1. If $n = 1$, $A = [a_{11}]$, and has determinant $\det A = a_{11}$.
2. Inductive case: for $n > 1$,

$$\det A = a_{11} \det A_{11} - a_{12} \det A_{12} + \dots + (-1)^{1+n} a_{1n} \det A_{1n}$$

where A_{ij} is the submatrix obtained by eliminating row i and column j of A .

minor

Example

$$A = \begin{pmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{pmatrix} \Rightarrow A_{2,3} = \begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \end{pmatrix}$$

Section 3.1 Slide 174

5	9/18 - 9/22	2.3.2.4	WS2.2.2.3	2.5	WS2.4.2.5	2.8
6	9/25 - 9/29	2.9	WS2.8.2.9	3.1.3.2	WS3.1.3.2	3.3
7	10/2 - 10/6	4.9	WS3.3.4.9	5.1.5.2	WS5.1.5.2	5.2
8	10/9 - 10/13	Break	Break	Exam 2, Review	Cancelled	5.3

Example 1

Compute $\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\det \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} = 1 \cdot 2 - 1 \cdot 2 = 0$$

$$\det \begin{bmatrix} 1 & 0 \\ 4 & 3 \end{bmatrix} = 3 \cdot 1 - 0 \cdot 4 = 3$$

Section 3.1 Slide 176

Example 2 Cofactor expansion

Compute $\det \begin{pmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{pmatrix} = \begin{vmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{vmatrix}$

① Choose any row or any column

Cofactors

Cofactors give us a more convenient notation for determinants.

Definition: Cofactor

The (i, j) cofactor of an $n \times n$ matrix A is

$$C_{ij} = (-1)^{i+j} \det A_{ij}$$

The pattern for the negative signs is

$$\begin{pmatrix} + & - & + & \dots \\ - & + & - & \dots \\ + & - & + & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

$$a_{ij} \rightarrow (-1)^{i+j}$$

add $i+j$ -
over $i+j$ +

$$\det \begin{pmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{pmatrix} = a_{11} \det(A_{11}) - a_{12} \det(A_{12}) + a_{13} \det(A_{13})$$

$$= (1 \cdot 0 - (-1)(2)) + 5 \left(2 \cdot 0 - (-1)(0) \right) + 0(2 \cdot 2 - 4 \cdot 0)$$

$$= 2 + 5(0) + 0(4) = 2 + 0 + 0 = 2$$



$$\det \begin{pmatrix} 1 & -5 & 0 \\ 2 & 4 & -1 \\ 0 & 2 & 0 \end{pmatrix} = -a_{12} \det(A_{12}) + a_{22} \det(A_{22}) - a_{32} \det(A_{32})$$


$$= -(-5) \begin{vmatrix} 2 & -1 \\ 0 & 0 \end{vmatrix} + 4 \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} - 2 \begin{vmatrix} 1 & 0 \\ 2 & -1 \end{vmatrix}$$

$$= 5(0) + 4(0) - 2(-1) = 0 + 0 + 2 = 2$$

Theorem
The determinant of a matrix A can be computed down any row or column of the matrix. For instance, down the j^{th} column, the determinant is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj}$$

This gives us a way to calculate determinants more efficiently.



Example 3

Compute the determinant of

$$\begin{pmatrix} 5 & 4 & 3 \\ 0 & 1 & 2 \\ 0 & -1 & 1 \\ 0 & 1 & 3 \end{pmatrix} = A$$

See says!! Zeros are great!!

$$\det A = 5 \begin{vmatrix} 2 & 0 \\ -1 & 1 \\ 1 & 3 \end{vmatrix} - 0 + 0 - 0$$

$$= 5 \cdot (0 - 0 + 3 \begin{vmatrix} 1 & 2 \\ -1 & 1 \end{vmatrix})$$

$$= 5 \cdot 3(1 - (-2)) = 5 \cdot 3^2 = 45$$

Triangular Matrices

Theorem

If A is a triangular matrix then

$$\det A = a_{11}a_{22}a_{33} \cdots a_{nn}$$

Example 4

Compute the determinant of the matrix. Empty elements are zero.

$$\begin{bmatrix} 2 & 1 & & & \\ & 2 & 1 & & \\ & & 2 & 1 & \\ & & & 2 & 1 \\ & & & & 2 \end{bmatrix}$$

Section 1.1 Slide 188

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 3 & 2 & 0 & 0 \\ 4 & -1 & 5 & 0 \\ 1 & 0 & -3 & 1 \end{bmatrix}$$

$$\det A = 1 \begin{vmatrix} 2 & 0 & 0 \\ -1 & 5 & 0 \\ 0 & -3 & -1 \end{vmatrix} = -0 + 0 - 0$$

$$= 1 \cdot 2 \cdot \begin{vmatrix} 5 & 0 \\ -3 & -1 \end{vmatrix} = -0 + 0$$

$$= 1 \cdot 2 \cdot (5 \cdot (-1) - 0 \cdot (-3))$$

$$= 1 \cdot 2 \cdot 5 \cdot (-1) = -10$$

Computational Efficiency

Note that computation of a co-factor expansion for an $N \times N$ matrix requires roughly $N!$ multiplications.

- A 10×10 matrix requires roughly $10! = 3.6$ million multiplications
- A 20×20 matrix requires $20! \approx 2.4 \times 10^{18}$ multiplications

This doesn't mean that determinants are not useful.

- We will explore other methods that further the efficiency of their calculation.
- Determinants are very useful in multivariable calculus for solving certain integration problems.

10×10

$10!$

$$5 \times 5 \begin{bmatrix} * & * & * & * & * \\ * & & & & \\ * & & & & \\ * & & & & \\ * & & & & \end{bmatrix}$$

$$\begin{bmatrix} * & * & * & * & * \\ * & & & & \\ * & & & & \\ * & & & & \\ * & & & & \end{bmatrix} = \begin{bmatrix} * & | & * & * & * \\ * & & & & \\ * & & & & \\ * & & & & \\ * & & & & \end{bmatrix} \begin{matrix} 5 \\ 4 \times 4's \end{matrix}$$

$$\begin{bmatrix} * & * & * \\ * & * & * \\ * & * & * \end{bmatrix} \begin{matrix} 4 \\ 3 \times 3's \end{matrix}$$

$$\begin{bmatrix} * & * \\ * & * \end{bmatrix} \begin{matrix} 2 \times 2's \\ 2 \times 2's!! \end{matrix}$$

60
 $2 \times 2's!!$

3.1 EXERCISES

Compute the determinants in Exercises 1–8 using a cofactor expansion across the first row. In Exercises 1–4, also compute the determinant by a cofactor expansion down the second column.

$$1. \begin{vmatrix} 3 & 0 & 4 \\ 2 & 3 & 2 \\ 0 & 5 & -1 \end{vmatrix} \quad 2. \begin{vmatrix} 0 & 4 & 1 \\ 2 & -3 & 0 \\ 2 & 3 & 1 \end{vmatrix}$$

$$3. \begin{vmatrix} 2 & -2 & 3 \\ 3 & 1 & 2 \\ 1 & 3 & -1 \end{vmatrix} \quad 4. \begin{vmatrix} 1 & 2 & 4 \\ 3 & 1 & 1 \\ 2 & 4 & 2 \end{vmatrix}$$

$$5. \begin{vmatrix} 2 & 3 & -3 \\ 4 & 0 & 3 \\ 6 & 1 & 5 \end{vmatrix} \quad 6. \begin{vmatrix} 5 & -2 & 2 \\ 0 & 3 & -3 \\ 2 & -4 & 7 \end{vmatrix}$$

170 CHAPTER 3 Determinants

$$7. \begin{vmatrix} 4 & 3 & 0 \\ 6 & 5 & 2 \\ 9 & 7 & 3 \end{vmatrix} \quad 8. \begin{vmatrix} 4 & 1 & 2 \\ 4 & 0 & 3 \\ 3 & -2 & 5 \end{vmatrix}$$

Compute the determinants in Exercises 9–14 by cofactor expansions. At each step, choose a row or column that involves the least amount of computation.

$$9. \begin{vmatrix} 4 & 0 & 0 & 5 \\ 1 & 7 & 2 & -5 \\ 3 & 0 & 0 & 0 \\ 8 & 3 & 1 & 7 \end{vmatrix} \quad 10. \begin{vmatrix} 1 & -2 & 5 & 2 \\ 0 & 0 & 3 & 0 \\ 2 & -4 & -3 & 5 \\ 2 & 0 & 3 & 5 \end{vmatrix}$$

$$11. \begin{vmatrix} 3 & 5 & -6 & 4 \\ 0 & -2 & 3 & -3 \\ 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 3 \end{vmatrix} \quad 12. \begin{vmatrix} 3 & 0 & 0 & 0 \\ 7 & -2 & 0 & 0 \\ 2 & 6 & 3 & 0 \\ 3 & -8 & 4 & -3 \end{vmatrix}$$

$$13. \begin{vmatrix} 4 & 0 & -7 & 3 & -5 \\ 0 & 0 & 2 & 0 & 0 \\ 7 & 3 & -6 & 4 & -8 \\ 5 & 0 & 5 & 2 & -3 \\ 0 & 0 & 9 & -1 & 2 \end{vmatrix}$$

$$14. \begin{vmatrix} 6 & 3 & 2 & 4 & 0 \\ 9 & 0 & -4 & 1 & 0 \\ 8 & -5 & 6 & 7 & 1 \\ 2 & 0 & 0 & 0 & 0 \\ 4 & 2 & 3 & 2 & 0 \end{vmatrix}$$

The expansion of a 3×3 determinant can be remembered by the following device. Write a second copy of the first two columns to the right of the matrix, and compute the determinant by multiplying entries on six diagonals:

Add the downward diagonal products and subtract the upward products. Use this method to compute the determinants in Exercises 15–18. **Warning:** This trick does not generalize in any reasonable way to 4×4 or larger matrices.

$$15. \begin{vmatrix} 1 & 0 & 4 \\ 2 & 3 & 2 \\ 0 & 5 & -2 \end{vmatrix} \quad 16. \begin{vmatrix} 0 & 3 & 1 \\ 4 & -5 & 0 \\ 3 & 4 & 1 \end{vmatrix}$$

$$17. \begin{vmatrix} 2 & -3 & 3 \\ 3 & 2 & 2 \\ 1 & 3 & -1 \end{vmatrix} \quad 18. \begin{vmatrix} 1 & 3 & 4 \\ 2 & 3 & 1 \\ 3 & 3 & 2 \end{vmatrix}$$

In Exercises 19–24, explore the effect of an elementary row operation on the determinant of a matrix. In each case, state the row operation and describe how it affects the determinant.

$$19. \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \begin{bmatrix} c & d \\ a & b \end{bmatrix}$$

$$20. \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \begin{bmatrix} a+kc & b+kd \\ c & d \end{bmatrix}$$

$$21. \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \begin{bmatrix} a & b \\ kc & kd \end{bmatrix}$$

$$22. \begin{bmatrix} 3 & 2 \\ 5 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 5+3k & 4+2k \end{bmatrix}$$

$$23. \begin{bmatrix} a & b & c \\ 3 & 2 & 1 \\ 4 & 5 & 6 \end{bmatrix}, \begin{bmatrix} a & b & c \\ 3 & 2 & 1 \\ 4 & 5 & 6 \end{bmatrix}$$

$$24. \begin{bmatrix} 1 & 0 & 1 \\ -3 & 4 & -4 \\ 2 & -3 & 1 \end{bmatrix}, \begin{bmatrix} k & 0 & k \\ -3 & 4 & -4 \\ 2 & -3 & 1 \end{bmatrix}$$

Compute the determinants of the elementary matrices given in Exercises 25–30. (See Section 2.2.)

$$25. \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & k & 1 \end{bmatrix} \quad 26. \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$27. \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ k & 0 & 1 \end{bmatrix} \quad 28. \begin{bmatrix} k & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$29. \begin{bmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad 30. \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Use Exercises 25–28 to answer the questions in Exercises 31 and 32. Give reasons for your answers.

31. What is the determinant of an elementary row replacement matrix?

32. What is the determinant of an elementary scaling matrix with k on the diagonal?

In Exercises 33–36, verify that $\det EA = (\det E)(\det A)$, where E is the elementary matrix shown and $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

$$33. \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \quad 34. \begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$$

$$35. \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad 36. \begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix}$$

$$37. \text{ Let } A = \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix}. \text{ Write } 5A. \text{ Is } \det 5A = 5 \det A?$$

38. Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and let k be a scalar. Find a formula that relates $\det kA$ to k and $\det A$.

In Exercises 39 and 40, A is an $n \times n$ matrix. Mark each statement True or False. Justify each answer.

39. a. An $n \times n$ determinant is defined by determinants of $(n-1) \times (n-1)$ submatrices.

b. The (i, j) -cofactor of a matrix A is the matrix A_{ij} obtained by deleting from A its i th row and j th column.

Section 3.2 : Properties of the Determinant

Chapter 3 : Determinants

Math 1554 Linear Algebra

"A problem isn't finished just because you've found the right answer."
- Yōko Ogawa

We have a method for computing determinants, but without some of the strategies we explore in this section, the algorithm can be very inefficient.

Topics and Objectives

Topics

We will cover these topics in this section.

- The relationships between row reductions, the invertibility of a matrix, and determinants.

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Apply properties of determinants (related to row reductions, transpose, and matrix products) to compute determinants.
2. Use determinants to determine whether a square matrix is invertible.

Topics and Objectives

Section 3.2 : Properties of the Determinant

Chapter 3 : Determinants
Math 1554 Linear Algebra

"A problem isn't finished just because you've found the right answer."
- Yoko Ogawa

We have a method for computing determinants, but without some of the strategies we explore in this section, the algorithm can be very inefficient.

Topics

We will cover these topics in this section.

- The relationships between row reductions, the invertibility of a matrix, and determinants.

Objectives

For the topics covered in this section, students are expected to be able to do the following.

- Apply properties of determinants (related to row reductions, transpose, and matrix products) to compute determinants.
- Use determinants to determine whether a square matrix is invertible.

5	9/18 - 9/22	2.3,2.4	WS2.2.2.3	2.5	WS2.4.2.5	2.8
6	9/25 - 9/29	2.9	WS2.8.2.9	3.1,3.2	WS3.1.3.2	3.3
7	10/2 - 10/6	4.9	WS3.3.4.9	5.1,5.2	WS5.1.5.2	5.2
8	10/9 - 10/13	Break	Break	Exam 2, Review	Cancelled	5.3

Section 3.2 Slide 182

Row Operations

- We saw how determinants are difficult or impossible to compute with a cofactor expansion for large N .
- Row operations give us a more efficient way to compute determinants.

Theorem: Row Operations and the Determinant

Let A be a square matrix.

- If a multiple of a row of A is added to another row to produce B , then $\det B = \det A$.
- If two rows are interchanged to produce B , then $\det B = -\det A$.
- If one row of A is multiplied by a scalar k to produce B , then $\det B = k \det A$.

Example 1 Compute $\begin{vmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{vmatrix} = \det A =$

$$\begin{pmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & -4 & 2 \\ 0 & 0 & -5 \\ 0 & 3 & 2 \end{pmatrix}$$

No change yet

new det = (-1) * old det

$$\begin{pmatrix} 1 & -4 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & -5 \end{pmatrix}$$

$$E_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\det E_3 = 1 \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 \cdot (1 \cdot 1) = 1$$

$$\sim \begin{pmatrix} 1 & -4 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{pmatrix} = U$$

$\frac{1}{3}R_2$

$$E_1 = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad "2R_1 + R_2 \rightarrow R_2"$$

~~"2R_1 - R_2 \rightarrow R_2"~~

$$\det E_1 = 1$$

$$E_2 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad "R_1 \leftrightarrow R_2"$$

$$\det(E_2) = -1$$

$$E_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad "1/3 R_2 \rightarrow R_2"$$

$$\det(E_3) = 1/3$$

new det = (-1/3) * old det

det U = 3

det A = 15

$$\begin{matrix} \frac{1}{3} & -1 & 1 & 1 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ E_4 & E_3 & E_2 & E_1 \end{matrix} \quad A = U$$

$$\det A = ? \quad \det U = 3$$

$$\det(E_4 E_3 E_2 E_1 A) = \det U$$

$$\downarrow \quad (1/3) \quad (-1) \quad (1) \quad (1) \quad ? \quad 3$$

$$\det(E_4) \det(E_3) \det(E_2) \det(E_1) \det A = \det U$$

Section 3.2 Slide 184

Section 3.2 Slide 188

Invertibility

Important practical implication: If A is reduced to echelon form, by P interchanges of rows and columns, then

$$|A| = \begin{cases} (-1)^r \times (\text{product of pivots}) & \text{when } A \text{ is invertible} \\ 0 & \text{when } A \text{ is singular.} \end{cases}$$

of REF.

Example 2 Compute the determinant

$$\begin{vmatrix} 0 & 1 & 2 & -1 \\ 2 & 5 & -7 & 3 \\ 0 & 3 & 6 & 2 \\ -2 & -5 & 4 & 2 \end{vmatrix}$$

IMT DLC
a. A invertible
P.
g. $\det A \neq 0$

THEOREM 3

Row Operations

Let A be a square matrix.

- a. If a multiple of one row of A is added to another row to produce a matrix B , then $\det B = \det A$.
- b. If two rows of A are interchanged to produce B , then $\det B = -\det A$.
- c. If one row of A is multiplied by k to produce B , then $\det B = k \cdot \det A$.

$$U = \begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

$\det U \neq 0$

$$U = \begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & 0 & \blacksquare \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\det U = 0$

FIGURE 1
Typical echelon forms of square matrices.

THEOREM 4

A square matrix A is invertible if and only if $\det A \neq 0$.

THEOREM 6

Multiplicative Property

If A and B are $n \times n$ matrices, then $\det AB = (\det A)(\det B)$.

Properties of the Determinant

For any square matrices A and B , we can show the following.

1. $\det A = \det A^T$.
2. A is invertible if and only if $\det A \neq 0$.
3. $\det(AB) = \det A \cdot \det B$.

Additional Example (if time permits)

Use a determinant to find all values of λ such that matrix C is not invertible.

$$C = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} - \lambda I_3$$

Compute $\det(C)$
for what λ 's is C not invertible?

$$C = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} - \begin{pmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{pmatrix}$$

$$C = \begin{pmatrix} 5-\lambda & 0 & 0 \\ 0 & -\lambda & 1 \\ 1 & 1 & -\lambda \end{pmatrix}$$

$$\det C = (5-\lambda) \begin{vmatrix} -\lambda & 1 \\ 1 & -\lambda \end{vmatrix} - 0 + 0$$

$$= (5-\lambda)(\lambda^2 - 1)$$

$$= -(\lambda - 5)(\lambda - 1)(\lambda + 1) = 0$$

$$\lambda = 5, 1, -1$$

Additional Example (if time permits)

Determine the value of

$$\det A = \det \begin{pmatrix} 0 & 2 & 0 & 4 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 3 \end{pmatrix}$$

3.2 EXERCISES

Each equation in Exercises 1–4 illustrates a property of determinants. State the property.

$$1. \begin{vmatrix} 0 & 5 & -2 \\ 1 & -3 & 6 \\ 4 & -1 & 8 \end{vmatrix} = - \begin{vmatrix} 1 & -3 & 6 \\ 0 & 5 & -2 \\ 4 & -1 & 8 \end{vmatrix}$$

$$2. \begin{vmatrix} 1 & 2 & 2 \\ 0 & 3 & -4 \\ 3 & 7 & 4 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 2 \\ 0 & 3 & -4 \\ 0 & 1 & -2 \end{vmatrix}$$

$$3. \begin{vmatrix} 3 & -6 & 9 \\ 3 & 5 & -5 \\ 1 & 3 & 3 \end{vmatrix} = 3 \begin{vmatrix} 1 & -2 & 3 \\ 3 & 5 & -5 \\ 1 & 3 & 3 \end{vmatrix}$$

$$4. \begin{vmatrix} 1 & 3 & -4 \\ 2 & 0 & -3 \\ 3 & -5 & 2 \end{vmatrix} = \begin{vmatrix} 1 & 3 & -4 \\ 0 & -6 & 5 \\ 3 & -5 & 2 \end{vmatrix}$$

Find the determinants in Exercises 5–10 by row reduction to echelon form.

$$5. \begin{vmatrix} 1 & 5 & -4 \\ -1 & -4 & 5 \\ -2 & -8 & 7 \end{vmatrix}$$

$$6. \begin{vmatrix} 3 & 3 & -3 \\ 3 & 4 & -4 \\ 2 & -3 & -5 \end{vmatrix}$$

$$7. \begin{vmatrix} 1 & 3 & 0 & 2 \\ -2 & -5 & 7 & 4 \\ 3 & 5 & 2 & 1 \\ 1 & -1 & 2 & -3 \end{vmatrix}$$

$$8. \begin{vmatrix} 1 & 3 & 2 & -4 \\ 0 & 1 & 2 & -5 \\ 2 & 7 & 6 & -3 \\ -3 & -10 & -7 & 2 \end{vmatrix}$$

$$9. \begin{vmatrix} 1 & -1 & -3 & 0 \\ 0 & 1 & 5 & 4 \\ -1 & 0 & 5 & 3 \\ 3 & -3 & -2 & 3 \end{vmatrix}$$

$$10. \begin{vmatrix} 1 & 3 & -1 & 0 & -2 \\ 0 & 2 & -4 & -2 & -6 \\ -2 & -6 & 2 & 3 & 10 \\ 1 & 5 & -6 & 2 & -3 \\ 0 & 2 & -4 & 5 & 9 \end{vmatrix}$$

Combine the methods of row reduction and cofactor expansion to compute the determinants in Exercises 11–14.

$$11. \begin{vmatrix} 3 & 4 & -3 & -1 \\ 3 & 0 & 1 & -3 \\ -6 & 0 & -4 & 3 \\ 6 & 8 & -4 & -1 \end{vmatrix}$$

$$12. \begin{vmatrix} -1 & 2 & 3 & 0 \\ 3 & 4 & 3 & 0 \\ 11 & 4 & 6 & 6 \\ 4 & 2 & 4 & 3 \end{vmatrix}$$

$$13. \begin{vmatrix} 2 & 5 & 4 & 1 \\ 4 & 7 & 6 & 2 \\ 6 & -2 & -4 & 0 \\ -6 & 7 & 7 & 0 \end{vmatrix}$$

$$14. \begin{vmatrix} 1 & 5 & 4 & 1 \\ 0 & -2 & -4 & 0 \\ 3 & 5 & 4 & 1 \\ -6 & 5 & 5 & 0 \end{vmatrix}$$

Find the determinants in Exercises 15–20, where

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = 7.$$

$$15. \begin{vmatrix} a & b & c \\ d & e & f \\ 3g & 3h & 3i \end{vmatrix}$$

$$16. \begin{vmatrix} a & b & c \\ 5d & 5e & 5f \\ g & h & i \end{vmatrix}$$

$$17. \begin{vmatrix} a+d & b+e & c+f \\ d & e & f \\ g & h & i \end{vmatrix}$$

$$18. \begin{vmatrix} d & e & f \\ a & b & c \\ g & h & i \end{vmatrix}$$

$$19. \begin{vmatrix} a & b & c \\ 2d+a & 2e+b & 2f+c \\ g & h & i \end{vmatrix}$$

$$20. \begin{vmatrix} a & b & c \\ d+3g & e+3h & f+3i \\ g & h & i \end{vmatrix}$$

In Exercises 21–23, use determinants to find out if the matrix is invertible.

$$21. \begin{bmatrix} 2 & 6 & 0 \\ 1 & 3 & 2 \\ 3 & 9 & 2 \end{bmatrix}$$

$$22. \begin{bmatrix} 5 & 1 & -1 \\ 1 & -3 & -2 \\ 0 & 5 & 3 \end{bmatrix}$$

$$23. \begin{bmatrix} 2 & 0 & 0 & 6 \\ 1 & -7 & -5 & 0 \\ 3 & 8 & 6 & 0 \\ 0 & 7 & 5 & 4 \end{bmatrix}$$

In Exercises 24–26, use determinants to decide if the set of vectors is linearly independent.

$$24. \begin{bmatrix} 4 \\ 6 \\ 2 \end{bmatrix}, \begin{bmatrix} -7 \\ 0 \\ 7 \end{bmatrix}, \begin{bmatrix} -3 \\ -5 \\ -2 \end{bmatrix}$$

$$25. \begin{bmatrix} 7 \\ -4 \\ -6 \end{bmatrix}, \begin{bmatrix} -8 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 0 \\ 7 \\ -5 \end{bmatrix}$$

$$26. \begin{bmatrix} 3 \\ 5 \\ -6 \\ 4 \end{bmatrix}, \begin{bmatrix} 2 \\ -6 \\ 0 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ -1 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ -2 \end{bmatrix}$$

In Exercises 27 and 28, A and B are $n \times n$ matrices. Mark each statement True or False. Justify each answer.

27. a. A row replacement operation does not affect the determinant of a matrix.

b. The determinant of A is the product of the pivots in any echelon form U of A , multiplied by $(-1)^r$, where r is the number of row interchanges made during row reduction from A to U .

c. If the columns of A are linearly dependent, then $\det A = 0$.

d. $\det(A+B) = \det A + \det B$.

28. a. If three row interchanges are made in succession, then the new determinant equals the old determinant.

b. The determinant of A is the product of the diagonal entries in A .

c. If $\det A$ is zero, then two rows or two columns are the same, or a row or a column is zero.

d. $\det A^{-1} = (-1) \det A$.

$$29. \text{ Compute } \det B^4, \text{ where } B = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{bmatrix}.$$

30. Use Theorem 3 (but not Theorem 4) to show that if two rows of a square matrix A are equal, then $\det A = 0$. The same is true for two columns. Why?

In Exercises 31–36, mention an appropriate theorem in your explanation.

31. Show that if A is invertible, then $\det A^{-1} = \frac{1}{\det A}$.

32. Suppose that A is a square matrix such that $\det A^3 = 0$. Explain why A cannot be invertible.

33. Let A and B be square matrices. Show that even though AB and BA may not be equal, it is always true that $\det AB = \det BA$.

34. Let A and P be square matrices, with P invertible. Show that $\det(PAP^{-1}) = \det A$.

35. Let U be a square matrix such that $U^T U = I$. Show that $\det U = \pm 1$.

36. Find a formula for $\det(rA)$ when A is an $n \times n$ matrix.

Verify that $\det AB = (\det A)(\det B)$ for the matrices in Exercises 37 and 38. (Do not use Theorem 6.)

$$37. A = \begin{bmatrix} 3 & 0 \\ 6 & 1 \end{bmatrix}, B = \begin{bmatrix} 2 & 0 \\ 5 & 4 \end{bmatrix}$$

$$38. A = \begin{bmatrix} 3 & 6 \\ -1 & -2 \end{bmatrix}, B = \begin{bmatrix} 4 & 3 \\ -1 & -3 \end{bmatrix}$$

39. Let A and B be 3×3 matrices, with $\det A = -3$ and $\det B = 4$. Use properties of determinants (in the text and in the exercises above) to compute:

- a. $\det AB$ b. $\det 5A$ c. $\det B^T$
d. $\det A^{-1}$ e. $\det A^3$

40. Let A and B be 4×4 matrices, with $\det A = -3$ and $\det B = -1$. Compute:

- a. $\det AB$ b. $\det B^5$ c. $\det 2A$
d. $\det A^T B A$ e. $\det B^{-1} A B$

41. Verify that $\det A = \det B + \det C$, where

$$A = \begin{bmatrix} a+e & b+f \\ c & d \end{bmatrix}, B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, C = \begin{bmatrix} e & f \\ c & d \end{bmatrix}$$

42. Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Show that $\det(A+B) = \det A + \det B$ if and only if $a+d=0$.

Section 3.3 : Volume, Linear Transformations

Chapter 3 : Determinants

Math 1554 Linear Algebra

NOTE: Cramer's rule and Adjoint of a matrix are NOT covered in Math 1554

Topics and Objectives

Topics

We will cover these topics in this section.

1. Relationships between area, volume, determinants, and linear transformations.

Objectives

For the topics covered in this section, students are expected to be able to do the following.

1. Use determinants to compute the area of a parallelogram, or the volume of a parallelepiped, possibly under a given linear transformation.

Students are not expected to be familiar with Cramer's rule.

Section 3.3 : Volume, Linear Transformations

Topics and Objectives

5	9/18 - 9/22	2.3,2.4	WS2.2.2.3	2.5	WS2.4.2.5	2.8
6	9/25 - 9/29	2.9	WS2.8.2.9	3.1,3.2	WS3.1.3.2	3.3
7	10/2 - 10/6	4.9	WS3.3.4.9	5.1,5.2	WS5.1.5.2	5.2
8	10/9 - 10/13	Break	Break	Exam 2, Review	Cancelled	5.3

Chapter 3 : Determinants
Math 1554 Linear Algebra

Topics
We will cover these topics in this section.
1. Relationships between area, volume, determinants, and linear transformations.

Objectives
For the topics covered in this section, students are expected to be able to do the following:
1. Use determinants to compute the area of a parallelogram, or the volume of a parallelepiped, possibly under a given linear transformation.



Students are not expected to be familiar with Cramer's rule.

NOTE: Cramers rule and Adjoint of a matrix are NOT covered in Math 1554

Supplementary FREE textbook

<https://textbooks.math.gatech.edu/ila/>

NOTE: Cramers rule and Adjoint of a matrix are NOT covered in Math 1554



Parallelograms and Parallelepipeds


The determinant computes the volume of the following kind of geometric object.

Definition. The *parallelepiped* determined by n vectors v_1, v_2, \dots, v_n in \mathbb{R}^n is the subset

$$P = \{a_1x_1 + a_2x_2 + \dots + a_nx_n \mid 0 \leq a_1, a_2, \dots, a_n \leq 1\}.$$

In other words, a parallelepiped is the set of all linear combinations of n vectors with coefficients in $[0, 1]$. We can draw parallelepipeds using the parallelogram law for vector addition.

Example (The unit cube). The parallelepiped determined by the standard coordinate vectors e_1, e_2, \dots, e_n is the unit n -dimensional cube.



Topics and Objectives

Topics

We will cover these topics in this section.

1. Relationships between area, volume, determinants, and linear transformations.

Objectives

For the topics covered in this section, students are expected to be able to do the following.


1. Use determinants to compute the area of a parallelogram, or the volume of a parallelepiped, possibly under a given linear transformation.

Students are not expected to be familiar with Cramer's rule.

Section 3.3 Slide 192

Determinants, Area and Volume

In \mathbb{R}^2 , determinants give us the area of a parallelogram.



$$A = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$$

$$\det(A) = ad - bc$$

Section 3.3 Slide 193






FIGURE 3
Volume = $|abc|$.

$$\det \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} = abc$$


Example (Parallelograms). When $n = 2$, a parallelepiped is just a parallelogram in \mathbb{R}^2 . Note that the edges come in parallel pairs.



Example. When $n = 3$, a parallelepiped is a kind of a skewed cube. Note that the faces come in parallel pairs.



When does a parallelepiped have zero volume? This can happen only if the parallelepiped is flat, i.e., it is squashed into a lower dimension.




This means exactly that $\{v_1, v_2, \dots, v_n\}$ is *linearly dependent*, which by this [corollary in Section 4.1](#) means that the matrix with rows v_1, v_2, \dots, v_n has determinant zero. To summarize:

Key Observation. The parallelepiped defined by v_1, v_2, \dots, v_n has zero volume if and only if the matrix with rows v_1, v_2, \dots, v_n has zero determinant.


Example (Area of a triangle). \blacktriangleleft

Find the area of the triangle with vertices $(-1, -2)$, $(2, -1)$, $(1, 3)$.



Solution

Doubling a triangle makes a parallelogram. We choose two of its sides to be the rows of a matrix.



$$\det \begin{bmatrix} 2 & 3 \\ 5 & 1 \end{bmatrix} = 2 \cdot 1 - 3 \cdot 5$$

$$= 2 - 15$$

$$= -13/2$$

$$\det \begin{bmatrix} 3 & 2 \\ 1 & 5 \end{bmatrix} = 15 - 2 = 13$$

So triangle area is

$$13/2$$

Determinants as Area, or Volume

Theorem

The volume of the parallelepiped spanned by the columns of an $n \times n$ matrix A is $|\det A|$.

Key Geometric Fact (which works in any dimension). The area of the parallelogram spanned by two vectors \vec{a}, \vec{b} is equal to the area spanned by $\vec{a}, c\vec{a} + \vec{b}$, for any scalar c .




FIGURE 2 Two parallelograms of equal area.

Section 3.3 Slide 194

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Any 3×3 matrix A can be transformed into a diagonal matrix using column operations that do not change $|\det(A)|$.





FIGURE 4 Two parallelepipeds of equal volume.

Section 3.3



Example 1

Calculate the area of the parallelogram determined by the points $(-2, -2), (0, 3), (4, -1), (6, 4)$.




FIGURE 5 Translating a parallelogram does not change its area.

$$\begin{aligned} |\det(\vec{v}_1, \vec{v}_2)| &= \left| \det \begin{bmatrix} 2 & 4 \\ 5 & 1 \end{bmatrix} \right| \begin{matrix} \leftarrow 4 - (-2) \\ \leftarrow 1 - (-2) \end{matrix} \\ &= |2 - 20| = |-18| \\ &= \boxed{18} \end{aligned}$$

Section 3.3 Slide 196


Linear Transformations

Theorem

If $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$, and S is some parallelogram in \mathbb{R}^n , then

$$\text{volume}(T_A(S)) = |\det(A)| \cdot \text{volume}(S)$$

An example that applies this theorem is given in this week's worksheets.



Section 3.3 Slide 197



THEOREM 10

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation determined by a 2×2 matrix A . If S is a parallelogram in \mathbb{R}^2 , then

$$\{\text{area of } T(S)\} = |\det A| \cdot \{\text{area of } S\} \quad (5)$$

If T is determined by a 3×3 matrix A , and if S is a parallelepiped in \mathbb{R}^3 , then

$$\{\text{volume of } T(S)\} = |\det A| \cdot \{\text{volume of } S\} \quad (6)$$



$$ax^2 + bx + cy + dy = k$$

Example (Area of an ellipse). ^

Find the area of the interior E of the ellipse defined by the equation

$$\left(\frac{2x-y}{2}\right)^2 + \left(\frac{y+3x}{3}\right)^2 = 1.$$

If $a^2 + b^2 = 1$

then $\left(\frac{2x-y}{2}\right)^2 + \left(\frac{y+3x}{3}\right)^2 = 1$

$$a = \frac{2x-y}{2} \quad \begin{cases} x - \frac{1}{2}y = a \\ x + \frac{1}{3}y = b \end{cases}$$

So Area ellipse

area circle \downarrow
 $= \det A^{-1} \cdot \pi \cdot 1^2$

J ("circle") = "ellipse"
 $T(x) = A^{-1}x$

$\pi \cdot 6/5$

<https://www.geogebra.org/calculator/mkxegfjy>

$$A^{-1}A \begin{pmatrix} x \\ y \end{pmatrix} = A \begin{pmatrix} a \\ b \end{pmatrix}$$

$$A = \begin{pmatrix} 1 & -1/2 \\ 1 & 1/3 \end{pmatrix}$$


$[x] = A^{-1}[a]$
 ellipse $\frac{6}{5}\pi$
 circle

$$\det A^{-1} = \frac{6}{5}$$

$$\det A = \frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6}$$

$$\det(A^{-1}A) = \det I$$

$$\det(A^{-1}) \det A = 1$$



Example (Area of an ellipse). ^

Find the area of the interior E of the ellipse defined by the equation

$$\left(\frac{2x-y}{2}\right)^2 + \left(\frac{y+3x}{3}\right)^2 = 1.$$

Vol (Parallelepiped) = |det A|?

① $\det(AB) = \det A \cdot \det B.$

② use elementary matrices.


$$A = E_k E_{k-1} \dots E_2 E_1$$

§4.9 MARKOV CHAINS Preview

Example 1

- A small town has two libraries, A and B .
- After 1 month, among the books checked out of A ,
 - 80% returned to A
 - 20% returned to B
- After 1 month, among the books checked out of B ,
 - 30% returned to A
 - 70% returned to B

If both libraries have 1000 books today, how many books does each library have after 1 month? After one year? After n months? A place to simulate this is <http://setosa.io/markov/index.html>



Example 1 Continued

The books are equally divided by between the two branches, denoted by $\vec{x}_0 = \begin{bmatrix} .5 \\ .5 \end{bmatrix}$. What is the distribution after 1 month, call it \vec{x}_1 ? After two months?

After k months, the distribution is \vec{x}_k , which is what in terms of \vec{x}_0 ?

$$\text{Nul}(A - I)$$

Section 4.9 Slide 200

From A: $(.80)1000 + (.20)1000 = 800 + 200 = 1000$

From B: $(.30)1000 + (.70)1000 = 300 + 700 = 1000$

$$1000 \begin{bmatrix} .80 \\ .20 \end{bmatrix} + 1000 \begin{bmatrix} .30 \\ .70 \end{bmatrix} = \begin{bmatrix} 1100 \\ 900 \end{bmatrix}$$

$$\begin{bmatrix} .80 \\ .20 \end{bmatrix} \cdot 300 + \begin{bmatrix} .30 \\ .70 \end{bmatrix} \cdot 700 = \begin{bmatrix} 1100 \\ 900 \end{bmatrix}$$

$$(.80)1100 + (.20)900 = 1150$$

$$(.20)1100 + (.80)900 = 850$$

$$1100 \begin{bmatrix} .60 \\ .20 \end{bmatrix} + 900 \begin{bmatrix} .30 \\ .70 \end{bmatrix} = \begin{bmatrix} 1150 \\ 850 \end{bmatrix}$$

@A:

Section 4.9 Slide 201

$$A \vec{x}_0 = \vec{x}_1$$

$$A \vec{x}_1 = \vec{x}_2$$

$$A \vec{x}_2 = \vec{x}_3$$

$$A^n \vec{x}_0 = \vec{x}_n$$

In Exercises 19–22, find the area of the parallelogram whose vertices are listed.

19. $(0, 0), (5, 2), (6, 4), (11, 6)$
20. $(0, 0), (-2, 4), (4, -5), (2, -1)$
21. $(-2, 0), (0, 3), (1, 3), (-1, 0)$
22. $(0, -2), (5, -2), (-3, 1), (2, 1)$
23. Find the volume of the parallelepiped with one vertex at the origin and adjacent vertices at $(1, 0, -3), (1, 2, 4),$ and $(5, 1, 0)$.
24. Find the volume of the parallelepiped with one vertex at the origin and adjacent vertices at $(1, 3, 0), (-2, 0, 2),$ and $(-1, 3, -1)$.
25. Use the concept of volume to explain why the determinant of a 3×3 matrix A is zero if and only if A is not invertible. Do not appeal to Theorem 4 in Section 3.2. [Hint: Think about the columns of A .]
26. Let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation, and let \mathbf{p} be a vector and S a set in \mathbb{R}^m . Show that the image of $\mathbf{p} + S$ under T is the translated set $T(\mathbf{p}) + T(S)$ in \mathbb{R}^n .
27. Let S be the parallelogram determined by the vectors $\mathbf{b}_1 = \begin{bmatrix} -2 \\ 3 \end{bmatrix}$ and $\mathbf{b}_2 = \begin{bmatrix} -2 \\ 5 \end{bmatrix}$, and let $A = \begin{bmatrix} 6 & -3 \\ -3 & 2 \end{bmatrix}$. Compute the area of the image of S under the mapping $\mathbf{x} \mapsto A\mathbf{x}$.

28. Repeat Exercise 27 with $\mathbf{b}_1 = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$, $\mathbf{b}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, and $A = \begin{bmatrix} 5 & 2 \\ 1 & 1 \end{bmatrix}$.

29. Find a formula for the area of the triangle whose vertices are $\mathbf{0}, \mathbf{v}_1,$ and \mathbf{v}_2 in \mathbb{R}^2 .
30. Let R be the triangle with vertices at $(x_1, y_1), (x_2, y_2),$ and (x_3, y_3) . Show that

$$\{\text{area of triangle}\} = \frac{1}{2} \det \begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$$

[Hint: Translate R to the origin by subtracting one of the vertices, and use Exercise 29.]

31. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation determined by the matrix $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$, where $a, b,$ and c are