

# Circumference of 3-connected claw-free graphs and large Eulerian subgraphs of 3-edge-connected graphs

Mark Bilinski\*

Bill Jackson†

Jie Ma ‡

Xingxing Yu §

## Abstract

The circumference of a graph is the length of its longest cycles. Results of Jackson, and Jackson and Wormald, imply that the circumference of a 3-connected cubic  $n$ -vertex graph is  $\Omega(n^{0.694})$ , and the circumference of a 3-connected claw-free graph is  $\Omega(n^{0.121})$ . We generalise and improve the first result by showing that every 3-edge-connected graph with  $m$  edges has an Eulerian subgraph with  $\Omega(m^{0.753})$  edges. We use this result together with the Ryjáček closure operation to improve the lower bound on the circumference of a 3-connected claw-free graph to  $\Omega(n^{0.753})$ . Our proofs imply polynomial time algorithms for finding large Eulerian subgraphs of 3-edge-connected graphs and long cycles in 3-connected claw-free graphs.

---

\*School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA. Partially supported by NSF VIGRE Grant

†School of Mathematical Sciences, Queen Mary University of London, London E1 4NS, England

‡School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA.

§School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA. Partially supported by an NSA grant and NSFC Project 10628102

# 1 Introduction

Motivated by the Four Color Problem, Tait [43] conjectured in 1880 that every 3-connected cubic planar graph contains a Hamilton cycle. His conjecture remained open until a counterexample was constructed by Tutte [45] in 1946. There has since been much interest and extensive research concerning longest cycles in (special families of) graphs. We use  $|G|$  to denote the number of vertices in a graph  $G$  and refer to the length of a longest cycle in  $G$  as the *circumference* of  $G$ . We will be concerned with bounds on the circumference of 3-connected graphs which are either cubic or claw-free.

Barnette [3] showed that every 3-connected cubic  $n$ -vertex graph has circumference  $\Omega(\log n)$ . Bondy and Simonovits [9] improved this lower bound to  $\exp(\Omega(\sqrt{\log n}))$ , and conjectured that it can be improved further to  $\Omega(n^c)$  for some constant  $0 < c < 1$ . This conjecture was established by Jackson [29], with  $c = \log_2(1 + \sqrt{5}) - 1 \approx 0.694$ . A construction given by Bondy and Simonovits in [9] gives an infinite family of 3-connected cubic graphs with circumference  $\Theta(n^{\log_9 8})$ , where  $\log_9 8 \approx 0.946$ . Our first theorem improves the exponent in the lower bound on circumference given in [29], and also generalises the result to graphs which are not necessarily cubic. We use  $K_2^3$  to denote the graph with two vertices joined by three parallel edges.

**Theorem 1.1** *Let  $G$  be a 3-edge-connected graph,  $e, f \in E(G)$ , and assume  $G \neq K_2^3$ . Then  $G$  contains an Eulerian subgraph  $H$  such that  $e, f \in E(H)$  and  $|E(H)| \geq (|E(G)|/6)^\alpha + 2$ , where  $\alpha \approx 0.753$  is the real root of  $4^{1/x} - 3^{1/x} = 2$ .*

Given graphs  $G, H$ , we say that  $G$  is  $H$ -free if  $G$  has no induced subgraph isomorphic to  $H$ . In the special case when  $H = K_{1,3}$  we say that  $G$  is *claw-free*. Jackson and Wormald [30] proved a general lower bound on the circumference of 3-connected  $K_{1,d}$ -free graphs, which reduces to  $\frac{1}{2}|G|^c$ , where  $c = \log_{150} 2 \approx 0.121$ , when  $G$  is claw-free. We will obtain the following stronger result.

**Theorem 1.2** *If  $G$  is a 3-connected claw-free graph, then the circumference of  $G$  is at least  $(|G|/12)^\alpha + 2$ , where  $\alpha \approx 0.753$  is the real root of  $4^{1/x} - 3^{1/x} = 2$ .*

Note that if  $G$  is a cubic graph then blowing up each vertex of  $G$  to a triangle in an obvious way we obtain a claw-free cubic graph  $H$ ; and it is easy to see that the circumference of  $G$  is  $\Theta(|G|^c)$  if and only if the circumference of  $H$  is  $\Theta(|H|^c)$ . Thus the above mentioned construction of Bondy and Simonovits implies that the exponent  $\alpha$  in Theorem 1.2 cannot exceed  $\log_9 8$ .

We prove Theorem 1.2 by reducing the problem to line graphs using the closure result of Ryjáček [40]. For  $x$  a vertex in a graph  $G$  we use  $N_G(x)$  (or simply  $N(x)$  if there is no confusion) to denote the neighborhood of  $x$ ; and for each  $S \subseteq V(G)$  we use  $G[S]$  to denote the subgraph of  $G$  induced by  $S$ . Let  $G_0, \dots, G_k$  be a maximal sequence of graphs such that  $G_0 = G$  and for each  $1 \leq i \leq k$ ,  $G_i$  is obtained from  $G_{i-1}$  by taking some  $x \in V(G)$  for which  $G_{i-1}[N_{G_{i-1}}(x)]$  is connected and adding edges between all pairs of nonadjacent vertices in  $N_{G_{i-1}}(x)$ . Then  $G_k$  is said to be a *Ryjáček closure* of  $G$ .

**Theorem 1.3** [40] *The Ryjáček closure of a claw-free simple graph  $G$  is uniquely determined, and is equal to the line graph  $L(H)$  of a triangle-free simple graph  $H$ . Furthermore, for every cycle  $C'$  of  $L(H)$  there exists a cycle  $C$  of  $G$  with  $V(C') \subseteq V(C)$ .*

The final conclusion of this theorem is a slightly stronger statement than that given by Ryjáček in [40, Theorem 3] (that the circumferences of  $G$  and  $L(H)$  are the same), but it follows from his proof, see [14, Proposition G] and [13, Lemma 8].

It is clear that in a graph  $H$  any Eulerian subgraph with  $m$  edges gives rise to a cycle with  $m$  vertices in  $L(H)$ . In addition we will see that  $L(H)$  is 3-connected if and only if the removal of all degree one vertices from  $H$  results in a graph obtained from a 3-edge-connected graph by subdividing each edge at most once. Thus Theorem 1.2 will follow from Theorem 1.3 and an edge-weighted version of Theorem 1.1.

Bounds on the circumference of order  $|G|^c$  have also been obtained for other families of 3-connected graphs  $G$ . For graphs embedded on a fixed surface, Chen and Yu [17] proved that every 3-connected  $n$ -vertex graph embeddable in the torus or Klein bottle has circumference at least  $n^{\log_3 2}$ , establishing a conjecture of Moon and Moser [38] and Grünbaum and Walther [26]. This was generalized in [41] to locally planar graphs on orientable surfaces. Infinite families of 3-connected cubic planar graphs  $G$  with circumference  $\Theta(|G|^c)$  have been constructed (for various constants  $0 < c < 1$ ), see for example [25, 26, 47, 48].

For graphs of bounded maximum degree, Jackson and Wormald [30] proved that every 3-connected  $n$ -vertex graph with maximum degree at most  $d$  has circumference  $\Omega(n^{\log_b 2})$ , with  $b = 6d^2$ . This result was improved to  $b = 2(d - 1)^2 + 1$  by Chen, Xu and Yu [16], and further improved to  $b = 4d + 1$  by Chen, Gao, Zang and Yu [15]. When  $d \geq 4$ , Jackson and Wormald conjecture that the correct value for  $b$  is  $d - 1$ , and construct an infinite family of 3-connected  $n$ -vertex graphs with maximum degree  $d$  and circumference  $\Theta(n^{\log_{d-1} 2})$  in [30].

One may also consider families of graphs of connectivity other than three. Bounds on the circumference of families of 2-connected  $n$ -vertex graphs tend to be of order  $\log n$ . In particular Bondy and Entringer [8] showed that every 2-connected graph with maximum degree at most  $d$  has circumference at least  $\log_{d-1} n$ , and constructs an infinite family of such graphs with circumferences of the same order of magnitude. Broersma et al [12] showed that the circumference of a 2-connected claw-free  $n$ -vertex graph is also  $\Omega(\log n)$ . (Note that there can be no analogous result for 2-connected graphs embeddable on a fixed surface since  $K_{2,n-2}$  is 2-connected and planar, and has circumference four.)

On the other hand, bounds on the circumference of families of  $n$ -vertex graphs of connectivity greater than three may be of order  $n$ . Bondy, see [29, Conjecture 1], conjectured that if  $G$  is a 3-connected cubic graph and every 3-edge-cut of  $G$  is trivial, then  $G$  has circumference  $\Omega(n)$ . A stronger conjecture due to Fleischner, see [29, Conjecture 2], is that every such graph  $G$  has a cycle  $C$  such that  $G - C$  is an independent set of vertices. Both conjectures are true for planar cubic graphs by Tutte's bridge theorem [46]. Fleischner and Jackson [22] showed that Fleischner's conjecture is equivalent to a conjecture of Thomassen [44] that every 4-connected line graph is Hamiltonian. Ryjáček [40] used Theorem 1.3 to show that Thomassen's conjecture is in turn equivalent to the conjecture of Mathews and Sumner [36] that every 4-connected claw-free graph is Hamiltonian. Zhang [49] has verified Thomassen's conjecture for the special case of 7-connected line graphs. This result was extended to 7-connected claw-free graphs by Ryjáček in [40].

An outline of the paper is as follows. Section 2 contains some preliminary results. We introduce a reduction technique called 'edge-splitting' in Subsection 2.1 and characterize when it can be used to split away two edges from a vertex in such a way that 3-edge-connectivity is preserved. In Subsection 2.2, we characterize when a 3-edge-connected graph has an Eulerian subgraph which contains two given edges and four given vertices. In Subsection 2.3, we prove

some inequalities based on the concavity of the function  $n \rightarrow n^c$  when  $0 < c < 1$  which we will use in our induction. We prove the aforementioned edge-weighted version of Theorem 1.1 in Section 3 by applying the edge-splitting lemmas to reduce to the case when each of the endvertices of  $e$  and  $f$  has degree three, and then extending the proof technique for cubic graphs given in [29]. Theorem 1.2 is derived in Section 4. Our proofs of Theorems 1.1 and 1.2 are constructive and give rise to polynomial algorithms. These will be outlined in Section 5.

## 2 Definitions and preliminary results

Unless specified otherwise all graphs considered may contain loops and multiple edges. We will refer to graphs without loops and multiple edges as *simple graphs*. For any edge  $e$  in a graph  $G$ , we use  $V(e)$  to denote the set of vertices of  $G$  that are incident with  $e$ . For  $S \subseteq E(G)$  we use  $G - S$  to denote the graph obtained from  $G$  by deleting  $S$ . For  $H$  and  $L$  subgraphs of  $G$ , we use  $H - L$  to denote the graph obtained from  $H$  by deleting  $V(H) \cap V(L)$  and all edges of  $H$  incident with vertices in  $V(H) \cap V(L)$ . If  $L$  consists of one vertex, say  $v$ , then we also write  $H - v$  for  $H - L$ .

### 2.1 Edge splitting

Let  $G$  be a graph,  $v \in V(G)$ , and  $e, f$  be distinct edges of  $G$  with  $V(e) = \{u, v\}$  and  $V(f) = \{v, w\}$ . When  $d(v) = 2$ , the operation of *suppressing  $v$  in  $G$*  deletes  $v$  (and hence also  $e, f$ ) and adds a new edge between  $u$  and  $w$  (which may be a loop if  $u = w$ ). When  $d(v) \geq 4$  the operation of *splitting  $e, f$  at  $v$*  deletes  $e, f$  from  $G$ , adds a new edge between  $u$  and  $w$ , and suppresses  $v$  if  $v$  has degree 2 in  $G - \{e, f\}$ . We use  $G_v^{e,f}$  to denote the graph obtained from  $G$  by splitting  $e, f$  at  $v$ . Note that if  $e$  is a loop at  $v$  then  $G_v^{e,f}$  is isomorphic to  $G - e$  when  $d(v) > 4$ , and to the graph obtained from  $G - e$  by suppressing  $v$  when  $d(v) = 4$ . When  $G$  is  $k$ -edge-connected, we say that  $e, f$  form a  *$k$ -splittable pair at  $v$*  if  $G_v^{e,f}$  is also  $k$ -edge-connected. (Note that loops have no effect on edge-connectivity so a pair containing a loop will always be  $k$ -splittable.) If there is no confusion, we will simply say that  $e, f$  is a *splittable pair at  $v$* . We need the following consequence of a more general result of Frank (Theorem B, [23]).

**Lemma 2.1** *Let  $G$  be a 3-edge-connected graph and  $v \in V(G)$  such that  $d(v) \geq 4$ . If  $d(v)$  is even then each edge incident with  $v$  belongs to a splittable pair at  $v$ . If  $d(v)$  is odd then there is at most one edge incident with  $v$  that does not belong to any splittable pair at  $v$ .*

For our purpose, we also need to describe the structure when an edge is not contained in any splittable pair. This structure is illustrated in Figure 1. To describe it more precisely we need some more notation. Given a graph  $G$  and disjoint subsets  $X, Y$  of  $V(G)$ , we use  $E(X, Y)$  to denote the set, and  $\delta(X, Y)$  the number, of edges of  $G$  incident with both  $X$  and  $Y$ . When  $X = \{x\}$  or  $Y = \{y\}$ , we write  $\delta(x, Y)$  or  $\delta(X, y)$ . We also put  $\delta(X) = \delta(X, V(G) - X)$ . We write  $\delta_G(X)$  when the underlying graph  $G$  is not clear from the context.

The lemma below is similar to a result for local edge-connectivity due to Szigeti (Theorem 1.6, [42]). We will need the  $k = 3$  case (see Figure 1) but we state it for general  $k$  as it may be of independent interest.

**Lemma 2.2** *Let  $G$  be a  $k$ -edge-connected graph ( $k \geq 3$ ) and  $e \in E(G)$  with  $V(e) = \{u, v\}$ . Suppose that  $d(v) \geq k + 2$ , and  $e$  belongs to no splittable pair at  $v$ . Then  $k$  is odd,  $d(v) = k + 2$ ,*




Figure 1:  $k = 3$  and the edge  $e$  belongs to no splittable pair at  $v$ .

and there exists a partition  $Y_0, Y_1, Y_2$  of  $V(G) - \{v\}$  such that  $u \in Y_0$ ,  $\delta(v, Y_0) = 1$ ,  $\delta(v, Y_1) = \delta(v, Y_2) = (k + 1)/2$ ,  $\delta(Y_0, Y_1) = \delta(Y_0, Y_2) = (k - 1)/2$ , and  $\delta(Y_1, Y_2) = 0$ .

*Proof.* Since  $e$  is contained in no splittable pair,  $v$  is incident with no loops and there exists a family of sets  $\mathcal{F} = \{X_1, \dots, X_t\}$  such that  $N(v) \subseteq \bigcup_{i=1}^t X_i$  and, for  $1 \leq i \leq t$ ,  $u \in X_i \subseteq V - \{v\}$  and  $\delta(X_i) \leq k + 1$ . We choose  $\mathcal{F}$  such that

- (1)  $t$  is minimum, and
- (2) subject to (1),  $\sum_{i=1}^t |X_i|$  is maximum.

Since  $d(v) \geq k + 2$  and  $v$  is not incident with any loop, we have  $t \geq 2$ . Let  $Y_0 = X_1 \cap X_2$ ,  $Y_1 = X_1 - X_2$  and  $Y_2 = X_2 - X_1$ . By (1),  $Y_i \neq \emptyset$  for  $i = 1, 2$ . Note that  $\delta(Y_0) \geq k$  since  $u \in Y_0$  and  $G$  is  $k$ -edge-connected. Also  $\delta(X_1 \cup X_2) \geq k + 2$ , for otherwise  $\delta(X_1 \cup X_2) \leq k + 1$  and  $(\mathcal{F} - \{X_1, X_2\}) \cup \{X_1 \cup X_2\}$  contradicts the choice of  $\mathcal{F}$  (via (1)). So

$$(k + 1) + (k + 1) \geq \delta(X_1) + \delta(X_2) = \delta(Y_0) + \delta(X_1 \cup X_2) + 2\delta(Y_1, Y_2) \geq k + (k + 2).$$

Therefore, equality must hold throughout; so  $\delta(X_1) = \delta(X_2) = k + 1$ ,  $\delta(Y_0) = k$ ,  $\delta(Y_1, Y_2) = 0$ , and  $\delta(X_1 \cup X_2) = k + 2$ .

Since  $u \in Y_0$  and  $v \in V(G) - (X_1 \cup X_2)$ ,  $\delta(Y_0, V(G) - (X_1 \cup X_2)) \geq 1$ . Because  $G$  is  $k$ -edge-connected,  $\delta(Y_i) \geq k$  for  $i = 1, 2$ ; and hence

$$(k + 1) + (k + 1) = \delta(X_1) + \delta(X_2) = \delta(Y_1) + \delta(Y_2) + 2\delta(Y_0, V(G) - (X_1 \cup X_2)) \geq k + k + 2.$$

Equality holds throughout; so  $\delta(Y_1) = \delta(Y_2) = k$  and  $\delta(v, Y_0) = \delta(Y_0, V(G) - (X_1 \cup X_2)) = 1$ .

Since  $G$  is  $k$ -edge-connected and  $\delta(X_1) = k + 1$ ,  $G[X_1]$  is  $\lceil (k - 1)/2 \rceil$ -edge-connected. Hence  $\delta(Y_0, Y_1) \geq \lceil (k - 1)/2 \rceil$ . Similarly,  $\delta(Y_0, Y_2) \geq \lceil (k - 1)/2 \rceil$ . Because  $\delta(Y_0) = k$ ,  $v \notin X_1 \cup X_2$ , and  $\delta(v, Y_0) = 1$ , we must have  $\delta(Y_0, Y_1) = \delta(Y_0, Y_2) = (k - 1)/2$ . In particular,  $k$  is odd.

We may assume  $t \geq 3$ . For, suppose  $t = 2$ . Then  $N(v) \subseteq Y_0 \cup Y_1 \cup Y_2$ . Since  $\delta(Y_1, Y_2) = 0$ ,  $\delta(Y_1) = \delta(Y_2) = k$ , and  $\delta(Y_0, Y_1) = \delta(Y_0, Y_2) = (k - 1)/2$ , we have  $\delta(v, Y_1) = \delta(v, Y_2) = (k + 1)/2$ . Hence  $d(v) = k + 2$ . Therefore, there are no edges of  $G$  leaving  $Y_0 \cup Y_1 \cup Y_2 \cup \{v\}$ ; so  $\{Y_0, Y_1, Y_2\}$  is a partition of  $V(G) - \{v\}$ , and the assertion of the lemma holds.

Suppose  $Y_0 \not\subseteq X_3$ . Note that  $\delta(X_3 \cup Y_0) \geq k + 2$  as otherwise  $(\mathcal{F} - \{X_3\}) \cup \{X_3 \cup Y_0\}$  contradicts the choice of  $\mathcal{F}$  (via (2)). Since  $u \in X_3 \cap Y_0$  and  $G$  is  $k$ -edge-connected,  $\delta(X_3 \cap Y_0) \geq k$ . Therefore, we have the following contradiction

$$(k + 1) + k \geq \delta(X_3) + \delta(Y_0) \geq \delta(X_3 \cup Y_0) + \delta(X_3 \cap Y_0) \geq (k + 2) + k.$$

So  $Y_0 \subseteq X_3$ , i.e.,  $X_1 \cap X_2 \subseteq X_3$ . Hence by symmetry among  $X_1, X_2, X_3$ , we also have  $X_2 \cap X_3 \subseteq X_1$  and  $X_1 \cap X_3 \subseteq X_2$ . So  $X_1 \cap X_2 = X_1 \cap X_3 = X_2 \cap X_3 = Y_0$  and  $\delta(Y_0, X_1 - Y_0) = \delta(Y_0, X_2 - Y_0) = \delta(Y_0, X_3 - Y_0) = (k - 1)/2$ . This is impossible since we also have  $\delta(v, Y_0) = 1$  and  $\delta(Y_0) = k$ . ■

We also need to know when an edge is contained in a unique splittable pair at a vertex of degree four in a 3-edge-connected graph, see Figure 2. This follows from a more general result of Jordán [32, Theorem 3.6].




Figure 2: The edge  $e$  belongs to a unique splittable pair at  $v$ .

**Lemma 2.3** *Let  $G$  be a 3-edge-connected graph and  $e, f \in E(G)$  with  $V(e) = \{u, v\}$  and  $V(f) = \{v, w\}$ . Suppose that  $d(v) = 4$  and that  $e, f$  is the unique splittable pair at  $v$  which contains  $e$ . Then there exists a partition  $Y_0, Y_1, Y_2, Y_3$  of  $V(G) - \{v\}$  such that  $u \in Y_0$ ,  $w \in Y_2$ ,  $\delta(v, Y_i) = 1$  for all  $0 \leq i \leq 3$ ,  $\delta(Y_0, Y_1) = \delta(Y_1, Y_2) = \delta(Y_2, Y_3) = \delta(Y_3, Y_0) = 1$ , and  $\delta(Y_0, Y_2) = \delta(Y_1, Y_3) = 0$ .*

## 2.2 Cyclability

Let  $G$  be a graph and  $e \in E(G)$  with  $V(e) = \{u, v\}$ . Then the graph  $G/e$  obtained from  $G$  by contracting  $e$  to a single vertex  $z$  (where  $z \notin V(G)$ ) is the graph obtained from  $G - \{u, v\}$  by adding the new vertex  $z$  and replacing each edge  $f$  in  $G - e$  with at least one end in  $\{u, v\}$  by an edge in which the corresponding end vertex/vertices are equal to  $z$ . We denote the edge of  $G/e$  corresponding to  $f$  by the same label  $f$ . Note that an edge  $f$  of  $G - e$  with  $V(f) = \{u, v\}$  will be replaced by a loop at  $z$  in  $G/e$ . More generally, if  $H$  is a subgraph of  $G$ , then graph  $G/H$  obtained from  $G$  by contracting  $H$  to a single vertex  $z$  (where  $z \notin V(G)$ ) is the graph obtained from  $G - H$  by adding the new vertex  $z$  and replacing each edge  $f$  in  $G - E(H)$  with at least one end in  $V(H)$  by an edge in which the corresponding end vertex/vertices are equal to  $z$ . We again denote the edge of  $G/e$  corresponding to  $f$  by the same label  $f$ . Note that: contracting a subgraph cannot reduce the edge-connectivity of  $G$ ; contracting a subgraph of an Eulerian graph results in another Eulerian graph; and, when  $H$  is connected,  $G/H$  can be obtained from  $G$  by successively contracting each edge of  $H$ .

Ellingham, Holton and Little obtained the following characterization of 3-connected cubic graphs  $G$  with the property that no cycle of  $G$  contains a given set of two edges and at most four vertices of  $G$ .

**Lemma 2.4** [18] *Let  $G$  be a 3-connected cubic graph,  $X \subseteq V(G)$  with  $|X| \leq 4$  and  $F = \{e, f\} \subseteq E(G)$ . Then no cycle of  $G$  contains  $X \cup F$  if and only if  $|X| = 4$  and  $G$  has pairwise disjoint subgraphs  $Z_1, Z_2, \dots, Z_m$  such that  $V(Z_1), V(Z_2), \dots, V(Z_m)$  partitions  $V(G)$ ,  $|X \cap Z_i| = 1$  for  $1 \leq i \leq 4$ ,  $e \in E(Z_5, Z_6)$ ,  $f \in E(Z_7, Z_8)$ ,  $\delta(Z_i) = 3$  for all  $1 \leq i \leq m$ , and either:*

- (a)  $m = 8$ , the graph obtained by contracting each  $Z_i$  to a single vertex is the Wagner graph, and  $G$  has the structure illustrated in Figure 3(a), or
- (b)  $m = 10$ , the graph obtained by contracting each  $Z_i$  to a single vertex is the Petersen graph, and  $G$  has the structure illustrated in Figure 3(b).




Figure 3: Graphs in which no Eulerian subgraph contains  $e, f$  and any four given vertices in  $Z_1, Z_2, Z_3$  and  $Z_4$ .

We will need the following extension of Theorem 2.4 to 3-edge-connected graphs which are not necessarily cubic. We use the term *trail* to mean a walk between two vertices in a graph which may repeat vertices but not edges. A *closed trail* is a trail which begins and ends at the same vertex.

**Lemma 2.5** *Let  $G$  be a 3-edge-connected graph,  $X \subseteq V(G)$  with  $|X| \leq 4$  and  $F = \{e, f\} \subseteq E(G)$ . Then no Eulerian subgraph of  $G$  contains  $X \cup F$  if and only if  $|X| = 4$  and  $G$  has pairwise disjoint subgraphs  $Z_1, Z_2, \dots, Z_m$  such that  $V(Z_1), V(Z_2), \dots, V(Z_m)$  partitions  $V(G)$ ,  $|X \cap Z_i| = 1$  for  $1 \leq i \leq 4$ ,  $e \in E(Z_5, Z_6)$ ,  $f \in E(Z_7, Z_8)$ ,  $\delta(Z_i) = 3$  for all  $1 \leq i \leq m$ , and either:*

- (a)  $m = 8$ , the graph obtained by contracting each  $Z_i$  to a single vertex is the Wagner graph, and  $G$  has the structure illustrated in Figure 3(a), or
- (b)  $m = 10$ , the graph obtained by contracting each  $Z_i$  to a single vertex is the Petersen graph, and  $G$  has the structure illustrated in Figure 3(b).

*Proof.* It is not difficult to check that if  $G$  has the specified subgraphs  $Z_1, Z_2, \dots, Z_m$  then no Eulerian subgraph of  $G$  can contain  $X \cup F$ . Hence suppose that no Eulerian subgraph of  $G$  contains  $X \cup F$ . We use induction on  $a(G) := \sum_{v \in V(G)} (d(v) - 3)$  to show that the specified subgraphs  $Z_1, Z_2, \dots, Z_m$  exist. If  $a(G) = 0$  then  $G$  is cubic and the assertion follows immediately from Lemma 2.4. Hence suppose  $a(G) > 0$  and choose  $v \in V(G)$  with  $d(v) \geq 4$ . By Lemma 2.1, we may choose edges  $e_1, e_2$  incident to  $v$  such that the graph  $G_v^{e_1, e_2}$  obtained

by splitting  $e_1, e_2$  at  $v$  in  $G$  is 3-edge-connected. Let  $V(e_i) = \{v, u_i\}$  for  $i = 1, 2$  and let  $G'$  be obtained from  $G - \{e_1, e_2\}$  by adding a new vertex  $z$  and three new edges  $e_1, e_2, e_3$  where  $V(e_i) = \{z, u_i\}$  for  $i = 1, 2$ , and  $V(e_3) = \{z, v\}$ . We give two of the new edges the same labels as the deleted edges so that we have  $E(G) \subseteq E(G')$ . Note that  $G' = G/e_3$ ,  $e_3 \notin \{e, f\}$ , and, if  $e_1$  is a loop in  $G$ , then  $e_1$  is an edge between  $z$  and  $v$  in  $G'$ .

The 3-edge-connectivity of  $G_v^{e_1, e_2}$  implies that  $G'$  is 3-edge-connected, and we have  $X \subseteq V(G) \subseteq V(G')$  and  $F \subseteq E(G) \subseteq E(G')$ . Since no Eulerian subgraph of  $G$  can contain  $X \cup F$ , no Eulerian subgraph of  $G'$  can contain  $X \cup F$ . Since  $a(G') < a(G)$  we may use induction to deduce that the specified subgraphs  $Z'_1, Z'_2, \dots, Z'_m$  exist for  $G'$ . If  $e_3 \in E(Z'_i)$  for some  $1 \leq i \leq m$  then we may construct the required subgraphs  $Z_1, Z_2, \dots, Z_m$  for  $G$  by putting  $Z_i = Z'_i/e_3$  and  $Z_j = Z'_j$  for all  $i \neq j$ . Thus we may assume that  $e_3 \notin E(Z'_i)$  for all  $1 \leq i \leq m$ . We will show that this case cannot occur by constructing an Eulerian subgraph  $H$  of  $G$  which contains  $X \cup F$ . Let  $\tilde{G}$  be the graph obtained from  $G'$  by contracting each subgraph  $Z'_i$  to a single vertex  $z_i$ .

Suppose  $m = 8$ . Then  $\tilde{G}$  is isomorphic to the Wagner graph and we may assume by symmetry that  $e_3$  is incident to either  $z_1$  and  $z_5$ , or  $z_1$  and  $z_3$ . Consider the cycle  $C = z_5 z_6 z_3 z_7 z_8 z_4 z_2 z_5$  of  $\tilde{G}$ . We may extend  $E(C)$  to the Eulerian subgraph  $H$  of  $G = G'/e_3$  which contains  $X \cup F$  as follows. We first assume that  $V(e_3) = \{z_1, z_5\}$ . For  $i \neq 1, 5$  we construct a trail  $P_i$  in  $Z'_i$  joining the two vertices incident to  $C$  and passing through any vertex in  $X \cap V(Z_i)$ . For  $i = 5$  we construct a trail  $P_5$  in  $Z'_5$  joining the vertices incident to  $C$  and passing through the vertex incident to  $e_3$ . For  $i = 1$  we construct a closed trail  $C_1$  in  $Z'_1$  containing the vertex incident to  $e_3$  and the vertex in  $X \cap V(Z'_1)$ . (These trails exist since  $G'$  is 3-edge-connected and hence  $Z_i^* = G'/(G' - Z'_i)$  is 3-edge-connected for all  $1 \leq i \leq 8$ .) We then choose  $H$  to be the subgraph of  $G$  induced by  $\bigcup_{i=2}^8 E(P_i) \cup E(C_1) \cup E(C)$ . We proceed similarly when  $V(e_3) = \{z_1, z_3\}$  by interchanging the roles of  $Z_5$  and  $Z_3$  in the construction.

Suppose that  $m = 10$ . Then  $\tilde{G}$  is isomorphic to the Petersen graph and we may assume by symmetry that  $e_3$  is incident to either  $z_1$  and  $z_7$ , or  $z_1$  and  $z_9$ , or  $z_9$  and  $z_{10}$ . In the first two cases we may proceed as in the previous paragraph, using the cycle  $C = z_5 z_6 z_2 z_7 z_8 z_3 z_9 z_{10} z_4 z_5$  of  $\tilde{G}$ . In the case when  $V(e) = \{z_9, z_{10}\}$ , we proceed similarly using the two disjoint cycles  $C_1 = z_1 z_9 z_3 z_8 z_7 z_1$  and  $C_2 = z_{10} z_2 z_6 z_5 z_4 z_{10}$  in  $\tilde{G}$ . (These cycles give rise to two disjoint Eulerian subgraphs of  $G'$  which become one Eulerian subgraph in  $G = G'/e_3$ .) ■

### 2.3 Three inequalities

The purpose of this subsection is to present three inequalities that will be used to estimate the weight of an Eulerian subgraph obtained by combining several smaller Eulerian subgraphs. The first is elementary.

**Lemma 2.6** *Let  $n_1, n_2$  be nonnegative reals. Then for any  $0 < c \leq 1$ ,*

$$n_1^c + n_2^c \geq (n_1 + n_2)^c.$$

**Lemma 2.7** *Let  $s$  be a positive real number and  $\beta$  be the root of  $(s+2)^x - s^x = 1$  in  $(0, 1)$ . Then for any real numbers  $n_1, n_2, n_3, \gamma$  satisfying  $n_1 \geq sn_3$ ,  $n_2 \geq n_3 \geq 0$ , and  $0 < \gamma \leq \beta$  we have*

$$n_1^\gamma + n_2^\gamma \geq (n_1 + n_2 + n_3)^\gamma.$$



*Proof.* It is not difficult to check that  $(s+2)^x - s^x = 1$  has a unique root  $\beta \in (0, 1)$  and that  $(s+2)^\gamma - s^\gamma - 1 \leq 0$  for all  $0 < \gamma \leq \beta$ . Let  $f(n_1, n_2, n_3) = n_1^\gamma + n_2^\gamma - (n_1 + n_2 + n_3)^\gamma$ . We show that  $f(n_1, n_2, n_3) \geq 0$  when  $n_1 \geq sn_3$  and  $n_2 \geq n_3 \geq 0$ . We have  $\partial f / \partial n_1 \geq 0$  and  $\partial f / \partial n_2 \geq 0$  since  $0 < \gamma < 1$ , so  $f$  is minimised when  $n_1 = sn_3$  and  $n_2 = n_3$ . Thus

$$f(n_1, n_2, n_3) \geq f(sn_3, n_3, n_3) = (s^\gamma + 1 - (s+2)^\gamma)n_3^\gamma \geq 0.$$

■

**Lemma 2.8** *Suppose  $n_1, \dots, n_k, t, \gamma$  are real numbers with  $k \geq 3$ ,  $0 \leq n_k \leq t \min\{n_1, \dots, n_{k-1}\}$ , and  $0 < \gamma \leq \log_{t+k-1}(k-1)$ . Then*

$$\sum_{i=1}^{k-1} n_i^\gamma \geq \left( \sum_{i=1}^k n_i \right)^\gamma.$$

*Proof.* The assertion of this lemma follows from Lemma 2.6 when  $n_k = 0$ . Thus we may assume  $n_k > 0$ . Hence  $t > 0$  and  $\sum_{i=1}^k n_i > 0$ . Define  $x_i = n_i / \sum_{j=1}^k n_j$ , for  $i = 1, \dots, k$ . Then  $x_1, \dots, x_k \in [0, 1]$ ,  $\sum_{i=1}^k x_i = 1$ , and  $x_k \leq t \min\{x_1, \dots, x_{k-1}\}$ . It suffices to show that  $\sum_{i=1}^{k-1} x_i^\gamma \geq 1$ .

Let  $f(x_1, \dots, x_{k-1}) = \sum_{i=1}^{k-1} x_i^\gamma$ . We first show that the minimum of  $f(x_1, \dots, x_{k-1})$  subject to the constraints that  $x_i \geq x_k/t \geq 0$  for all  $1 \leq i \leq k-1$ ,  $\sum_{i=1}^k x_i = 1$ , and  $x_k$  is fixed, occurs when  $x_i = x_k/t$  for all  $i = 2, \dots, k$ . Let  $(a_1, a_2, \dots, a_{k-1})$  be a point at which this minimum occurs and is such that  $a_1$  is as large as possible. By symmetry we have  $a_1 \geq a_i$  for all  $2 \leq i \leq k-1$ . We may use elementary calculus and the facts that  $a_1 \geq a_i$  and  $0 \leq \gamma < 1$  to deduce that  $(a_1 + \epsilon)^\gamma + (a_i - \epsilon)^\gamma \leq a_1^\gamma + a_i^\gamma$  for all  $\epsilon \geq 0$ . The choice of  $(a_1, a_2, \dots, a_{k-1})$  now implies that  $a_i = x_k/t$  for all  $2 \leq i \leq k-1$ ,  $a_1 = 1 - (t+k-2)x_k/t$ , and  $f(a_1, \dots, a_{k-1}) = (1 - (t+k-2)x_k/t)^\gamma + (k-2)(x_k/t)^\gamma =: g(x_k)$ .

Since  $\sum_{i=1}^k x_i = 1$  we have  $\sum_{i=1}^k tx_i = t$ . We can now use the fact that  $x_k \leq tx_i$  for all  $1 \leq i \leq k-1$  to deduce that  $x_k \leq t/(t+k-1)$ . We complete the proof by showing that  $g(x_k) \geq 1$  for  $0 \leq x_k \leq t/(t+k-1)$ . It is not difficult to see that  $g'(x_k) = 0$  has a unique solution, and that  $g''(x_k) < 0$  for  $0 \leq x_k \leq t/(t+k-1)$ . Hence, the minimum of  $g(x_k)$  is achieved at  $x_k = 0$  or  $x_k = t/(t+k-1)$ . We have  $g(0) = 1$ , and

$$g\left(\frac{t}{t+k-1}\right) = (k-1)(t+k-1)^{-\gamma} \geq (k-1)(t+k-1)^{-\log_{t+k-1}(k-1)} = 1.$$

Therefore,  $f(x_1, \dots, x_{k-1}) \geq 1$ .

■

We will use the following special cases of Lemmas 2.7 and 2.8.

**Corollary 2.9** *Let  $\alpha \approx 0.753$  be the real root of  $4^{1/x} - 3^{1/x} = 2$ . Then:*

(a) *for all real numbers  $n_1, n_2, n_3$  satisfying  $n_1 \geq 3^{1/\alpha}n_3$  and  $n_2 \geq n_3 \geq 0$  we have*

$$n_1^\alpha + n_2^\alpha \geq (n_1 + n_2 + n_3)^\alpha;$$

(b) *for all real numbers  $n_1, n_2, n_3, n_4$  satisfying  $0 \leq n_4 \leq \min\{n_1, n_2, n_3\}$  we have*

$$n_1^\alpha + n_2^\alpha + n_3^\alpha \geq (n_1 + n_2 + n_3 + n_4)^\alpha;$$

(c) for all real numbers  $n_1, n_2, n_3, n_4, n_5$  satisfying  $0 \leq n_5 \leq (4^{1/\alpha} - 4) \min\{n_1, n_2, n_3, n_4\}$  we have

$$n_1^\alpha + n_2^\alpha + n_3^\alpha + n_4^\alpha \geq (n_1 + n_2 + n_3 + n_4 + n_5)^\alpha.$$

*Proof.* Part (a) follows from Lemma 2.7 by taking  $s = 3^{1/\alpha}$  and using the fact that  $(3^{1/\alpha} + 2)^\alpha - (3^{1/\alpha})^\alpha = 4 - 3 = 1$ . Parts (b) and (c) follow from Lemma 2.8 by taking  $k = 4$  and  $t = 1$ , and  $k = 5$  and  $t = 4^{1/\alpha} - 4$ , respectively. ■

### 3 Eulerian subgraphs of 3-edge-connected graphs

In this section we prove an edge weighted version of Theorem 1.1. Let  $G$  be a graph and let  $w : E(G) \rightarrow \{1, 2\}$ . For any  $H \subseteq G$  let  $w(H) = \sum_{e \in E(H)} w(e)$ , and for any  $S \subseteq E(G)$  let  $w(S) = \sum_{e \in S} w(e)$ . We will show

**Theorem 3.1** *Let  $G$  be a 3-edge-connected graph,  $e, f \in E(G)$ , and  $w : E(G) \rightarrow \{1, 2\}$ . Suppose  $G \neq K_2^3$ . Then  $G$  contains an Eulerian subgraph  $H$  such that  $e, f \in E(H)$  and  $w(H) \geq (w(G)/6)^\alpha + 2$ , where  $\alpha \approx 0.753$  is the real root of  $4^{1/x} - 3^{1/x} = 2$ .*

The multiplicative constant  $(1/6)^\alpha$  in Theorem 1.1 is chosen to simplify its proof; it may be improved by considering other exceptional graphs in addition to  $K_2^3$ . Note that the conclusion of Theorem 3.1 does not hold for  $K_2^3$  because of the additive constant 2. We need this additive constant for the inductive step in our proof.

We first need to deal with graphs with few edges to provide a basis for our induction.

**Lemma 3.2** *Theorem 3.1 holds for graphs with at most 6 edges.*

*Proof.* The assertion of Theorem 3.1 clearly holds if  $G$  is Eulerian. So assume that  $u, v$  are vertices of  $G$  with odd degree. Since  $G$  is 3-edge-connected and  $|E(G)| \leq 6$ ,  $|G| \leq 4$ .

If  $|G| = 4$  then  $G = K_4$ . If  $|G| = 2$  then, since  $G \neq K_2^3$ ,  $G$  is obtained from  $K_2^3$  by adding one, two or three edges, which can be either two more  $uv$ -edges and at most one loop, or all loops. In each case it is easy to check that the desired Eulerian subgraph  $H$  exists.

Now assume  $|G| = 3$ . Let  $w$  denote the vertex of  $G$  other than  $u, v$ . Since  $G$  is 3-edge-connected and  $|E(G)| \leq 6$ , we see that  $G$  has at most two edges between  $u$  and  $v$ . If there is no edge between  $u$  and  $v$ , then  $G$  is obtained from a path of length 2 by tripling each edge, and it is easy to find the desired  $H$ . If there is exactly one edge between  $u$  and  $v$ , then  $d(u) = d(v) = 3$  (as  $|E(G)| \leq 6$ ) and  $d(w) = 4$  or  $6$  (if  $d(w) = 6$  then there is a loop on  $w$ ); and the desired  $H$  can be found directly. Finally, assume that there are precisely two edges between  $u$  and  $v$ . Since  $G$  is 3-edge-connected and by symmetry, we may assume  $d(u) = 5$  (so that there are 3 edges between  $u$  and  $w$ ). Then  $d(v) = 3$  (since  $|E(G)| \leq 6$ ), and there is just one edge between  $v$  and  $w$ . Again the desired  $H$  exists. ■

The next lemma will be used to construct the desired Eulerian subgraph of  $G$  from an Eulerian subgraph of a graph obtained from  $G$  by contracting several disjoint induced subgraphs.

**Lemma 3.3** *Let  $G$  be a 3-edge-connected graph,  $w : E(G) \rightarrow \{1, 2\}$ , and let  $C_1, \dots, C_k$  be disjoint induced subgraphs of  $G$  such that  $\delta(C_i) = 3$  and  $|E(C_i)| < |E(G)| - 3$  for all  $i = 1, \dots, k$ . Let  $\tilde{G}$  denote the graph obtained from  $G$  by contracting each subgraph  $C_i$  to a single*

vertex  $c_i$ . Suppose Theorem 3.1 holds for all graphs with fewer edges than  $G$ , and assume that  $\tilde{G}$  contains an Eulerian subgraph  $\tilde{H}$  such that  $c_i \in V(\tilde{H})$  for all  $i$ . Then  $G$  contains an Eulerian subgraph  $H$  such that the edges of  $G$  corresponding to the edges in  $\tilde{H}$  are in  $H$  and

$$w(H) \geq \sum_{i=1}^k (w(C_i)/6)^\alpha + w(\tilde{H}).$$

*Proof.* For each  $i$ , let  $e_i, f_i$  denote the edges of  $\tilde{H}$  incident with  $c_i$ . Let  $C_i^*$  be obtained from  $G$  by contracting  $G - C_i$  to a single vertex  $c_i^*$ . Since  $\delta(C_i) = 3$  and  $G$  is 3-edge-connected,  $C_i^*$  is 3-edge-connected. Assign the edges incident with  $c_i^*$  weight 1.

Since  $|E(C_i)| < |E(G)| - 3$ , we have  $|E(C_i^*)| < |E(G)|$ . If  $C_i^* \neq K_2^3$  then, by assumption,  $C_i^*$  contains an Eulerian subgraph  $H_i$  such that  $e_i, f_i \in E(H_i)$  and  $w(H_i) \geq (w(C_i^*)/6)^\alpha + 2 \geq (w(C_i)/6)^\alpha + 2$ . On the other hand, if  $C_i^* = K_2^3$  then  $E(C_i) = \emptyset$ ,  $w(C_i) = 0$  and we may also construct an Eulerian subgraph  $H_i$  such that  $e_i, f_i \in E(H_i)$  and  $w(H_i) = 2 = (w(C_i)/6)^\alpha + 2$ .

Since  $d(c_i^*) = 3$ , we see that  $H_i$  uses exactly two edges at  $c_i^*$ , namely  $e_i$  and  $f_i$ . Then  $\bigcup_{i=1}^k E(H_i) \cup E(\tilde{H})$  induces an Eulerian subgraph  $H$  of  $G$  such that

$$w(H) \geq \sum_{i=1}^k (w(H_i) - 2) + w(\tilde{H}) \geq \sum_{i=1}^k (w(C_i)/6)^\alpha + w(\tilde{H}).$$

**Lemma 3.4** *Let  $L$  be a 3-edge-connected graph,  $w : E(G) \rightarrow \{1, 2\}$ , and  $z_1, z_2$  be two adjacent vertices of degree three in  $L$ . Let  $L'$  be obtained from  $L$  by deleting the edge joining  $z_1$  and  $z_2$ , and then suppressing  $z_1, z_2$  to two edges  $k_1, k_2$ , respectively, of weight 1. Suppose Theorem 3.1 holds for all graphs with fewer edges than  $L$ . Then  $L'$  has an Eulerian subgraph  $H$  with  $k_1, k_2 \in E(H)$  and  $w(H - \{k_1, k_2\}) \geq (w(L')/6)^\alpha$ .*

*Proof.* We use an inner induction on  $|E(L)|$ . If  $L'$  is 3-edge-connected then we may apply Theorem 3.1 to  $L'$  to find an Eulerian subgraph  $H$  with  $k_1, k_2 \in E(H)$  and  $w(H') \geq (w(L')/6)^\alpha + 2$ . Then  $w(H' - \{k_1, k_2\}) \geq (w(L')/6)^\alpha$  as required.

Hence suppose that  $L'$  is not 3-edge-connected. Since  $L$  is 3-edge-connected,  $L'$  is 2-edge-connected and every 2-edge-cut of  $L'$  separates  $k_1$  and  $k_2$ . Choose a 2-edge-cut  $\{g, h\}$  of  $L'$  and let  $L_1^*, L_2^*$  be the components of  $L' - \{g, h\}$  with  $k_1 \in E(L_1^*)$  and  $k_2 \in E(L_2^*)$ . For  $i = 1, 2$ , construct  $L_i'$  from  $L_i^*$  by adding a new edge  $f_i$  of weight 1 between the endvertices of  $g$  and  $h$  in  $L_i^*$ . Let  $L_i$  be obtained from  $L_i'$  by subdividing  $k_i$  and  $f_i$  with two new vertices  $z_1'$  and  $z_2'$  and then adding an edge between  $z_1'$  and  $z_2'$ . Then  $L_i$  is 3-edge-connected since it can be obtained from  $L$  by contracting  $L_{3-i} \cup \{z_{3-i}\}$  to a single vertex. We may apply the inner induction to  $L_i$  to deduce that  $L_i'$  has an Eulerian subgraph  $H_i$  with  $k_i, f_i \in E(H_i)$  and  $w(H_i - \{k_i, f_i\}) \geq (w(L_i')/6)^\alpha$ . Then  $E(H_1 - f_1) \cup E(H_2 - f_2) \cup \{g, h\}$  induces an Eulerian subgraph  $H$  of  $L'$  with  $k_1, k_2 \in E(H)$  and

$$w(H - \{k_1, k_2\}) \geq (w(L_1')/6)^\alpha + (w(L_2')/6)^\alpha + w(g) + w(h) \geq (w(L')/6)^\alpha$$

by Lemma 2.6. ■

**Proof of Theorem 3.1.** We use induction on  $|E(G)|$ . By Lemma 3.2, we may assume:

$$|E(G)| \geq 7. \tag{3.1}$$

As induction hypothesis, we assume that:

$$\text{the theorem holds for all graphs with fewer than } |E(G)| \text{ edges.} \quad (3.2)$$

We may also assume that:

$$\text{neither } e \text{ nor } f \text{ belongs to a splittable pair in } G. \quad (3.3)$$

For, suppose by symmetry that  $\{e, g\}$  is a splittable pair in  $G$ , with  $V(e) = \{v, u\}$  and  $V(g) = \{v, x\}$ . Let  $G' := G_v^{e,g}$  be obtained from  $G$  by splitting  $\{e, g\}$  at  $v$ , and assign weight 1 to the new edge  $e'$  which corresponds to  $e$  and  $g$ , and also to the other new edge  $e''$  when  $d(v) = 4$ . Let  $f' = f$  if none of  $V(f)$  is suppressed; otherwise let  $f' = e'$  (if  $f = g$ ) or  $f' = e''$  (if  $f \neq g$ ).

Note that  $w(G') \geq w(G) - 6$ . Also note that  $|E(G')| \geq |E(G)| - 2 \geq 5$  (by (3.1)); so  $G' \neq K_2^3$ . Hence, by (3.2),  $G'$  contains an Eulerian subgraph  $H'$  such that  $e', f' \in E(H')$  and  $w(H') \geq (w(G')/6)^\alpha + 2$ . Let  $H$  be obtained from  $H'$  by replacing  $e'$  with  $e$  and  $g$  and replacing  $e''$  (if it exists in  $H'$ ) with the corresponding edges in  $G$ . Then by Lemma 2.6,

$$w(H) \geq w(H') + 1 \geq (w(G')/6)^\alpha + 2 + 1 \geq (w(G)/6)^\alpha + 2.$$

□

Assumption (3.3) implies in particular that neither  $e$  nor  $f$  is a loop or is adjacent to a loop.

We may further assume that:

$$e \text{ and } f \text{ are not adjacent.} \quad (3.4)$$

Suppose on the contrary that  $V(e) = \{v, u\}$  and  $V(f) = \{v, x\}$ . Then  $d(v) = 3$  by (3.3) and Lemma 2.1. So  $x \neq u$ ; for otherwise, by (3.3) and Lemma 2.1 we would also have  $d(u) = 3$ , and (since  $G \not\cong K_2^3$ )  $G$  would not be 3-edge-connected.

Let  $g$  denote the edge incident with  $v$  other than  $e$  and  $f$ , and let  $y$  be the end of  $g$  other than  $v$ . Note that  $y \neq v$  as  $d(v) = 3$ . Let  $G'$  be obtained from  $G - g$  by suppressing degree 2 vertices (namely,  $v$  and possibly  $y$ ) and assign weight 1 to the new edge(s) which resulted from the vertex suppression(s). So  $w(G') \geq w(G) - 6$  if both  $e$  and  $f$  have weight 1 in  $G$ ; otherwise  $w(G') \geq w(G) - 8$  and  $e$  or  $f$  has weight 2 in  $G$ . By (3.1),  $|E(G')| \geq 4$ , and hence  $G' \neq K_2^3$ . Let  $e'$  denote the edge of  $G'$  obtained by suppressing  $v$ , and if  $d(y) = 3$  let  $e''$  denote the edge of  $G'$  obtained by suppressing  $y$ .

First, consider the case when  $G'$  is 3-edge-connected. Let  $f'$  be an arbitrary edge of  $G'$  that is adjacent to  $e'$ . By (3.2),  $G'$  contains an Eulerian subgraph  $H'$  such that  $e', f' \in E(H')$  and  $w(H') \geq (w(G')/6)^\alpha + 2$ . Let  $H$  be obtained from  $H'$  by replacing  $e'$  with  $e$  and  $f$  and by replacing  $e''$  (if  $e''$  exists and belongs to  $H'$ ) with the edges of  $G - g$  incident with  $y$ . Now  $H$  is an Eulerian subgraph of  $G$  and  $e, f \in H$ . If both  $e$  and  $f$  have weight 1 in  $G$  then  $w(H) \geq w(H') + 1 \geq ((w(G) - 6)/6)^\alpha + 2 + 1 \geq (w(G)/6)^\alpha + 2$  (by Lemma 2.6). Otherwise,  $w(H) \geq w(H') + 2 \geq ((w(G) - 8)/6)^\alpha + 2 + 2 \geq (w(G)/6)^\alpha + 2$  (by Lemma 2.6).

Thus we may assume that  $G'$  is not 3-edge-connected. Then  $G'$  has a 2-edge-cut  $S = \{g_1, g_2\}$  such that  $u, x$  are contained in the same component of  $G' - S$ , say  $G_1$ . We choose  $S$  such that  $G_1$  is minimal (under subgraph containment). Let  $G_2$  denote the other component of  $G' - S$ , and let  $V(g_1) = \{u_1, u_2\}$  and  $V(g_2) = \{v_1, v_2\}$  with  $u_i, v_i \in G_i$  for  $i = 1, 2$ .

Let  $G'_1$  be obtained from  $G_1$  by adding an edge  $f'$  between  $u_1$  and  $v_1$  (which may be a loop) and assign  $f'$  weight 1. By the minimality of  $G_1$  we see that  $G'_1$  is 3-edge-connected. When  $G'_1 \neq K_2^3$  we may use (3.2) to deduce that  $G'_1$  contains an Eulerian subgraph  $H'_1$  such that  $e', f' \in H'_1$  and  $w(H'_1) \geq (w(G'_1)/6)^\alpha + 2$ . In the case when  $G'_1 = K_2^3$ , we choose  $H'_1$  to be the Eulerian subgraph of  $G'_1$  with  $E(H'_1) = \{e', f'\}$  and  $w(H'_1) = w(e') + w(f') = 2$ .

Let  $G'_2$  be obtained from  $G$  by contracting  $G[V(G_1) \cup \{v\}]$  to a single vertex  $z$ . Then  $G'_2$  is 3-edge-connected. Assign weight 1 to  $g, g_1, g_2$  in  $G'_2$ . Since  $G'$  is not 3-edge-connected, we see that  $|E(G_2)| \geq 1$ ; so  $G'_2 \neq K_2^3$ . Hence by (3.2),  $G'_2$  contains an Eulerian subgraph  $H'_2$  such that  $g_1, g_2 \in H'_2$  and  $w(H'_2) \geq (w(G'_2)/6)^\alpha + 2$ .

Let  $H$  be the subgraph of  $G$  induced by  $E(H'_1 - \{e', f'\}) \cup \{e, f\} \cup E(H'_2)$ . Then  $H$  is an Eulerian subgraph of  $G$  (as both  $H'_1$  and  $H'_2$  are 2-edge-connected),  $e, f \in E(H)$  and  $w(H) \geq w(H'_1) + w(H'_2)$ . If  $G'_1 = K_2^3$  then  $w(G'_2) \geq w(G) - 9$  and  $w(H) \geq 2 + ((w(G) - 9)/6)^\alpha + 2 \geq (w(G)/6)^\alpha + 2$  by Lemma 2.6. So assume  $G'_1 \neq K_2^3$ . Note that  $w(G'_1) + w(G'_2) \geq w(G_1) + 1 + w(G_2) + 3 \geq w(G) - 8$ . Then  $w(H) \geq (w(G'_1)/6)^\alpha + 2 + (w(G'_2)/6)^\alpha + 2 \geq (w(G)/6)^\alpha + 2$  again by Lemma 2.6.  $\square$

We say that a 3-edge-cut  $S$  of  $G$  is *trivial* if some component of  $G - S$  consists of a single vertex and no edge. Otherwise we say that  $S$  is *non-trivial*. We may assume that:

$$\text{neither } e \text{ nor } f \text{ is contained in a non-trivial 3-edge-cut of } G. \quad (3.5)$$

For, suppose  $S = \{e, g_1, g_2\}$  is a 3-edge-cut of  $G$  and let  $G_1, G_2$  be the components of  $G - S$  such that  $|E(G_i)| \geq 1$  for  $i = 1, 2$ . Let  $V(e) = \{u_1, u_2\}$ ,  $V(g_1) = \{x_1, x_2\}$  and  $V(g_2) = \{y_1, y_2\}$  with  $u_i, x_i, y_i \in V(G_i)$ ,  $i = 1, 2$ . Let  $G'_i$  be obtained from  $G$  by contracting  $G_{3-i}$ , for  $i = 1, 2$ . By symmetry, assume  $f \in E(G_1) \cup S$ . Assign weight 1 to  $e, g_1, g_2$  in both  $G'_1$  and  $G'_2$ . Then  $w(G'_1) + w(G'_2) \geq w(G)$  as the weight of every edge of  $G$  is at most 2.

Note that for  $i = 1, 2$ ,  $G'_i$  is 3-edge-connected, and  $G'_i \neq K_2^3$  (since  $|E(G_i)| \geq 1$ ). So by (3.2),  $G'_1$  contains an Eulerian subgraph  $H'_1$  such that  $e, f \in H'_1$  and  $w(H'_1) \geq (w(G'_1)/6)^\alpha + 2$ . Without loss of generality, we may assume that  $g_1 \in H'_1$ . By (3.2),  $G'_2$  contains an Eulerian subgraph  $H'_2$  such that  $e, g_1 \in H'_2$  and  $w(H'_2) \geq (w(G'_2)/6)^\alpha + 2$ .

Let  $H$  be the subgraph of  $G$  induced by  $E(H'_1) \cup E(H'_2)$ . Then  $H$  is an Eulerian subgraph of  $G$  containing  $e, f$  and  $w(H) \geq w(H'_1) + w(H'_2) - 2 \geq (w(G'_1)/6)^\alpha + (w(G'_2)/6)^\alpha + 2 \geq (w(G)/6)^\alpha + 2$  by Lemma 2.6.  $\square$

We may also assume that:

$$\text{for any 3-edge-cut } S \text{ of } G, e \text{ and } f \text{ are contained in the same component of } G - S. \quad (3.6)$$

Suppose on the contrary that  $S = \{g_1, g_2, g_3\}$  is a 3-edge-cut of  $G$  such that  $e \in G_1$  and  $f \in G_2$ , where  $G_1, G_2$  are the components of  $G - S$ . Let  $V(g_1) = \{x_1, x_2\}$ ,  $V(g_2) = \{y_1, y_2\}$ , and  $V(g_3) = \{z_1, z_2\}$  such that  $x_i, y_i, z_i \in G_i$  for  $i = 1, 2$ .

Let  $G'_i$  be obtained from  $G$  by contracting  $G_{3-i}$ , for  $i = 1, 2$ . In both  $G'_1$  and  $G'_2$ , assign weight 1 to  $g_1, g_2$  and  $g_3$ . Then  $w(G_i) = w(G'_i) - 3$ ; so  $w(G'_1) + w(G'_2) = w(G_1) + w(G_2) + 6 \geq w(G)$ .

Note that  $G'_i$  is 3-edge-connected and, since  $|E(G_i)| \geq 1$ ,  $G'_i \neq K_2^3$ . By symmetry, we may assume  $|G'_1| \leq |G'_2|$ .<sup>1</sup> By (3.2),  $G'_1$  contains an Eulerian subgraph  $H'_1$  such that  $e, g_1 \in H'_1$

<sup>1</sup>This assumption will not be used in the proof of (3.6) but will be important when we convert the proof into a polynomial time algorithm in Section 5.

and  $w(H'_1) \geq (w(G'_1)/6)^\alpha + 2$ . Without loss of generality, we may assume  $g_2 \in H'_1$  (so  $g_3 \notin H'_1$ ). By (3.2) again,  $G'_1$  contains an Eulerian subgraph  $H''_1$  such that  $e, g_3 \in H''_1$  and  $w(H''_1) \geq (w(G'_1)/6)^\alpha + 2$ . We now have a symmetry between  $g_1$  and  $g_2$ , and we may thus assume that  $g_1 \in H''_1$ .

In  $G'_2$  we find an Eulerian subgraph  $H'_2$  such that  $f, g_1 \in H'_2$  and  $w(H'_2) \geq (w(G'_2)/6)^\alpha + 2$ . If  $g_2 \in H'_2$ , let  $H$  be the subgraph of  $G$  induced by  $E(H'_1) \cup E(H'_2)$ ; otherwise we have  $g_3 \in H'_2$  and we let  $H$  be the subgraph of  $G$  induced by  $E(H''_1) \cup E(H'_2)$ . Then  $H$  is an Eulerian subgraph of  $G$  such that  $e, f \in H$ , and  $w(H) = w(H'_1) + w(H'_2) - 2$  or  $w(H) = w(H''_1) + w(H'_2) - 2$ . Hence  $w(H) \geq (w(G'_1)/6)^\alpha + (w(G'_2)/6)^\alpha + 2 \geq (w(G)/6)^\alpha + 2$  by Lemma 2.6.  $\square$

We may further assume that:

$$\text{the vertices incident to } e \text{ and } f \text{ all have degree 3 in } G. \quad (3.7)$$

Suppose on the contrary that  $V(e) = \{u, v\}$  and  $d(v) \geq 4$ . By (3.3),  $e$  is not in any splittable pair of  $G$ . Lemmas 2.1 and 2.2 now imply that  $d(v) = 5$ , and  $V(G) - \{v\}$  has a partition  $Y_0, Y_1, Y_2$  such that  $u \in Y_0$ ,  $\delta(v, Y_0) = 1$ ,  $\delta(v, Y_1) = \delta(v, Y_2) = 2$ ,  $\delta(Y_0, Y_1) = \delta(Y_0, Y_2) = 1$ , and  $\delta(Y_1, Y_2) = 0$ . See Figure 1. By (3.5),  $|Y_0| = 1$ . So by (3.4),  $f \in Y_1$  or  $f \in Y_2$ . By symmetry, we may assume  $f \in Y_1$ . But then the edges from  $Y_1$  to  $\{u, v\}$  form a non-trivial 3-edge-cut in  $G$  which separates  $e$  from  $f$ , contradicting (3.6).  $\square$

Let  $V(e) = \{u, v\}$ . By (3.7),  $d(u) = d(v) = 3$ . Let  $g_i, i = 1, 2$ , denote the other two edges incident with  $u$  with  $V(g_i) = \{u, u_i\}$ ; and let  $h_i, i = 1, 2$ , denote the other two edges incident with  $v$  with  $V(h_i) = \{v, v_i\}$ . Since  $G$  is 3-edge-connected and  $d(v) = d(u) = 3$ ,  $e$  is the only edge between  $u$  and  $v$ . So  $v \neq u_i$  and  $u \neq v_i$  for  $i = 1, 2$ .

Let  $G_i, i = 1, 2$ , be obtained from  $G - g_i$  by suppressing  $u$  to  $e'$  and, if  $d_G(u_i) = 3$ , suppressing  $u_i$  to  $e_i$ . Define  $f' = f$  if  $u_i \notin V(f)$  or  $u_i$  is not suppressed, and otherwise let  $f' = e_i$ . Similarly, let  $H_i, i = 1, 2$ , be obtained from  $G - h_i$  by suppressing  $v$  to  $e'$  and, if  $d_G(v_i) = 3$ , suppressing  $v_i$  to  $f_i$ . Define  $f' = f$  if  $v_i \notin V(f)$  or  $v_i$  is not suppressed, and otherwise let  $f' = f_i$ .

We may assume that:

$$G_1, G_2, H_1, \text{ and } H_2 \text{ are not 3-edge-connected.} \quad (3.8)$$

Suppose on the contrary that  $G_1$  is 3-edge-connected. Assign weight 1 to the edges of  $G_1$  which resulted from vertex suppressions. Note that  $w(G_1) \geq w(G) - 6$  if both  $e$  and  $g_2$  have weight 1 in  $G$ ; otherwise  $w(G_1) \geq w(G) - 8$ . By (3.2),  $G_1$  contains an Eulerian subgraph  $H'$  such that  $e', f' \in E(H')$  and  $w(H') \geq (w(G'_1)/6)^\alpha + 2$ . Let  $H$  be obtained from  $H'$  by replacing  $e'$  with  $e$  and  $g_1$  and, if  $e_1$  exists and belongs to  $H'$ , replacing it with the suppressed edges at  $u_1$ . Then  $H$  is an Eulerian subgraph of  $G$  such that  $e, f \in H$ . If  $e$  and  $g_2$  both have weight 1 in  $G$  then  $w(H) \geq w(H') + 1 \geq ((w(G) - 6)/6)^\alpha + 1 + 2 \geq (w(G)/6)^\alpha + 2$  by Lemma 2.6. So assume that  $e$  or  $g_2$  has weight 2 in  $G$ . Then  $w(H) \geq w(H') + 2 \geq ((w(G) - 8)/6)^\alpha + 2 + 2 \geq (w(G)/6)^\alpha + 2$ , again by Lemma 2.6.  $\square$

Since  $G$  is 3-edge-connected,  $G_i, H_i$  are all 2-edge-connected. By (3.8), we may choose a 2-edge-cut  $S_i$  of  $G_i$ . Note that  $S_i \cup \{g_i\}$  is a 3-edge-cut in  $G$ ; so by (3.6), some component  $C_i$  of  $G - S_i$  satisfies  $e, f \notin C_i$ . We choose  $S_i$  and  $C_i$  such that  $C_i$  is maximal. Then  $|E(C_i)| \geq 1$ ; as otherwise,  $G_i$  would be 3-edge-connected (by the maximality of  $C_i$ ). Similarly, we choose  $T_i$  to be a 2-edge-cut of  $H_i$ ,  $D_i$  to be the component of  $H_i - T_i$  such that  $e, f \notin D_i$ , and suppose

that  $T_i, D_i$  have been chosen such that  $D_i$  is maximal (so  $|E(D_i)| \geq 1$ ). We remark that the argument given below to verify (3.9) does not use the maximality of  $C_i$  and  $D_i$ ; this maximality will be used later to ensure that the graph obtained from  $G_i$ , or  $H_i$ , by contracting  $C_i$ , or  $D_i$ , to a single vertex of degree two and then suppressing this vertex, is 3-edge-connected.

We next show that:

$$C_1, C_2, D_1 \text{ and } D_2 \text{ are pairwise disjoint.} \quad (3.9)$$

First, suppose  $C_1 \cap C_2 \neq \emptyset$ . Since  $u, v \notin V(C_1 \cup C_2)$ ,  $C_1 \cup C_2 \neq V(G)$ . Since  $G$  is 3-edge-connected, we have

$$3 + 3 = \delta_G(C_1) + \delta_G(C_2) \geq \delta_G(C_1 \cap C_2) + \delta_G(C_1 \cup C_2) \geq 3 + 3.$$

Thus equality must hold throughout and, in particular,  $\delta_G(C_1 \cup C_2) = 3$ . Since  $d_G(u) = 3$  and  $\delta_G(u, C_1 \cup C_2) = 2$  we have  $\delta_G(C_1 \cup C_2 \cup \{u\}) = 2$ . Since  $v \notin V(C_1 \cup C_2) \cup \{u\}$ , this contradicts the fact that  $G$  is 3-edge-connected.

Next, suppose  $C_1 \cap D_1 \neq \emptyset$ . We may deduce as above that  $\delta_G(C_1 \cup D_1) = 3$ . Since  $d_G(u) = 3 = d_G(v)$  and  $\delta_G(\{u, v\}, C_1 \cup D_1) = 2$ , we have  $\delta_G(C_1 \cup D_1 \cup \{u, v\}) = 3$ . This contradicts (3.6) since  $f$  is an edge of  $G - (C_1 \cup D_1 \cup \{u, v\})$ .

Similar arguments apply to all other pairs.  $\square$

Our current knowledge on the structure of  $G$  is illustrated in Figure 4.




Figure 4: The structure of  $G$  around  $e$ . Note that for each  $i = 1, 2$ , one of the edges leaving  $C_i$  may be incident to  $D_j$ , for some  $j = 1, 2$ .

Let  $V(f) = \{u', v'\}$  and let  $g'_i, h'_i, S'_i, C'_i, D'_i$  be defined with respect to  $f$  in the same way that  $g_i, h_i, S_i, C_i, D_i$  were defined with respect to  $e$ . Then  $|E(C'_i)| \geq 1$  and  $|E(D'_i)| \geq 1$  for  $i = 1, 2$ , and  $C'_1, C'_2, D'_1, D'_2$  are pairwise disjoint by (3.9) and symmetry. Let  $\mathcal{S} = \{C_1, C_2, D_1, D_2\}$ ,  $\mathcal{S}' = \{C'_1, C'_2, D'_1, D'_2\}$ , and  $K = G - \{u, v, u', v'\} - \bigcup_{X \in \mathcal{S} \cup \mathcal{S}'} X$ .

We next show that:

$$\text{for all } X \in \mathcal{S} \text{ and } X' \in \mathcal{S}' \text{ we have either } X = X' \text{ or } X \cap X' = \emptyset. \quad (3.10)$$

Suppose  $X \cap X' \neq \emptyset$ . Then  $\delta_G(X \cap X') \geq 3$  since  $G$  is 3-edge-connected. Since  $u, v \notin V(X \cup X')$  we also have  $\delta_G(X \cup X') \geq 3$ . Hence

$$3 + 3 \geq \delta_G(X) + \delta_G(X') \geq \delta_G(X \cap X') + \delta_G(X \cup X') \geq 3 + 3.$$

This implies that  $\delta_G(X \cup X') = 3$ . The maximality of  $X$  and  $X'$  now gives  $X = X'$ .  $\square$

We may further assume that:

$$\{C_1, C_2\} \neq \{C'_1, C'_2\}. \quad (3.11)$$

Suppose on the contrary that  $\{C_1, C_2\} = \{C'_1, C'_2\}$ . Relabeling if necessary we have  $C_1 = C'_1$  and  $C_2 = C'_2$ . See the first graph in Figure 5. Let  $k_i$  be the edge from  $C_i$  to  $G - (C_i \cup \{u, u'\})$ ,  $i = 1, 2$ . Let  $G^*$  be obtained from  $G$  by contracting  $G[C_1 \cup C_2 \cup \{u, u'\}]$  to a single vertex  $z$ . Then  $e, f, k_1, k_2$  are the only edges of  $G^*$  incident with  $z$ . See the second graph in Figure 5. The graph  $G^*$  is 3-edge-connected since contraction cannot reduce edge-connectivity.




Figure 5:  $C_1 = C'_1$  and  $C_2 = C'_2$ .

We first consider the case when  $\{e, k_1\}$  is splittable at  $z$  in  $G^*$ . Let  $\tilde{G}$  be the graph obtained from  $G^*$  by splitting  $e, k_1$  at  $z$ , and let  $\tilde{e}, \tilde{f}$  be the edges of  $\tilde{G}$  which correspond to  $e, k_1$  and  $f, k_2$ , respectively. Assign weight 1 to  $\tilde{e}$  and to  $\tilde{f}$ . By induction,  $\tilde{G}$  has an Eulerian subgraph  $\tilde{H}$  containing  $\tilde{e}, \tilde{f}$  and with  $w(\tilde{H}) \geq (w(\tilde{G})/6)^\alpha + 2$ . For  $i = 1, 2$  let  $C_i^*$  be the 3-edge-connected graph obtained from  $G$  by contracting  $G - C_i$  to a single vertex  $z_i$ . By (3.2),  $C_1^*$  has an Eulerian subgraph  $H_1$  containing  $g_1, k_1$  and with  $w(H_1) \geq (w(C_1^*)/6)^\alpha + 2$ . Similarly,  $C_2^*$  has an Eulerian subgraph  $H_2$  containing  $g_2', k_2$  and with  $w(H_2) \geq (w(C_2^*)/6)^\alpha + 2$ . Let  $H$  be the Eulerian subgraph of  $G$  with  $E(H) = (E(\tilde{H}) - \{\tilde{e}, \tilde{f}\}) \cup E(H_1) \cup E(H_2) \cup \{e, f\}$ . Then

$$w(H) \geq w(\tilde{H}) + w(H_1) + w(H_2) \geq (w(\tilde{G})/6)^\alpha + (w(C_1^*)/6)^\alpha + 2 + (w(C_2^*)/6)^\alpha + 2.$$

Since  $w(G) \geq (w(\tilde{G}) - 2) + w(C_1^*) + w(C_2^*) + w(\{e, f\})$ , we may use Lemma 2.6 to deduce that  $w(H) \geq (w(G)/6)^\alpha + 2$ .

Hence we may assume that  $\{e, k_1\}$  is not splittable at  $z$  in  $G^*$ , and, by symmetry,  $\{e, k_2\}$  is not splittable at  $z$  in  $G^*$ . Thus  $\{e, f\}$  is the only splittable pair at  $z$  in  $G^*$  that contains  $e$ .






Figure 6:  $\{e, f\}$  is the only splittable pair at  $z$ .

We may now choose a partition  $Y_0, Y_1, Y_2, Y_3$  of  $V(G^*) - z$  satisfying the conclusions of Lemma 2.3, and with  $v \in Y_0$  and  $v' \in Y_2$ . See the first graph in Figure 6. We have  $\delta_G(Y_i) = 3$  for all  $0 \leq i \leq 3$ . Thus (3.5) implies that  $Y_0 = \{v\}$  and  $Y_2 = \{v'\}$ . We may now deduce that  $G$  has the structure illustrated in the second graph of Figure 6, and that the graph  $\tilde{G}$  obtained from  $G$  by contracting  $Y_1, Y_3, C_1, C_2$  to single vertices  $y_1, y_3, c_1, c_2$ , respectively, is isomorphic to the cube. We may construct a Hamilton cycle  $\tilde{H} = uv y_3 c_2 u' v' y_1 c_1 u$  in  $\tilde{G}$  which contains  $e, f$ . By Lemma 3.3,  $G$  has an Eulerian subgraph  $H$  such that all edges of  $\tilde{H}$  are in  $H$  and

$$\begin{aligned}
w(H) &\geq (w(C_1)/6)^\alpha + (w(C_2)/6)^\alpha + (w(Y_1)/6)^\alpha + (w(Y_3)/6)^\alpha + w(\tilde{H}) \\
&\geq ([w(C_1) + w(C_2) + w(Y_1) + w(Y_3)]/6)^\alpha + 8 \\
&\geq ([w(G) - 24]/6)^\alpha + 8 \\
&\geq (w(G)/6)^\alpha + 2
\end{aligned}$$

by Lemma 2.6. □

We may further assume that:

$$\mathcal{S} \neq \mathcal{S}'. \quad (3.12)$$

Suppose on the contrary that  $\mathcal{S} = \mathcal{S}'$ . By (3.11) and symmetry we may assume that  $C'_1 = C_1, C'_2 = D_2, D'_1 = C_2, D'_2 = D_1$  and

$$w(D_2) = \min\{w(C_1), w(C_2), w(D_1), w(D_2)\}.$$

We first consider the case when  $K$  is empty. See the first graph in Figure 7. Then (3.5) implies that  $\delta_G(C_1, C_2) = \delta_G(D_1, D_2) = \delta_G(C_2, D_1) = \delta_G(C_1, D_2) = 0$ . Hence  $\delta_G(C_1, D_1) = \delta_G(C_2, D_2) = 1$  and the graph  $\tilde{G}$  obtained from  $G$  by separately contracting each of  $C_1, C_2, D_1, D_2$  to single vertices  $c_1, c_2, d_1, d_2$ , respectively, is isomorphic to the Wagner graph. In  $\tilde{G}$  there is a cycle  $\tilde{H} = u v d_1 c_1 u' v' c_2 u$  containing  $e, f$ . By Lemma 3.3,  $G$  has an Eulerian subgraph  $H$  such that all edges of  $\tilde{H}$  are in  $H$  and

$$\begin{aligned}
w(H) &\geq (w(C_1)/6)^\alpha + (w(C_2)/6)^\alpha + (w(D_1)/6)^\alpha + w(\tilde{H}) \\
&\geq ([w(C_1) + w(C_2) + w(D_1) + w(D_2)]/6)^\alpha + 7 \\
&\geq ([w(G) - 24]/6)^\alpha + 7 \\
&\geq (w(G)/6)^\alpha + 2,
\end{aligned}$$

where the second inequality uses the minimality of  $w(D_2)$  and Corollary 2.9(b), the third inequality uses the fact that there are 12 edges in  $G$  which do not belong to  $C_1, C_2, D_1$  or  $D_2$ , and the last inequality uses Lemma 2.6.




Figure 7: The structure of  $G$  when  $\mathcal{S} = \mathcal{S}'$ .

Hence we may assume that  $K$  is not empty. The 3-edge-connectivity of  $G$  now implies that  $\delta_G(X, K) = 1$  and  $\delta_G(X, Y) = 0$  for all  $X, Y \in \mathcal{S}$ . Let  $e_1, e_2, f_1, f_2$  be the edges from  $K$  to  $C_1, C_2, D_1, D_2$ , respectively, and let  $x_1, x_2, y_1, y_2$  be the endvertices of  $e_1, e_2, f_1, f_2$  in  $K$ , respectively. See the second graph in Figure 7. Since  $G$  is 3-edge-connected,  $K$  is connected.

Suppose  $K$  has a cut edge, say  $k$ , separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$ . Let  $J_i$  denote the component of  $K - k$  containing  $\{x_i, y_i\}$ , for  $i = 1, 2$ . See Figure 8. Then  $\delta(J_i) = 3$  for  $i = 1, 2$ . Let  $\tilde{G}$  be obtained from  $G$  by separately contracting  $C_1, C_2, D_1, D_2, J_1, J_2$  to single vertices  $c_1, c_2, d_1, d_2, j_1, j_2$ , respectively. Then  $\tilde{G}$  is isomorphic to the Petersen graph and we may construct a cycle  $\tilde{H}$  in  $\tilde{G}$  which contains  $e, f$  and all vertices of  $\tilde{G}$  other than  $d_2$ . By Lemma 3.3,  $G$  has an Eulerian subgraph  $H$  such that all edges of  $\tilde{H}$  are in  $H$  and

$$\begin{aligned}
w(H) &\geq (w(C_1)/6)^\alpha + (w(C_2)/6)^\alpha + (w(D_1)/6)^\alpha + (w(J_1)/6)^\alpha + (w(J_2)/6)^\alpha + w(\tilde{H}) \\
&\geq ([w(C_1) + w(C_2) + w(D_1) + w(D_2)]/6)^\alpha + (w(J_1)/6)^\alpha + (w(J_2)/6)^\alpha + 9 \\
&\geq ([w(C_1) + w(C_2) + w(D_1) + w(D_2) + w(J_1) + w(J_2)]/6)^\alpha + 9 \\
&\geq ([w(G) - 30]/6)^\alpha + 9 \\
&\geq (w(G)/6)^\alpha + 2,
\end{aligned}$$

where the second inequality uses the minimality of  $w(D_2)$  and Corollary 2.9(b), the third and fifth inequalities use Lemma 2.6, and the fourth inequality uses the fact that there are 15 edges in  $G$  which do not belong to  $C_1, C_2, D_1, D_2, J_1$  or  $J_2$ .




Figure 8: The case when  $K$  has a cut edge separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$

Hence we may assume that  $K$  has a no cut edge separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$ . Let  $L$  be the graph obtained from  $K$  by adding two new vertices  $z_1, z_2$ , an edge  $g$  from  $z_1$  to  $z_2$ , and four other edges joining  $z_1$  to  $x_1, y_1$  and  $z_2$  to  $x_2, y_2$ . Let  $F$  be obtained from  $L$  by contracting  $g$ , and let  $K^*$  be obtained from  $L - g$  by suppressing  $z_1$  and  $z_2$  to edges  $k_1, k_2$ , respectively, of weight 1. Then  $F$  is 3-edge-connected since it can be obtained from  $G$  by contracting  $G - K$  to a single vertex. The fact that  $K$  has no cut edge separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$  now implies that  $L$  is also 3-edge-connected. We may now apply Lemma 3.4 to  $L$  to deduce that  $K^*$  has an Eulerian subgraph  $H^*$  such that  $k_1, k_2 \in E(H^*)$  and  $w(H^* - \{k_1, k_2\}) \geq (w(K)/6)^\alpha$ .

Let  $G'$  be obtained from  $G$  by contracting  $K$  to a single vertex  $z$ , and  $\tilde{G}'$  be obtained from  $G'$  by contracting  $C_1, C_2, D_1, D_2$  to single vertices  $c_1, c_2, d_1, d_2$ , respectively. Then  $\tilde{H}' := uvd_1zd_2u'v'c_2zc_1u$  is an Eulerian subgraph of  $\tilde{G}'$  which contains  $e, f$ . By Lemma 3.3, there is an Eulerian subgraph  $H'$  of  $G'$  such that all edges of  $\tilde{H}'$  are contained in  $H'$  and

$$\begin{aligned} w(H') &\geq (w(C_1)/6)^\alpha + (w(C_2)/6)^\alpha + (w(D_1)/6)^\alpha + (w(D_2)/6)^\alpha + w(\tilde{H}') \\ &\geq ([w(C_1) + w(C_2) + w(D_1) + w(D_2)]/6)^\alpha + 10 \\ &\geq ([w(G) - w(K) - 28]/6)^\alpha + 10 \end{aligned}$$

where the second inequality uses Lemma 2.6, and last inequality uses the fact that there are 14 edges in  $G$  which do not belong to  $C_1, C_2, D_1, D_2$ , or  $K$ .

The facts that  $k_1, k_2 \in E(H^*)$  and that  $E(\tilde{H}') \subseteq E(H')$  imply that  $E(H^* - \{k_1, k_2\}) \cup E(H')$  induces an Eulerian subgraph  $H$  of  $G$  with  $e, f \in H$  and

$$w(H) \geq w(H^* - \{k_1, k_2\}) + w(H') \geq (w(K)/6)^\alpha + ([w(G) - w(K) - 28]/6)^\alpha + 10 \geq (w(G)/6)^\alpha + 2$$

by Lemma 2.6.  $\square$

Let  $\mathcal{S} \cup \mathcal{S}' = \{X_1, X_2, \dots, X_q\}$  where  $w(X_1) \geq w(X_2) \geq \dots \geq w(X_q)$ . By (3.12),  $q \geq 5$ . Relabeling if necessary we may suppose that  $X_q = C_1$ . Let  $r = 4^{1/\alpha} - q$ . We may assume that:

$$w(K) \leq rw(C_1). \tag{3.13}$$

Suppose on the contrary that  $w(K) \geq rw(C_1)$ . Since

$$w(G) \geq \sum_{i=1}^q w(X_i) + w(K) + w(e, f, g_1, g_2, h_1, h_2, g'_1, g'_2, h'_1, h'_2)$$

we have

$$w(G) - w(C_1) - w(C_2) - 10 \geq (q - 2 + r)w(C_1) = (4^{1/\alpha} - 2)w(C_1) = 3^{1/\alpha}w(C_1).$$

Recall that, for  $i = 1, 2$ ,  $S_i$  is the 2-edge-cut which separates  $C_i$  from  $e$  in  $G - g_i$ . See Figure 4. Let  $S_1 = \{e_1, e_2\}$ ,  $V(e_i) = \{x_i, y_i\}$  with  $x_i \in C_1$  and  $y_i \notin C_1$ . Similarly let  $S_2 = \{l_1, l_2\}$ ,  $V(l_i) = \{a_i, b_i\}$  with  $a_i \in C_2$  and  $b_i \notin C_2$ . Let  $G'$  be obtained from  $G$  by deleting  $C_1$ , suppressing  $u$  to  $e'$ , and adding an edge  $g$  with  $V(g) = \{y_1, y_2\}$  (which may be a loop). Assign weight 1 to both  $e'$  and  $g$  in  $G'$ . Recall that the maximality of  $C_1$  implies that  $G'$  is 3-edge-connected.

Let  $G''$  be obtained from  $G'$  by contracting  $C_2$  to a vertex  $c_2$  and assign weight one to  $e', l_1, l_2$  in  $G''$ . Since  $G'$  is 3-edge-connected,  $G''$  is 3-edge-connected. We also have  $f \in E(G'')$  by (3.5) and (3.6). By (3.2),  $G''$  has an Eulerian subgraph  $H''$  such that  $e', f \in E(H'')$  and  $w(H'') \geq (w(G'')/6)^\alpha + 2 \geq ([w(G) - w(C_1) - w(C_2) - 10]/6)^\alpha + 2$ .

Without loss of generality, we may assume  $l_1 \in H''$ . Let  $C_2^*$  be the 3-edge-connected graph obtained from  $G$  by contracting  $G - C_2$  to the single vertex  $z$ . Assign weight 1 to  $g_2, l_1, l_2$  in  $C_2^*$ . Recall that  $E(C_2) \neq \emptyset$ , and hence  $C_2^* \neq K_2^3$ . So by (3.2),  $C_2^*$  contains an Eulerian subgraph  $H'$  such that  $g_2, l_1 \in E(H')$  and  $w(H') \geq (w(C_2^*)/6)^\alpha + 2 \geq ([w(C_2) + 3]/6)^\alpha + 2$ .

Let  $J = (H'' - c_2) \cup (H' - z) \cup \{l_1, e'\}$ . Then  $J \subseteq G'$  and  $w(J) \geq w(H') + w(H'') - 2$ . Let  $H = (J - e') \cup \{u, e, g_2\}$  if  $g \notin E(J)$ , and otherwise let  $H$  be the Eulerian subgraph of  $G$  obtained from  $(J - e') \cup \{u, e, g_2\}$  by replacing  $g$  by a path  $P$  between  $y_1$  and  $y_2$  and with  $E(P) \subseteq E(C_1) \cup S_1$ . Then  $e, f \in E(H)$  and  $w(H) \geq w(J) + 1 \geq w(H') + w(H'') - 1$ . Now Corollary 2.9(a) and the facts that  $w(C_2) \geq w(C_1)$ , and  $w(G) - w(C_1) - w(C_2) - 10 \geq 3^{1/\alpha}w(C_1)$ , give:

$$\begin{aligned} w(H) &\geq w(H'') + w(H') - 1 \\ &\geq ([w(G) - w(C_1) - w(C_2) - 10]/6)^\alpha + ((w(C_2) + 3)/6)^\alpha + 3 \\ &\geq (w(G)/6)^\alpha + 2. \end{aligned}$$

□

We can now complete the proof of the theorem. Note that  $4^{1/\alpha} < 7$  so the fact that  $0 \leq w(K) \leq (4^{1/\alpha} - q)w(C_1)$  by (3.13) implies that  $q \leq 6$ . For all  $1 \leq i \leq 4$  we have

$$\sum_{j=5}^q w(X_j) + w(K) \leq (q - 4 + r)w(X_i) = (4^{1/\alpha} - 4)w(X_i) \quad (3.14)$$

by (3.13). Choose  $x_i \in V(X_i)$  for  $1 \leq i \leq 4$ .

Suppose that no Eulerian subgraph of  $G$  contains  $\{x_1, x_2, x_3, x_4, e, f\}$ . Then, by Lemma 2.5,  $G$  has the structure depicted in Figure 3(a) or (b). Since all 3-edge-cuts which contain  $e$  or  $f$  are trivial by (3.6), we have  $|Z_i| = 1$  for  $5 \leq i \leq 8$ . We may now deduce that  $\mathcal{S} = \{Z_1, Z_2, Z_3, Z_4\} = \mathcal{S}'$ , which contradicts the fact that  $\mathcal{S} \neq \mathcal{S}'$  by (3.12).

Thus  $\{x_1, x_2, x_3, x_4, e, f\}$  is contained in an Eulerian subgraph  $H'$  of  $G$ . Let  $\tilde{G}$  be the graph obtained from  $G$  by contracting  $X_i$  to the single vertex  $x_i$  for  $1 \leq i \leq 4$ . We may obtain an Eulerian subgraph  $\tilde{H}$  of  $\tilde{G}$  which contains  $\{x_1, x_2, x_3, x_4, e, f\}$  from  $H'$  by contracting the edges which belong to  $Z_i$  for all  $1 \leq i \leq 4$ . By Lemma 3.3, there is an Eulerian subgraph  $H$  of  $G$  such that all edges of  $\tilde{H}$  are contained in  $H$  and

$$\begin{aligned}
w(H) &\geq (w(X_1)/6)^\alpha + (w(X_2)/6)^\alpha + (w(X_3)/6)^\alpha + (w(X_4)/6)^\alpha + w(\tilde{H}) \\
&\geq \left( \left[ w(X_1) + w(X_2) + w(X_3) + w(X_4) + \sum_{j=5}^q w(X_j) + w(K) \right] / 6 \right)^\alpha + 8 \\
&\geq ([w(G) - 40]/6)^\alpha + 8 \\
&\geq ([w(G) - 40]/6 + 6^{1/\alpha})^\alpha + 2 \\
&\geq (w(G)/6)^\alpha + 2
\end{aligned}$$

where the second inequality uses (3.14) and Corollary 2.9(c), the third inequality uses the fact that there are at most 20 edges of  $G$  which do not belong to  $X_1, X_2, \dots, X_q$  or  $K$ , and the fourth inequality uses Lemma 2.6.  $\blacksquare$

## 4 Corollaries

It is easy to see that Theorem 1.1 is simply the case of Theorem 3.1 when all edges have the same weight 1. Theorem 1.1 in turn has the following consequence.

**Corollary 4.1** *Let  $G$  be a 3-edge-connected graph with  $|G| \geq 2$ , and let  $e, f \in E(G)$ . Then  $G$  contains an Eulerian subgraph  $H$  such that  $e, f \in H$  and  $|H| \geq (|G|/12)^\alpha + 1$ , where  $\alpha \approx 0.753$  is the real root of  $4^{1/x} - 3^{1/x} = 2$ .*

*Proof.* Choose a counterexample  $G$  so that  $|E(G)|$  is minimum. It is easy to see that the corollary holds if  $|G| = 2$  and hence  $|G| \geq 3$ .

Suppose  $G$  has a vertex  $v$  of degree at least 5. Then by Lemma 2.1 there exists a splittable pair  $g, h$  at  $v$ . By splitting  $g, h$  at  $v$ , we arrive at a 3-edge-connected graph  $G' := G_v^{g,h}$ . Since  $d(v) \geq 5$ ,  $|G'| = |G|$ . Let  $e' = e$  if  $e \notin \{g, h\}$ ; otherwise let  $e'$  denote the edge resulted from suppressing  $v$ . Define  $f'$  analogously. By the choice of  $G$ ,  $G'$  contains an Eulerian subgraph  $H'$  such that  $e', f' \in H'$  and  $|H'| \geq (|G'|/12)^\alpha + 1$ . Then  $H'$  gives rise the desired Eulerian subgraph in  $G$ .

So we may assume  $\Delta(G) \leq 4$ . By Theorem 1.1,  $G$  contains an Eulerian subgraph  $H$  such that  $e, f \in H$  and  $|E(H)| \geq (|E(G)|/6)^\alpha + 2$ . Since  $\Delta(G) \leq 4$ ,  $\Delta(H) \leq 4$ ; and so,  $|E(H)| \leq 2|H|$ . Hence  $|H| \geq |E(H)|/2 \geq (|G|/12)^\alpha + 1$ , a contradiction.  $\blacksquare$

Theorem 1.2 follows directly from the next result.

**Theorem 4.2** *Let  $G$  be a 3-connected claw-free graph and let  $x, y \in V(G)$ . Then  $G$  contains a cycle  $C$  such that  $x, y \in C$  and  $|C| \geq (|G|/6)^\alpha + 2$ , where  $\alpha \approx 0.753$  is the real root of  $4^{1/x} - 3^{1/x} = 2$ .*

*Proof.* Choose a counterexample  $G, x, y$  so that  $|G|$  is minimum and, subject to this condition,  $|E(G)|$  is maximum.

We claim that  $G$  is the line graph of a simple graph  $G_1$ . Let  $G^*$  denote the Ryjáček closure of  $G$ . Suppose  $G^* \neq G$ . Then  $|E(G^*)| > |E(G)|$  so, by the choice of  $G$ ,  $G^*$  has a cycle  $C^*$  such that  $x, y \in V(C^*)$  and  $|C^*| \geq (|G|/6)^\alpha + 2$ . Then by Theorem 1.3,  $G$  has a cycle  $C$  such that  $V(C^*) \subseteq V(C)$ , a contradiction. So  $G = G^*$ , and the claim follows.

Since  $G$  is 3-connected, for each edge-cut  $S$  in  $G_1$  of size at most 2,  $G - S$  has exactly two components, one of which is trivial. Let  $U = \{v \in V(G_1) : d_{G_1}(v) \geq 3\}$ . Then  $U \neq \emptyset$ , and for any  $v \in V(G_1) - U$ , all neighbors of  $v$  are contained in  $G_1$  and the edges at  $v$  form a 1-edge-cut or 2-edge-cut in  $G_1$ .

Let  $G_2$  and  $w : E(G_2) \rightarrow \{1, 2\}$  be defined as follows. For each 1-edge-cut  $uv$  of  $G_1$  with  $u \in U$  (hence  $d_{G_1}(v) = 1$ ), we delete  $v$  and add a loop at  $u$ . For each 2-edge-cut  $\{ab, bc\}$  of  $G_1$  (hence  $d_{G_1}(b) = 2$ ), we delete  $b$  and add an edge between  $a$  and  $c$  with weight 2. The loops and all other edges in  $G_1[U]$  have weight 1. Then  $G_2$  is 3-edge-connected and  $w(G_2) = |G|$ .

Since  $x, y \in V(G)$  we have  $x, y \in E(G_1)$ . Let  $x' = x$  if  $x \in E(G_2)$ ; otherwise, let  $x'$  denote the edge of  $G_2$  used to replace  $x$ . Define  $y'$  analogously. By Theorem 3.1,  $G_2$  contains an Eulerian subgraph  $H_2$  such that  $x', y' \in H_2$  and  $w(H_2) \geq (w(G_2)/6)^\alpha + 2$ . Then  $H_2$  gives rise to a cycle  $C$  in  $G$  such that  $x, y \in C$  and  $|C| \geq (w(G_2)/6)^\alpha + 2 = (|G|/6)^\alpha + 2$ . ■

## 5 Algorithmic considerations

There is a large gap between best known polynomial algorithms for approximating the longest cycle in a graph and hardness results. The best known polynomial time approximation algorithm, due to Gabow [24], finds a cycle of length at least  $\exp(\Omega(\sqrt{\log c(G)}/\log \log c(G)))$  in any graph  $G$  (which gives a polynomial algorithm for constructing a cycle of length at least  $\min\{\log c(G)^t, c(G)\}$  for any fixed  $t$ ). Alon, Yuster and Zwick [1], give a polynomial algorithm for constructing a cycle of length at least  $\min\{\log |G|, c(G)\}$ . On the other hand, Karger, Motwani, and Ramkumar [33], show that it is NP-hard to find a path of length at least  $r\ell(G)$  for any fixed  $r > 0$ , where  $\ell(G)$  demotes the length of a longest path in  $G$ . Better approximation algorithms are known for graphs of bounded degree, see [20, 21]. In [21], Feder, Motwani and Subi give a polynomial time algorithm for finding a cycle of length at least  $c(G)^{(\log_3 2)/2} > c(G)^{0.315}$  in any graph of maximum degree three. Their algorithm is based on a polynomial-time algorithm for constructing a cycle of weight at least  $w(G)^{\log_3 2}$  in any 3-connected cubic graph  $G$  equipped with nonnegative edge-weights. On the other hand Bazgan, Santha, and Tuza [4], show that, for any fixed  $r > 0$ , it is NP-hard to find a path of length at least  $r|G|$  in a cubic Hamiltonian graph  $G$ .

The situation seems to be just as unclear for exact algorithms. Algorithms for solving the Travelling Salesman Problem, [6, 27, 34, 35], can be used to find a Hamilton cycle in a graph  $G$ , or deduce that no such cycle exists, in  $O^*(2^{|G|})$  time. (The  $O^*$ -notation means that factors which are polynomial in  $|G|$  are suppressed.) The time complexity can be improved to  $O^*(b^{|G|})$ , for various constants  $b$  with  $1 < b < 2$ , when  $G$  has bounded maximum degree, see [5, 19, 28]. It is conceivable that these algorithms could be modified to give similar results for constructing longest cycles but the only specific results we know of are an algorithm of Monien [37], subsequently improved by Bodlaender [7] to find a longest cycle in an arbitrary graph  $G$  in time  $O(c(G)! 2^{c(G)} |G|)$ , and a recent result of Broersma et al [10] which gives an  $O^*(1.8878^{|G|})$  algorithm for finding a longest cycle when  $G$  is claw-free.

We indicate in Subsection 5.1 below how our proof of Theorem 3.1 can be adapted to give a polynomial time algorithm for finding an Eulerian subgraph  $H$  in a  $\{1, 2\}$ -edge-weighted,

3-edge-connected graph  $G$  such that  $w(H) \geq (w(G)/6)^\alpha + 2$ . In particular, this finds a cycle of length at least  $(|G|/4)^\alpha > (|G|/4)^{0.753}$  in any 3-connected cubic graph  $G$ . Our algorithm uses a subroutine which finds an Eulerian subgraph containing two given edges and four given vertices in a 3-edge-connected graph (when such a subgraph exists). This will be described in Subsection 5.2. We then use the algorithm from Subsection 5.1 to obtain a polynomial algorithm for finding a cycle of length at least  $(|G|/6)^\alpha$  in any 3-connected claw-free graph  $G$  in Subsection 5.3.

## 5.1 Large Eulerian subgraphs containing two given edges

Recall the proof of Theorem 3.1. Let  $G$  be a 3-edge-connected graph,  $e, f \in E(G)$ , and  $w : E(G) \rightarrow \{1, 2\}$ . We outline an algorithm for finding an Eulerian subgraph  $H$  in  $G$  such that  $e, f \in H$  and  $w(H) \geq (w(G)/6)^\alpha + 2$ . For convenience, we write  $(G, e, f)$  for the input, with the understanding that edges are assigned weights 1 or 2. We will use the fact that, given a graph  $G$ , two disjoint subsets  $X, Y \subseteq V(G)$ , and a fixed integer  $k$ , we can use maxflow computations to find either  $k$  edge-disjoint paths joining  $X$  to  $Y$ , or a minimal set  $X' \subseteq V(G)$  with  $X \subseteq X'$ ,  $Y \subseteq V(G) \setminus X'$  and  $\delta(X') < k$ , in  $O(|E(G)|)$  time.

### Algorithm EULERIANSUBGRAPH

INPUT: A 3-edge-connected graph  $G$ ,  $e, f \in E(G)$ , and  $w : E(G) \rightarrow \{1, 2\}$ .

OUTPUT: An Eulerian subgraph  $H$  of  $G$  such that  $e, f \in H$  and  $w(H) \geq (w(G)/6)^\alpha + 2$ .

COMPLEXITY:  $f(|E(G)|) = O(|E(G)|)^3$ .

- Step 1. Check if  $e$  or  $f$  belongs to a splittable pair. If not, go to Step 2. If yes, we apply the argument in (3.3) to reduce the problem to  $(G', e', f')$ , with  $|E(G')| \leq |E(G)| - 1$ . This shows  $f(|E(G)|) \leq f(|E(G)| - 1) + O(|E(G)|^2)$ , as it takes  $O(|E(G)|)$  time to check whether a particular splitting preserves 3-edge-connectivity and there are  $O(|E(G)|)$  splittings to check.
- Step 2. Check if  $e$  and  $f$  are adjacent. If not, go to Step 3. If yes, then by the argument in (3.4) we reduce the problem to  $(G', e', f')$  with  $|E(G')| \leq |E(G)| - 1$  (when  $G'$  is 3-edge-connected), or  $(G'_1, e', f')$  and  $(G'_2, g_1, g_2)$  (when  $G'$  is not 3-edge-connected, with  $|E(G'_1)| + |E(G'_2)| = |E(G)|$ ). Note that  $G', G'_1, G'_2$  can be found in  $O(|E(G)|)$  time. So  $f(|E(G)|) \leq f(|E(G')| - 1) + O(|E(G)|)$  or  $f(|E(G)|) \leq f(|E(G'_1)|) + f(|E(G'_2)|) + O(|E(G)|)$ .
- Step 3. Check to see if  $e$  or  $f$  is contained in a non-trivial 3-edge-cut of  $G$ . If not, go to Step 4. If yes, we use the argument for (3.5) to reduce the problem to  $(G'_1, e, f)$  and  $(G'_2, e, g_1)$ , with  $|E(G'_1)| + |E(G'_2)| = |E(G)| + 3$  and  $|E(G'_i)| < |E(G)|$ . Note that  $G'_1, G'_2$  can be found in  $O(|E(G)|)$  time, so  $f(|E(G)|) \leq f(|E(G'_1)|) + f(|E(G'_2)|) + O(|E(G)|)$ .
- Step 4. Check if there is a 3-edge-cut  $S$  such that  $e$  and  $f$  are contained in different components of  $G - S$ . If not, go to Step 5. If yes, we use the argument for (3.6) to reduce the problem to  $(G'_1, e, g_1)$ ,  $(G'_1, e, g_3)$  and  $(G'_2, g_1, f)$ , with  $|E(G'_1)| \leq |E(G'_2)| < |E(G)|$  and  $|E(G'_1)| + |E(G'_2)| = |E(G)| + 3$ . This implies  $f(|E(G)|) \leq 2f(|E(G'_1)|) + f(|E(G'_2)|) + O(|E(G)|)$ . Note that the multiplicative factor of '2' in the first term on the right hand side of this inequality is compensated for by the fact that  $|E(G'_1)| \leq (|E(G)| + 3)/2$ .

- Step 5. Construct  $G_i, H_i$  with respect to  $e$  as in the paragraph above (3.8). Similarly construct  $G'_i, H'_i$  with respect to  $f$ . Check if there is some  $i \in \{1, 2\}$  such that  $G_i, H_i, G'_i$  or  $H'_i$  is 3-edge-connected. If not, got to Step 6. If yes, say,  $G_1$  is 3-edge-connected, then we use the argument for (3.8) to reduce the problem to  $(G_1, e', f')$  with  $|E(G_1)| \leq |E(G)| - 1$ . This implies  $f(|E(G)|) \leq f(|E(G)| - 1) + O(|E(G)|)$ .
- Step 6. Construct  $C_i, D_i$  with respect to  $e$  as in the paragraph above (3.9). Similarly construct  $C'_i, D'_i$  with respect to  $f$ . Check to see if  $\{C_1, C_2\} = \{C'_1, C'_2\}$  or  $\{C_1, C_2\} = \{D'_1, D'_2\}$  or  $\{D_1, D_2\} = \{C'_1, C'_2\}$  or  $\{D_1, D_2\} = \{D'_1, D'_2\}$ . If not, go to Step 7. If yes, say  $\{C_1, C_2\} = \{C'_1, C'_2\}$ , apply the argument in (3.11): we either reduce the problem to  $(\tilde{G}, \tilde{e}, f)$ ,  $(C_1^*, g_1, k_1)$  and  $(C_2^*, g_2, k_2)$ ; or find a partition  $Y_0, Y_1, Y_2, Y_3$  of  $V(G^*) - \{z\}$  given by Lemma 2.3, and reduce the problem to  $G/(G - Y_1), G/(G - Y_3), G/(G - C_1)$  and  $G/(G - C_2)$  via Lemma 3.3. In the first case,  $f(|E(G)|) \leq f(|E(\tilde{G})|) + f(|E(C_1^*)|) + f(|E(C_2^*)|) + O(|E(G)|)$  with  $|E(\tilde{G})|, |E(C_1^*)|, |E(C_2^*)| < |E(G)|$  and  $|E(\tilde{G})| + |E(C_1^*)| + |E(C_2^*)| \leq |E(G)| + 6$ . In the latter case,  $f(|E(G)|) \leq f(|E(Y_1)| + 3) + f(|E(Y_3)| + 3) + f(|E(C_1)| + 3) + f(|E(C_2)| + 3) + O(|E(G)|)$  with  $|E(Y_1)| + |E(Y_3)| + |E(C_1)| + |E(C_2)| \leq |E(G)| - 8$ .
- Step 7. Check to see if  $\mathcal{S} = \mathcal{S}'$ . If not go to Step 8. If yes, we apply the argument in (3.12). We find the member of  $\mathcal{S}$  with minimum weight, say  $D_2$ . When  $K = \emptyset$ , we reduce the problem to  $G/(G - C_1), G/(G - D_1), G/(G - C_2)$  via Lemma 3.3. We have  $f(|E(G)|) \leq f(|E(C_1)| + 3) + f(|E(C_2)| + 3) + f(|E(D_1)| + 3) + O(|E(G)|)$  with  $|E(C_1)| + |E(C_2)| + |E(D_1)| \leq |E(G)| - 8$ . When  $K \neq \emptyset$ , we reduce the problem to either  $G/(G - C_1), G/(G - C_2), G/(G - D_1), J_1, J_2$  via Lemma 3.3 (if  $K$  has a cut-edge separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$ ), or to  $K^*, G/(G - C_1), G/(G - C_2), G/(G - D_1), G/(G - D_2)$  via Lemmas 3.4 and 3.3 (if  $K$  has no cut-edge separating  $\{x_1, y_1\}$  from  $\{x_2, y_2\}$ ). In the former case,  $f(|E(G)|) \leq f(|E(C_1)| + 3) + f(|E(C_2)| + 3) + f(|E(D_1)| + 3) + f(|E(J_1)| + 3) + f(|E(J_2)| + 3) + O(|E(G)|)$  with  $|E(C_1)| + |E(C_2)| + |E(D_1)| + |E(J_1)| + |E(J_2)| = |E(G)| - |E(D_2)| - 15$ . In the latter case,  $f(|E(G)|) \leq f(|E(C_1)| + 3) + f(|E(C_2)| + 3) + f(|E(D_1)| + 3) + f(|E(D_2)| + 3) + f(|E(K^*)|) + O(|E(G)|)$  with  $|E(C_1)| + |E(C_2)| + |E(D_1)| + |E(D_2)| + |E(K^*)| = |E(G)| - 12$ .
- Step 8. Check to see if  $w(K) \geq r \min_{X \in \mathcal{S} \cup \mathcal{S}'} \{w(X)\}$ , where  $r = 4^{1/\alpha} - |\mathcal{S} \cup \mathcal{S}'|$ . If not, go to Step 9. If yes, we use the argument in (3.3) to reduce the problem to  $(G'', e', f')$ ,  $(C_2^*, g_2, l_1)$ , and possibly finding an  $x_1 x_2$ -path in  $C_1$ . We have  $|E(G'')| + |E(C_2^*)| < |E(G)|$  and  $f(|E(G)|) \leq f(|E(G'')|) + f(|E(C_2^*)|) + O(|E(G)|)$ .
- Step 9. We proceed as in the last paragraph of the proof of Theorem 3.1. Choose the four heaviest subgraphs  $X_1, X_2, X_3, X_4 \in \mathcal{S} \cup \mathcal{S}'$  and let  $\tilde{G}$  be obtained from  $G$  by contracting each  $X_i$  to a single vertex  $x_i$ , for  $1 \leq i \leq 4$ . We can use the algorithm COVER, given in Subsection 5.2 below, to construct an Eulerian subgraph  $\tilde{H}$  of  $\tilde{G}$  containing  $\{e, f, x_1, x_2, x_3, x_4\}$  in time  $O(|E(\tilde{G})|^3)$ . This allows us to reduce the problem to that for  $(X'_i, e_i, f_i)$ ,  $1 \leq i \leq 4$ , where  $X'_i = G/(G - X_i)$  and  $e_i, f_i$  are the edges of  $\tilde{H}$  incident to  $x_i$ . This gives  $f(|E(G)|) \leq \sum_{i=1}^4 f(|E(X'_i)|) + O(|E(\tilde{G})|^3)$ , where  $|E(X'_i)| < |E(G)|$  for  $1 \leq i \leq 4$  and  $|E(\tilde{G})| + \sum_{i=1}^4 |E(X'_i)| = |E(G)| + 12$ .

From Steps 1–9, we see that  $f(|E(G)|) = O(|E(G)|^3)$ . So given a 3-edge-connected graph



$G$ ,  $e, f \in E(G)$ , and weight function  $w : E(G) \rightarrow \{1, 2\}$ , one can, in  $O(|E(G)|^3)$  time, find an Eulerian subgraph  $H$  such that  $e, f \in H$  and  $w(H) \geq (w(G)/6)^\alpha + 2$ .

## 5.2 Eulerian subgraphs containing a given set of four vertices and two edges

Let  $G$  be a 3-edge-connected graph. We say that  $G$  is *essentially 4-edge-connected* if all 3-edge-cuts of  $G$  are trivial. Let  $F \subseteq E(G)$  and  $X \subseteq V(G)$  with  $|F| = 2$  and  $|X| \leq 4$ . We say that  $(G, F, X)$  is *admissible* if  $G$  has an Eulerian subgraph  $H$  which contains  $F \cup X$ . We will outline an  $O(|E(G)|^3)$  algorithm which constructs such a subgraph  $H$  given an admissible triple  $(G, F, X)$ . Note that we can check whether a given triple is admissible, and construct disjoint subgraphs  $Z_1, Z_2, \dots, Z_m$  as in Lemma 2.5 if it is not, in  $O(|E(G)|^3)$ , as follows. We use maxflow computations to check if  $G$  is essentially 4-edge-connected in  $O(|E(G)|^2)$  time. If yes then it suffices to check if  $G$  is the Wagner graph or the Peterson graph with  $F, X$  as indicated in Figure 3. If not then we find a non-trivial 3-edge-cut  $S$  of  $G$  and construct the components  $G_1, G_2$  of  $G - S$ . If  $|V(G_i) \cap X| \leq 1$  and  $E(G_i) \cap F = \emptyset$  for some  $1 \leq i \leq 2$ , then we reduce the problem to  $(G', F, X')$  where  $G' = G/G_i$  and  $X'$  is the image of  $X$  under this contraction. Otherwise we deduce that  $(G, F, X)$  is admissible.

We first give a special case of the algorithm for cubic graphs. We use a result of Andersen et al [2] that an essentially 4-edge-connected cubic graph  $G$  on at least fourteen vertices has at least  $(|E(G)| + 12)/5$  *removable edges* i.e. edges  $e$  such that  $G - e$  is homeomorphic to an essentially 4-edge-connected cubic graph.

### Algorithm CUBIC COVER

INPUT: An admissible triple  $(G, F, X)$  where  $G$  is cubic.

OUTPUT: A cycle  $C$  of  $G$  such that  $F \cup X$  is contained in  $C$ .

COMPLEXITY:  $f_1(|E(G)|) = O(|E(G)|^3)$ .

- Step 1 Let  $F = \{e, f\}$ . Put  $G^+ = G$  if  $e, f$  are adjacent and otherwise let  $G^+$  be obtained from  $G$  by subdividing  $e, f$  with two new vertices  $x_5, x_6$  and adding a new edge joining them. Check to see if  $G^+$  is essentially 4-edge-connected. If yes, go to Step 3. If not, go to Step 2.
- Step 2 Construct a non-trivial 3-edge-cut  $S^+$  in  $G^+$ . Then  $S^+$  gives rise to a non-trivial 3-edge-cut  $S$  in  $G$  such that  $G - S$  has two components  $G_1, G_2$  and at least one of them, say  $G_1$ , has  $E(G_1) \cap F = \emptyset$ . Let  $G'_1 = G/G_2$  and  $G'_2 = G/G_1$ . It is not difficult to see that the problem can be reduced to two admissible triples  $(G'_1, F_1, X_1), (G'_2, F_2, X_2)$  for suitably defined sets  $F_1, F_2, X_1, X_2$ . Hence  $f_1(|E(G)|) \leq f_1(|E(G'_1)|) + f_1(|E(G'_2)|) + O(|E(G)|^2)$  where  $|E(G'_1)|, |E(G'_2)| < |E(G)|$  and  $|E(G'_1)| + |E(G'_2)| = |E(G)| + 3$ .
- Step 3 Check to see if  $|G| \geq 16$  and if  $G^+$  has a removable edge  $h$  which does not belong to  $F$  (when  $e, f$  are adjacent) and is not incident with  $X \cup \{x_5, x_6\}$  (when  $e, f$  are not adjacent). If not, go to Step 4. If yes, let  $G_1$  be the cubic graph which is homeomorphic to  $G - h$ . Then  $(G_1, F, X)$  is admissible since  $|G_1| \geq 14$  and  $G_1^+$  is essentially 4-edge-connected. We have  $|E(G')| = |E(G)| - 3$  and  $f_1(|E(G)|) \leq f_1(|E(G')|) + O(|E(G)|^2)$ .
- Step 4 By the above mentioned result of [2], we have  $|G^+| \leq 48$ , so  $|G| \leq 46$ . We can now find  $C$  by exhaustive search.

We next give an algorithm based on the proof of Lemma 2.5 which reduces the general case to that of cubic graphs.

**Algorithm REDUCE TO CUBIC**

INPUT: An admissible triple  $(G, F, X)$ .

OUTPUT: Either an Eulerian subgraph  $H$  such that  $F \cup X$  is contained in  $H$ , or an admissible triple  $(G', F, X)$  such that  $G'$  is cubic,  $G = G' / \{e_1, e_2, \dots, e_s\}$  for some  $e_1, e_2, \dots, e_s \in E(G')$  with  $s = \sum_{v \in V(G)} (d_G(v) - 3)$ .

COMPLEXITY:  $f_2(|E(G)|) = O(|E(G)|^3)$ .

Step 1 We construct a sequence of graphs  $G = G_0, G_1, \dots, G_s = G'$  recursively. Given  $G_i$  we construct  $G_{i+1}$  as follows. Find a vertex  $v_i \in V(G_i)$  of degree at least four, and edges  $f_i = v_i u_i, g_i = v_i w_i$  incident to  $v_i$  such that the graph  $G_{i+1}$  obtained from  $G - \{f_i, g_i\}$  by adding a new vertex  $z_i$  and three new edges  $e_i, f_i, g_i$  from  $z_i$  to  $v_i, u_i, w_i$  respectively, is 3-edge-connected. (The edges  $f_i, g_i$  exist by the argument given in first paragraph in the proof of Lemma 2.5, with  $G, G', v, u_1, u_2, e_1, e_2, e_3$  replaced by  $G_i, G_{i+1}, v_i, u_i, w_i, f_i, g_i, e_i$  respectively, and  $G_i = G_{i+1}/e_i$ .) Each step in this recursion takes  $O(|E(G)|^2)$  time so the whole step takes  $O(|E(G)|^3)$  time.

Step 2 Check to see if  $(G', F, X)$  is admissible. If yes output  $(G', F, X)$ . If not, go to Step 3.

Step 3 Construct pairwise disjoint subgraphs  $Z_1, Z_2, \dots, Z_m$  of  $G'$  as described in Lemma 2.5. Choose  $e_i \in \{e_1, e_2, \dots, e_s\}$  such that  $e_i \notin E(Z_j)$  for all  $1 \leq j \leq m$  and  $i$  is as large as possible. (The edge  $e_i$  exists since  $(G, F, X)$  is admissible.) Then  $T := \{e_{i+1}, \dots, e_s\} \subseteq \bigcup_{j=1}^m E(Z_j)$  so  $(G_i, F, X)$  is not admissible. We may construct pairwise disjoint subgraphs  $Z'_1, Z'_2, \dots, Z'_m$  of  $G_i$  as described in Lemma 2.5 by putting  $Z'_j = Z_j / (E(Z_j) \cap T)$ . We may now use maxflow computations for each  $Z'_j$  to construct an Eulerian subgraph  $H_i$  of  $G_i$  with  $F \cup X$  contained in  $H_i$  in time  $O(|E(G_i)|^2)$  as in the proof of Lemma 2.5. We then construct the required subgraph  $H$  from  $H_i$  by contracting any edges of  $e_1, e_2, \dots, e_{i-1}$  which belong to  $E(H_i)$ .

It is straightforward to combine these two algorithms to obtain:

**Algorithm COVER**

INPUT: An admissible triple  $(G, F, X)$ .

OUTPUT: An Eulerian subgraph  $H$  such that  $F \cup X$  is contained in  $H$ .

COMPLEXITY:  $f_3(|E(G)|) = O(|E(G)|^3)$ .

### 5.3 Long cycles in claw-free graphs

Let  $G$  be a 3-connected claw-free graph. It takes  $O(|E(G)||V(G)|)$  time to find the Ryjáček closure  $G^*$  of  $G$ . We can find a graph  $G_1$  such that  $L(G_1) = G^*$  in  $O(|E(G^*)|)$  time by a result of Roussopolos [39]. From the proof of Theorem 4.2,  $G_1$  is obtained from a 3-edge-connected graph  $G'_1$  by adding some pendant edges and by subdividing certain edges of  $G'_1$  exactly once. By assigning appropriate weights to edges of  $G'_1$  and replacing pendant edges with loops of weight 1, we arrive at a  $\{1, 2\}$ -edge-weighted 3-edge-connected graph  $G_2$  with  $w(G_2) = |E(G_1)| = |G^*| = |G|$ . Applying Algorithm EULERIANSUBGRAPH to  $G_2$ , we find an Eulerian subgraph  $H$  of  $G_2$  such that  $w(H) \geq (w(G_2)/6)^\alpha + 2 = (|G|/6)^\alpha + 2$ . Now an Euler tour of  $H$  can be transformed into a cycle in  $G$  of length at least  $w(H)$  as in [11]. The complexity of the algorithm is  $O(|E(G_2)|^3) = O(|V(G)|^3)$ .

## References

- [1] N. Alon, R. Yuster and U. Zwick, Color-coding, *J. ACM*, **42** (1995) 844–856.
- [2] L. D. Andersen, H. Fleischner and B. Jackson, Removable edges in cyclically 4-connected cubic graphs, *Graphs and Combinatorics* **4** (1988) 1–21.
- [3] D. Barnette, Trees in polyhedral graphs, *Canad. J. Math.* **18** (1966) 731–736.
- [4] C. Bazgan, M. Santha, and Z. Tuza, On the approximation of finding a(nother) Hamiltonian cycle in cubic Hamiltonian graphs, *J. Algorithms* **31** (1999) 249–268.
- [5] A. Björklund, T. Husfeldt, P. Kaski, and M. Koivisto, The travelling salesman problem in bounded degree graphs, in *35th International Colloquium on Automata, Languages and Programming (ICALP 2008)*, Lecture Notes in Computer Science 5125 (2008) 198–209.
- [6] E.T. Bax, Inclusion and exclusion algorithm for the Hamiltonian path problem, *Information Processing Letters* **47** (1993) 203–207.
- [7] H. L. Bodlaender, On linear time minor tests with depth-first search, *J. Algorithms*, **14** (1993) 1–23.
- [8] J. A. Bondy and R. C. Entringer, Longest cycles in 2-connected graphs with prescribed maximum degree *Canad. J. Math.* **32** (1980) 1325–1332.
- [9] J. A. Bondy and M. Simonovits, Longest cycles in 3-connected cubic graphs, *Canad. J. Math.* **32** (1980) 987–992.
- [10] H. Broersma, F. V. Fomin, P. van 't Hof, and D. Paulusma, Exact algorithms for finding longest cycles in claw-free graphs, *Manuscript*.
- [11] H.J. Broersma and D. Paulusma, Computing sharp 2-factors in claw-free graphs, in *33th International Symposium on Mathematical Foundations of Computer Science (MFCS 2008)* Lecture Notes in Computer Science **5162** (2008) 193–204.
- [12] H. J. Broersma, J. Van den Heuvel, H. A. Jung and H. J. Veldman, Long paths and cycles in tough graphs, *Graphs Combin.* **9** (1993) 3–17.
- [13] R. Cada, The \*-closure for graphs and claw-free graphs, *Discrete Math.* **308** (2008) 5585–5596.
- [14] R. Cada, E. Flandrin, H. Li and Z. Ryjáček, Cycles through given vertices and closures, *Discrete Math.* **276** (2004) 65–80.
- [15] G. Chen, Z. Gao, X. Yu, and W. Zang, Approximating the longest cycle problem on graphs with bounded degree, *Lecture Notes in Comput. Sci.* **3595** (2005), pp. 870–884.
- [16] G. Chen, J. Xu, and X. Yu, Circumference of graphs with bounded degree, *SIAM J. Comput.*, **33** (2004), pp. 1136–1170.
- [17] G. Chen and X. Yu, Long cycles in 3-connected graphs, *J. Combin. Theory Ser. B* **86** (2002) 80–99.
- [18] M.N. Ellingham, D.A. Holton and C.H.C. Little, A ten vertex theorem for 3-connected cubic graphs, *Combinatorica* **4** (1984) 256–273.
- [19] D. Eppstein, The traveling salesman problem for cubic graphs, *Journal of Graph Algorithms and Applications* **11** (2007) 6181.

- [20] T. Feder and R. Motwani, Finding large cycles in Hamiltonian graphs, *Discrete Applied Math.* to appear.
- [21] T. Feder, R. Motwani, and C. Subi, Approximating the longest cycle problem in sparse graphs, *SIAM J. Comput.* **31** (2002) 1596–1607.
- [22] H. Fleischner and B. Jackson, A note concerning some conjectures on cyclically 4-connected cubic graphs, *Annals of Discrete Math.* **41** (1989) 171–178.
- [23] A. Frank, On a theorem of Mader, *Discrete Math.* **101** (1992) 49–57.
- [24] H. N. Gabow, Finding paths and cycles of superpolylogarithmic length, *SIAM J. Comput.* **36** (2007) 1648–1671.
- [25] B. Grünbaum and T. Motzkin, Longest simple paths in polyhedral graphs, *J. London Math. Soc. (2)* **37** (1962) 152–160.
- [26] B. Grünbaum and H. Walther, Shortness exponents of families of graphs, *J. Combin. Theory Ser. A* **14** (1973) 364–385.
- [27] M. Held and R.M. Karp, A dynamic programming approach to sequencing problems, *Journal of SIAM* **10**, (1962) 196–210.
- [28] K. Iwama and T. Nakashima, An improved exact algorithm for cubic graph TSP, in *13th Annual International Computing and Combinatorics Conference (COCOON 2007)*, Lecture Notes in Computer Science **4598** (2007) 108–117.
- [29] B. Jackson, Longest cycles in 3-connected cubic graphs, *J. Combin. Theory Ser. B* **41** (1986) 17–26.
- [30] B. Jackson and N. C. Wormald, Longest cycles in 3-connected graphs of bounded maximum degree, in *Graphs, Matrices, and Designs*, R. S. Rees, ed., Dekker, New York, 1993, pp. 237–254.
- [31] B. Jackson and N. C. Wormald, Long cycles in 3-connected spanning subgraphs of bounded degree in 3-connected  $K_{1,d}$ -free graphs, *J. Combin. Theory Ser. B* **63** (1995) 163–169.
- [32] T. Jordán, Constrained edge-splitting problems, *SIAM J. Discrete Math.* **17** (2003) 88–102.
- [33] D. Karger, R. Motwani, and G. D. S. Ramkumar, On approximating the longest path in a graph, *Algorithmica*, **18** (1997), pp. 82–98.
- [34] R.M. Karp, Dynamic programming meets the principle of inclusion and exclusion, *Operations Research Letters* **1** (1982) 4951.
- [35] S. Kohn, A. Gottlieb, and M. Kohn, A generating function approach to the traveling salesman problem, in *Proceedings of the ACM annual conference (ACM 1977)*, ACM Press, 1977, 294–300.
- [36] M. M. Mathews and D. P. Sumner, Hamiltonian results in  $K_{1,3}$ -free graphs, *J. Graph Theory* **8** (1984) 139 – 146.
- [37] B. Monien, How to find long paths efficiently, in *Analysis and design of algorithms for combinatorial problems (Udine, 1982)*, North-Holland Math. Stud., 109, North-Holland, Amsterdam, 1985, 239254.

- [38] J. W. Moon and L. Moser, Simple paths on polyhedra, *Pacific J. Math.* **13** (1963) 629–631.
- [39] A.D. Roussopolos, A  $\max\{m, n\}$  algorithm for determining the graph  $H$  from its line graph  $G$ , *Information Processing Letters* **2** (1973) 108–112.
- [40] Z. Ryjáček, On a closure concept in claw-free graphs, *J. Combin. Theory Ser. B* **70** (1997) 217–224.
- [41] L. Sheppardson and X. Yu, Long cycles in 3-connected graphs in orientable surfaces, *J. Graph Theory* **41** (2002) 69–84.
- [42] Z. Szigeti, Edge-splittings preserving local edge-connectivity in graphs, *Discrete Applied Math.* **156** (2008) 1011–1018.
- [43] P. G. Tait, Remarks on colouring maps, *Proc. Roy. Soc. Edinburgh Ser. A* **10** (1880) 729.
- [44] C. Thomassen, Reflections on graph theory, *J. Graph Theory* **10** (1986) 309–324.
- [45] W. T. Tutte, On hamiltonian circuits, *J. London Math. Soc.* **21** (1946) 98–101.
- [46] W. T. Tutte, A theorem on planar graphs, *Trans. Amer. Math. Soc.* **82** (1956) 99–116.
- [47] H. Walther, Über die Anzahl der Knotenpunkte eines längsten Kreises in planaren, kubischen dreifach knotenzusammenhängenden Graphen, *Studia Sci. Math. Hungar.* **2** (1967) 391–398.
- [48] H. Walther, Über Extremalkreise irregulären Landkarten, *Wiss. Z. Techn. Hochsch. Ilmenau* **15** (1969) 139–142.
- [49] S. Zhan, On hamiltonian line graphs and connectivity, *Discrete Math.* **89** (1991) 89–95.